

INTERACTION SOLUTIONS AND ABUNDANT EXACT SOLUTIONS FOR THE NEW (3+1)-DIMENSIONAL GENERALIZED KADOMTSEV-PETVIASHVILI EQUATION IN FLUID MECHANICS

Jian-Guo Liu^{1,†}, Wen-Hui Zhu^{2,†} and Li Zhou¹

Abstract In this work, we present the interaction solutions and abundant exact solutions for the new (3+1)-dimensional generalized Kadomtsev-Petviashvili equation based on the Hirota's bilinear form and a direct function. The obtained interaction solutions contain the interaction between the rational function and the tanh function and the interaction between the rational function and the cos function. The dynamical properties of these resulting solutions are analyzed and shown in three-dimensional plots, corresponding contour graphs and plane figures.

Keywords Interaction solutions, bilinear form, exact solutions, dynamical properties.

MSC(2010) 35C08, 68M07, 33F10.

1. Introduction

Profound changes happening in modern natural science, nonlinear science through the mathematical science, life science, space science and earth science, becomes an important frontier field for contemporary scientific research. Solitary wave and soliton are one of the important concepts to promote the development of nonlinear science [1–5]. Soliton originated in solitary wave, it has been applied in a series of high tech fields such as nonlinear optical, magnetic flux sub-device, biology, plasma and optical fiber isolation, and many of these applications can be represented by nonlinear partial differential equation (NPDE) [6–14]. So it is necessary to study the solitary wave solution or the soliton solution for the NPDE. Various method are proposed by the researchers [15–25].

Recently, the rational function solution called lump solution as a kind of soliton solution has attracted the attention of many scholars, especially interaction solution between the rational function and other functions, such as trigonometric functions, exponential functions, hyperbolic functions, and so on [26, 27, 30–32]. In

[†]the corresponding author.

Email address: 395625298@qq.com (J. Liu), 415422402@qq.com (W. Zhu)

¹College of Computer, Jiangxi University of Traditional Chinese Medicine, Jiangxi 330004, China

²Institute of artificial intelligence, Nanchang Institute of Science and Technology, Jiangxi 330108, China

this paper, we aim to research the new $(3+1)$ -dimensional generalized Kadomtsev-Petviashvili (ngKP) equation [28].

$$u_{ty} + u_{tx} + u_{tz} - u_{zz} + 3(u_x u_y)_x + u_{xxx} = 0. \quad (1.1)$$

The multiple soliton solutions were investigated by Wazwaz in Ref. [28]. New exact periodic solitary-wave solutions were obtained in Ref. [29]. As far as we know, interaction solutions among the rational function, the tanh function and the cos function have not been found in other literatures.

The organization of this paper is as follows: Section 2 lists the Hirota's bilinear form and obtains the interaction solutions between the rational function and the tanh function. Section 3 presents the interaction solutions between the rational function and the cos function. Section 4 obtains the abundant exact solutions. The dynamical properties of these obtained solutions are analyzed and shown in some corresponding figures. Finally, the conclusions are presented.

2. Hirota's bilinear form and interaction solutions between the rational function and the tanh function

Substituting $u = 2[\ln \zeta]_x$ into Eq. (1.1) and simplifying, we have the following Hirota's bilinear form [29]

$$(D_t D_x + D_t D_y + D_t D_z + D_x^3 D_y - D_z^2) \zeta \cdot \zeta = 0. \quad (2.1)$$

This is equivalent to:

$$\begin{aligned} &(\zeta_{xxx} + \zeta_{tx} + \zeta_{ty} + \zeta_{tz} - \zeta_{zz}) \zeta - 3\zeta_{xy} \zeta_x + 3\zeta_{xy} \zeta_{xx} \\ &- \zeta_y \zeta_{xxx} - \zeta_t \zeta_x - \zeta_t \zeta_y - \zeta_t \zeta_z + \zeta_z^2 = 0. \end{aligned} \quad (2.2)$$

Considering Eq. (2.2) has the following interaction solutions between the rational function and the tanh function

$$\begin{aligned} \varrho &= \iota_1 x + \iota_2 y + \iota_3 z + \iota_4 t + \iota_5, \\ \varsigma &= \iota_6 x + \iota_7 y + \iota_8 z + \iota_9 t + \iota_{10}, \\ \zeta_1 &= \varrho^2 + \varsigma^2 + k \tanh(j_1 x + j_2 y + j_3 z + j_4 t) + \iota_{11}, \end{aligned} \quad (2.3)$$

where $\iota_i (1 \leq i \leq 11)$ and $j_i (1 \leq i \leq 4)$ are undetermined constants. Substituting Eq. (2.3) into Eq. (2.2), we have the following relational expression

$$\begin{aligned} \iota_2 = \iota_7 = j_1 = 0, \iota_8 &= \frac{\iota_3 \iota_6}{\iota_1}, \iota_4 = \frac{\iota_3^2}{\iota_1 + \iota_3}, j_2 = \frac{\iota_1 j_3}{\iota_3}, \\ j_4 &= \frac{\iota_3 j_3}{\iota_1 + \iota_3}, \iota_9 = \frac{\iota_3^2 \iota_6}{\iota_1^2 + \iota_3 \iota_1}, \end{aligned} \quad (2.4)$$

where $\iota_1 \neq 0, \iota_3 \neq 0, \iota_1 + \iota_3 \neq 0$. Therefore, we have

$$\begin{aligned} \zeta_1 &= k \tanh\left(\frac{\iota_3 j_3 t}{\iota_1 + \iota_3} + \frac{\iota_1 j_3 y}{\iota_3} + j_3 z\right) + \left(\iota_5 + \frac{\iota_3^2 t}{\iota_1 + \iota_3} + \iota_1 x + \iota_3 z\right)^2 \\ &+ \left(\iota_{10} + \frac{\iota_6 \iota_3^2 t}{\iota_1^2 + \iota_3 \iota_1} + \iota_6 x + \frac{\iota_6 \iota_3 z}{\iota_1}\right)^2 + \iota_{11}. \end{aligned} \quad (2.5)$$

Substituting Eq. (2.5) into the transformation $u = 2[\ln \xi]_x$, we derive the interaction solutions of the ngKP equation

$$\begin{aligned}
 u = & [2[2\iota_1(\iota_5 + \frac{\iota_3^2 t}{\iota_1 + \iota_3} + \iota_1 x + \iota_3 z) + 2\iota_6(\iota_{10} + \frac{\iota_6 \iota_3^2 t}{\iota_1^2 + \iota_3 \iota_1} + \iota_6 x \\
 & + \frac{\iota_6 \iota_3 z}{\iota_1})]/[\iota_{11} + k \tanh(\frac{\iota_3 j_3 t}{\iota_1 + \iota_3} + \frac{\iota_1 j_3 y}{\iota_3} + j_3 z) + (\iota_5 + \frac{\iota_3^2 t}{\iota_1 + \iota_3} \\
 & + \iota_1 x + \iota_3 z)^2 + (\iota_{10} + \frac{\iota_6 \iota_3^2 t}{\iota_1^2 + \iota_3 \iota_1} + \iota_6 x + \frac{\iota_6 \iota_3 z}{\iota_1})^2]. \quad (2.6)
 \end{aligned}$$

The dynamical properties for interaction solution between rational function and tanh function are displayed in Fig. 1. Figures 1(a), 1(b) and 1(c) list the 3d graphs in the (y, z) -plane when $x = -5, 0, 5$ respectively, Figures 1(d), 1(e) and 1(f) present the corresponding contour diagrams of Figures 1(a), 1(b) and 1(c), and Figures 1(g), 1(h) and 1(i) show the corresponding plane figures of Figures 1(a), 1(b) and 1(c) with $y = -8$.




Figure 1. Plots of the interaction solutions (2.6) for $\iota_1 = \iota_6 = 2$, $k = t = 1$, $\iota_3 = -1$, $\iota_5 = 5$, $\iota_{11} = -6$, $\iota_{10} = 3$, $j_3 = -2$, when $x = -5$ in (a) (d) (g), $x = 0$ in (b) (e) (h) and $x = 5$ in (c) (f) (i). $y = -8$ in (g) (h) and (i).

3. Interaction solutions between the rational function and the cos function

Based on the collision of rational function and trigonometric function, we set

$$\begin{aligned}\varrho &= \iota_1 x + \iota_2 y + \iota_3 z + \iota_4 t + \iota_5, \\ \varsigma &= \iota_6 x + \iota_7 y + \iota_8 z + \iota_9 t + \iota_{10}, \\ \zeta_2 &= \varrho^2 + \varsigma^2 + k \cos(j_1 x + j_2 y + j_3 z + j_4 t) + \iota_{11}.\end{aligned}\quad (3.1)$$

Substituting Eq. (3.1) into Eq. (2.2), we have the following comparison expression

$$\begin{aligned}\iota_7 &= \frac{\iota_2 \iota_6}{\iota_1}, \iota_8 = \frac{\iota_3 \iota_6}{\iota_1}, \iota_4 = \frac{\iota_3^2}{\iota_1 + \iota_2 + \iota_3}, \iota_9 = \frac{\iota_3^2 \iota_6}{\iota_1 (\iota_1 + \iota_2 + \iota_3)}, \\ j_2 &= -\frac{\iota_2 j_1}{\iota_1}, j_4 = \frac{\iota_1 j_3^2 - \iota_2 j_1^4}{\iota_1 (j_1 + j_3) - \iota_2 j_1}, k = \frac{2(\iota_1^2 + \iota_6^2)}{j_1^2}, \\ \iota_3 &= -[\iota_1 (\iota_1^2 + \iota_6^2) (-j_1) [2\iota_2^2 j_1^3 + j_3 (\iota_1^2 - \iota_2^2 - \iota_1 \iota_2 j_1^2)] - \\ &\quad \sqrt{\iota_1 \iota_2 (\iota_1^2 + \iota_6^2)^2 j_1^4 [\iota_1 (j_1 + j_3) - \iota_2 j_1]^2 [-\iota_1^2 + 3\iota_2^2 + \iota_2 \iota_1 (j_1^2 + 2)]}] \\ &\quad / [(\iota_1^2 + \iota_6^2) j_1^2 [\iota_1^2 + \iota_2^2 - \iota_2 \iota_1 (j_1^2 + 2)]],\end{aligned}\quad (3.2)$$

where $\iota_1 \neq 0, j_1 + j_3 \neq 0, \iota_1 + \iota_2 + \iota_3 \neq 0, j_1 \neq 0, \iota_1^2 + \iota_6^2 \neq 0, \iota_1 (j_1 + j_3) - \iota_2 j_1 \neq 0, \iota_1^2 + \iota_2^2 - \iota_2 \iota_1 (j_1^2 + 2) \neq 0$. Substituting Eq. (3.1) and Eq. (3.2) into the transformation $u = 2[\ln \xi]_x$, we get another interaction solutions of the ngKP equation

$$\begin{aligned}u &= 4\iota_1(\iota_5 + \frac{\iota_3^2 t}{\iota_1 + \iota_2 + \iota_3} + \iota_1 x + \iota_2 y + \iota_3 z)/\zeta_2 - 4(\iota_1^2 + \iota_6^2) \\ &\quad \times \sin(\frac{t(\iota_1 j_3^2 - \iota_2 j_1^4)}{\iota_1 (j_1 + j_3) - \iota_2 j_1} + j_1 x - \frac{\iota_2 j_1 y}{\iota_1} + j_3 z)/(j_1 \zeta_2) \\ &\quad + 4\iota_6[\iota_{10} + \frac{\iota_6 \iota_3^2 t}{\iota_1 (\iota_1 + \iota_2 + \iota_3)} + \iota_6 x + \frac{\iota_2 \iota_6 y}{\iota_1} + \frac{\iota_6 \iota_3 z}{\iota_1}]/\zeta_2.\end{aligned}\quad (3.3)$$

The dynamical properties for interaction solution between rational function and cos function are demonstrated in Fig. 2 and Fig. 3. Figures 2(a), 2(b) and 2(c) show the 3d graphs in the (y, t) -plane when $x = -3, 0, 3$ respectively, Figures 2(d), 2(e) and 2(f) list the corresponding contour diagrams of Figures 2(a), 2(b) and 2(c). Fig. 3 display the 3d graphs and corresponding contour diagrams of interaction solution (3.3) in the (x, y) -plane when $z = -35, 0, 35$ respectively.

4. Abundant exact solutions

To study the exact solutions for Eq. (1.1), a direct test function is selected as follows in Eq. (2.2)

$$\zeta = \Lambda_1 e^{\Psi_1} + e^{-\Psi_1} + \Lambda_2 \tan(\Psi_2) + \Lambda_3 \tanh(\Psi_3), \quad (4.1)$$

where $\Psi_i = \kappa_i x + \lambda_i y + \mu_i z + \nu_i t, i = 1, 2, 3, 4$ and $\kappa_i, \lambda_i, \mu_i, \nu_i$ are undetermined constants. Substituting Eq. (4.1) into Eq. (2.2) and equating all the coefficients




Figure 2. Plots of the interaction solutions (3.3) for $\iota_1 = \iota_5 = j_1 = 2$, $\iota_2 = -1$, $z = \iota_6 = 0$, $\iota_{11} = -6$, $\iota_{10} = 3$, $j_3 = -2$, when $x = -3$ in (a) (d), $x = 0$ in (b) (e) and $x = 3$ in (c) (f).




Figure 3. Plots of the interaction solutions (3.3) for $\iota_1 = \iota_5 = j_1 = 2$, $\iota_2 = -1$, $\iota_6 = 1$, $t = 20$, $\iota_{11} = -6$, $\iota_{10} = 3$, $j_3 = -2$, when $z = -35$ in (a) (d), $z = 0$ in (b) (e) and $z = 35$ in (c) (f).

of different powers of e^{Ψ_1} , $e^{-\Psi_1}$, $\tan(\Psi_2)$, $\tanh(\Psi_3)$ and constant term to zero via symbolic computation [33-39], Eq. (1.1) has the following exact solutions

Case(1)

$$\begin{aligned}\kappa_2 = \kappa_3 = 0, \lambda_2 = \frac{\mu_2^2 - \mu_2\nu_2}{\nu_2}, \lambda_3 = \frac{\mu_3^2 - \mu_3\nu_3}{\nu_3}, \mu_3 = \frac{\mu_2\nu_3}{\nu_2}, \\ \lambda_1 = -\frac{\mu_3^2(\kappa_1^3 + \nu_1)}{\nu_3^2} + \frac{\mu_3(\kappa_1^3 + 2\mu_1)}{\nu_3} - \kappa_1 - \mu_1, \nu_1 = \frac{\mu_1^2}{\kappa_1 + \mu_1}, \\ \nu_2 = \frac{\mu_2 \left(-\sqrt{\kappa_1^2 - 4\kappa_1^2\epsilon_1 + \kappa_1^3 + 2\mu_1} \right)}{2(\kappa_1 + \mu_1)},\end{aligned}\quad (4.2)$$

$$\begin{aligned}u_1 = [2 \left(\kappa_1 \Lambda_1 e^{\frac{\mu_1^2 t}{\kappa_1 + \mu_1} + \kappa_1 x + \lambda_1 y + \mu_1 z} - \kappa_1 e^{-\frac{\mu_1^2 t}{\kappa_1 + \mu_1} - \kappa_1 x - \lambda_1 y - \mu_1 z} \right)] \\ / [\Lambda_1 e^{\frac{\mu_1^2 t}{\kappa_1 + \mu_1} + \kappa_1 x + \lambda_1 y + \mu_1 z} + e^{-\frac{\mu_1^2 t}{\kappa_1 + \mu_1} - \kappa_1 x - \lambda_1 y - \mu_1 z} + \Lambda_2 \tan[\nu_2 t + \mu_2 z \\ + \mu_2 y \left(\frac{\mu_2}{\nu_2} - 1 \right)] + \Lambda_3 \tanh[\nu_3 t + \frac{\mu_2 \nu_3 y \left(\frac{\mu_2}{\nu_2} - 1 \right)}{\nu_2} + \frac{\mu_2 \nu_3 z}{\nu_2}]],\end{aligned}\quad (4.3)$$

where $\epsilon_1 = \pm 1$.

Case(2)

$$\begin{aligned}\kappa_2 = \kappa_3 = \lambda_1 = 0, \lambda_2 = \frac{\mu_2^2 - \mu_2\nu_2}{\nu_2}, \lambda_3 = \frac{\mu_3^2 - \mu_3\nu_3}{\nu_3}, \kappa_1 = 2\epsilon_2, \\ \mu_3 = \frac{\nu_3(\mu_1 + 2\epsilon_2)}{\mu_1 + 4\epsilon_2}, \nu_1 = \frac{\mu_1^2}{\kappa_1 + \mu_1}, \nu_2 = \frac{\mu_2(\mu_1 + 2\epsilon_2)}{\mu_1 + 2\epsilon_2},\end{aligned}\quad (4.4)$$

$$\begin{aligned}u_2 = [2 \left(2\epsilon_2 \Lambda_1 e^{\frac{\mu_1^2 t}{\mu_1 + 2\epsilon_2} + 2x\epsilon_2 + \mu_1 z} - 2\epsilon_2 e^{-\frac{\mu_1^2 t}{\mu_1 + 2\epsilon_2} - 2x\epsilon_2 - \mu_1 z} \right)] / [\Lambda_1 e^{\frac{\mu_1^2 t}{\mu_1 + 2\epsilon_2} + 2x\epsilon_2 + \mu_1 z} \\ + e^{-\frac{\mu_1^2 t}{\mu_1 + 2\epsilon_2} - 2x\epsilon_2 - \mu_1 z} + \Lambda_3 \tanh[\nu_3 t - \frac{2\nu_3 y \epsilon_2 (\mu_1 + 2\epsilon_2)}{(\mu_1 + 4\epsilon_2)^2} + \frac{\nu_3 z (\mu_1 + 2\epsilon_2)}{\mu_1 + 4\epsilon_2}] \\ + \Lambda_2 \tan \left(\frac{\mu_2 t (\mu_1 + 4\epsilon_2)}{\mu_1 + 2\epsilon_2} - \frac{2\mu_2 y \epsilon_2}{\mu_1 + 4\epsilon_2} + \mu_2 z \right)],\end{aligned}\quad (4.5)$$

where $\epsilon_2 = \pm 1$.

Case(3)

$$\begin{aligned}\kappa_3 = \Lambda_2 = \lambda_1 = 0, \lambda_3 = \frac{\mu_3^2 - \mu_3\nu_3}{\nu_3}, \kappa_1 = \frac{\mu_1(\mu_1 - \nu_1)}{\nu_1}, \\ \nu_3 = -[\mu_3[\mu_1\epsilon_3(\mu_1 - \nu_1)^2 \sqrt{-2\mu_1^3\nu_1 + (\mu_1^2 - 4)\nu_1^2 + \mu_1^4 + 3\mu_1^4\nu_1} \\ - 3\mu_1^3\nu_1^2 + \mu_1^2\nu_1^3 - \mu_1^5 - 2\nu_1^3]] / (2\mu_1\nu_1^2),\end{aligned}\quad (4.6)$$

$$\begin{aligned}u_3 = \frac{2\mu_1(\mu_1 - \nu_1)}{\nu_1 \left[\frac{\Lambda_3 e^{\frac{\nu_1 t + \frac{\mu_1^2 x}{\nu_1} + \mu_1(z-x)}{\tanh(\nu_3 t + \lambda_3 y + \mu_3 z) + 2}} + 1 \right]} \\ \Lambda_1 e^{\frac{\mu_1^2 t}{\nu_1} + \mu_1(z-x)} - 1\end{aligned}\quad (4.7)$$

where $\epsilon_3 = \pm 1$.

Case(4)

$$\begin{aligned}\lambda_1 = \lambda_2 = \lambda_3 = 0, \mu_2 = \frac{\mu_1 \nu_2}{\nu_1}, \mu_3 = \frac{\mu_2 \nu_3}{\nu_2}, \\ \kappa_1 = \frac{\mu_1 (\mu_1 - \nu_1)}{\nu_1}, \kappa_2 = \frac{\mu_2 (\mu_2 - \nu_2)}{\nu_2}, \kappa_3 = \frac{\mu_3 (\mu_3 - \nu_3)}{\nu_3},\end{aligned}\quad (4.8)$$

$$\begin{aligned}u_4 = & [2[\kappa_1 \Lambda_1 e^{\nu_1 t + \kappa_1 x + \mu_1 z} + \kappa_2 \Lambda_2 \sec^2 \left(\nu_2 t + \kappa_2 x + \frac{\mu_1 \nu_2 z}{\nu_1} \right) \\ & + \kappa_3 \Lambda_3 \operatorname{sech}^2 \left(\nu_3 t + \kappa_3 x + \frac{\mu_1 \nu_3 z}{\nu_1} \right) - \kappa_1 e^{-\nu_1 t - \kappa_1 x - \mu_1 z}]] / [\Lambda_1 e^{\nu_1 t + \kappa_1 x + \mu_1 z} \\ & + \Lambda_2 \tan \left(\nu_2 t + \kappa_2 x + \frac{\mu_1 \nu_2 z}{\nu_1} \right) + \Lambda_3 \tanh \left(\nu_3 t + \kappa_3 x + \frac{\mu_1 \nu_3 z}{\nu_1} \right) \\ & + e^{-\nu_1 t - \kappa_1 x - \mu_1 z}].\end{aligned}\quad (4.9)$$

Case(5)

$$\begin{aligned}\lambda_1 = \Lambda_2 = \lambda_3 = 0, \mu_3 = \frac{\mu_1 \nu_3}{\nu_1}, \\ \kappa_1 = \frac{\mu_1 (\mu_1 - \nu_1)}{\nu_1}, \kappa_3 = \frac{\mu_3 (\mu_3 - \nu_3)}{\nu_3},\end{aligned}\quad (4.10)$$

$$\begin{aligned}u_5 = & [2[\kappa_1 \Lambda_1 e^{\nu_1 t + \kappa_1 x + \mu_1 z} + \kappa_3 \Lambda_3 \operatorname{sech}^2 \left(\nu_3 t + \kappa_3 x + \frac{\mu_1 \nu_3 z}{\nu_1} \right) - \kappa_1 e^{-\nu_1 t - \kappa_1 x - \mu_1 z}]] \\ & / [\Lambda_1 e^{\nu_1 t + \kappa_1 x + \mu_1 z} + \Lambda_3 \tanh \left(\nu_3 t + \kappa_3 x + \frac{\mu_1 \nu_3 z}{\nu_1} \right) + e^{-\nu_1 t - \kappa_1 x - \mu_1 z}].\end{aligned}\quad (4.11)$$

Case(6)

$$\lambda_1 = \Lambda_3 = \lambda_2 = 0, \kappa_1 = \frac{\mu_1 (\mu_1 - \nu_1)}{\nu_1}, \kappa_2 = \frac{\mu_2 (\mu_2 - \nu_2)}{\nu_2}, \nu_2 = \frac{\mu_2 \nu_1}{\mu_1},\quad (4.12)$$

$$\begin{aligned}u_6 = & [2[\kappa_1 \Lambda_1 e^{\nu_1 t + \kappa_1 x + \mu_1 z} + \kappa_2 \Lambda_2 \sec^2 (\nu_2 t + \kappa_2 x + \mu_2 z) - \kappa_1 e^{-\nu_1 t - \kappa_1 x - \mu_1 z}]] \\ & / [\Lambda_1 e^{\nu_1 t + \kappa_1 x + \mu_1 z} + \Lambda_2 \tan (\nu_2 t + \kappa_2 x + \mu_2 z) + e^{-\nu_1 t - \kappa_1 x - \mu_1 z}].\end{aligned}\quad (4.13)$$

Case(7)

$$\Lambda_2 = \Lambda_3 = 0, \lambda_1 = \frac{\mu_1^2 - \nu_1 (\kappa_1 + \mu_1)}{4\kappa_1^3 + \nu_1},\quad (4.14)$$

$$\begin{aligned}u_7 = & [2[\kappa_1 \Lambda_1 \exp[\nu_1 t + \kappa_1 x + \frac{y[\mu_1^2 - \nu_1 (\kappa_1 + \mu_1)]}{4\kappa_1^3 + \nu_1} + \mu_1 z] - \kappa_1 \exp[-\nu_1 t \\ & - \kappa_1 x - \frac{y[\mu_1^2 - \nu_1 (\kappa_1 + \mu_1)]}{4\kappa_1^3 + \nu_1} - \mu_1 z]] / [\Lambda_1 \exp[\nu_1 t + \kappa_1 x + \frac{y[\mu_1^2 - \nu_1 (\kappa_1 + \mu_1)]}{4\kappa_1^3 + \nu_1} \\ & + \mu_1 z] + \exp[-\nu_1 t - \kappa_1 x - \frac{y[\mu_1^2 - \nu_1 (\kappa_1 + \mu_1)]}{4\kappa_1^3 + \nu_1} - \mu_1 z]].\end{aligned}\quad (4.15)$$

As an example, we study the dynamical behaviors for Eq. (4.3) by choosing different values of parameters as follows

$$\kappa_1 = \nu_3 = 2, \mu_1 = -1, \mu_2 = -3.\quad (4.16)$$




Figure 4. Solution (4.3) with $\Lambda_1 = -2$, $\Lambda_2 = 1$, $\Lambda_3 = 0$, $x = t = 0$, (a) three-dimensional graph (b) contour graph.




Figure 5. Solution (4.3) with $\Lambda_1 = 0$, $\Lambda_2 = 1$, $\Lambda_3 = 1$, $x = t = 0$, (a) three-dimensional graph (b) contour graph.




Figure 6. Solution (4.3) with $\Lambda_1 = -2$, $\Lambda_2 = 0$, $\Lambda_3 = 1$, $x = t = 0$, (a) three-dimensional graph (b) contour graph.




Figure 7. Solution (4.3) with $\Lambda_1 = -2$, $\Lambda_2 = 1$, $\Lambda_3 = 1$, $z = 0$, when $t = -5$ in (a) (d), $t = 0$ in (e) and $t = 5$ in (c) (f).




Figure 8. Solution (4.9) when $z = -10$ in (a) (d), $z = 0$ in (b) (e) and $z = 10$ in (c) (f).

Substituting Eq. (4.16) into Eq. (4.3), the dynamical behaviors for solution (4.3) are shown in Figs. 4-7.

Setting

$$\nu_1 = \nu_3 = 2, \mu_1 = -1, \nu_2 = \Lambda_2 = \Lambda_3 = 1, \Lambda_1 = -2. \quad (4.17)$$

Substituting Eq. (4.17) into Eq. (4.9), the dynamical behaviors for solution (4.9) are shown in Fig. 8.

5. Conclusion

In this paper, the interaction solutions for the ngKP equation between rational function and tanh function or cos function are presented based on the Hirota's bilinear form. Abundant exact solutions are also obtained by using a direct test function. Equations using this method need to have the Hirota's bilinear form. The dynamical properties of the obtained solutions are analyzed and shown in figures, which contain 3d plots, 2d contour plots and the plane graphs.

Acknowledgements

We would like to thank the Editors and the Referees for their timely and valuable comments.

References

- [1] F. Baronio, A. Degasperis, M. Conforti et al., *Solutions of the vector nonlinear Schrödinger equations: evidence for deterministic rogue waves*, Phys. Rev. Lett., 2012, 109, 044102.
- [2] S. Chen and W. Ma, *Lump solutions to a generalized Bogoyavlensky-Konopelchenko equation*, Front. Math. China., 2018, 13(3), 525–534.
- [3] C. Dai, Y. Wang and J. Zhang, *Analytical spatiotemporal localizations for the generalized $(3 + 1)$ -dimensional nonlinear Schrödinger equation*, Opt. Lett., 2010, 35, 1437–1439.
- [4] C. Dai, X. Zhang, Y. Fan et al., *Localized modes of the $(n + 1)$ -dimensional schrödinger equation with power-law nonlinearities in PT-symmetric potentials*, Commun. Nonlinear. Sci., 2017, 43, 239–250.
- [5] B. Ghanbari, A. Bekir and R. K. Saeed, *Oblique optical solutions of mitigating internet bottleneck with quadratic-cubic nonlinearity*, Int. J. Mod. Phys. B., 2019, 33(20), 1950224.
- [6] B. Ghanbari and N. Raza, *An analytical method for soliton solutions of perturbed Schrödinger's equation with quadratic-cubic nonlinearity*, Mod. Phys. Lett. B., 2019, 33(3), 1950018.
- [7] B. Ghanbari, M. Inc and L. Rada, *Solitary wave solutions to the tzitzeica type equations obtained by a new efficient approach*, J. Appl. Anal. Comput., 2019, 9(2), 568–589.

- [8] J. Gao, L. Han and Y. Huang, *Solitary Waves for the Generalized Nonautonomous Dual-power Nonlinear Schrödinger Equations with Variable Coefficients*, Journal of Nonlinear Modeling and Analysis, 2019, 1, 251–260.
- [9] B. Ghanbari and M. Inc, *A new generalized exponential rational function method to find exact special solutions for the resonance nonlinear Schrödinger equation*, Eur. Phys. J. Plus., 2018, 133, 142.
- [10] B. Ghanbari, M. S. Osman and D. Baleanu, *Generalized exponential rational function method for extended Zakharov-Kuznetsov equation with conformable derivative*, Mod. Phys. Lett. A., 2019, 34(20), 1950155.
- [11] B. Ghanbari and J. F. Gómez-Aguilar, *New exact optical soliton solutions for nonlinear Schrödinger equation with second-order spatio-temporal dispersion involving M -derivative*, Mod. Phys. Lett. B., 2019, 33(20), 1950235.
- [12] B. Ghanbari, *Abundant soliton solutions for the Hirota-Maccari equation via the generalized exponential rational function method*, Mod. Phys. Lett. B., 2019, 33(9), 1950106.
- [13] B. Ghanbari and C. Kuo, *New exact wave solutions of the variable-coefficient $(1+1)$ -dimensional Benjamin-Bona-Mahony and $(2+1)$ -dimensional asymmetric Nizhnik-Novikov-Veselov equations via the generalized exponential rational function method*, Eur. Phys. J. Plus., 2019, 134, 334.
- [14] B. Ghanbari, A. Yusuf, M. Inc et al., *The new exact solitary wave solutions and stability analysis for the $(2+1)$ -dimensional zakharov-kuznetsov equation*, Adv. Differ. Equ., 2019, 49, 1–15.
- [15] B. Ghanbari, M. Inc, A. Yusuf et al., *Exact optical solitons of Radhakrishnan-Kundu-Lakshmanan equation with Kerr law nonlinearity*, Mod. Phys. Lett. B., 2019, 33(6), 1950061.
- [16] L. Huang and Y. Chen, *Lump solutions and interaction phenomenon for $(2+1)$ -dimensional sawada-kotera equation*, Commun. Theor. Phys., 2017, 67(5), 473–478.
- [17] Y. Kong, L. Xin, Q. Qiu et al., *Exact periodic wave solutions for the modified Zakharov equations with a quantum correction*, Appl. Math. Lett., 2019, 94, 140–148.
- [18] C. Kuo and B. Ghanbari, *Resonant multi-soliton solutions to new $(3+1)$ -dimensional Jimbo-Miwa equations by applying the linear superposition principle*, Nonlinear Dyn., 2019, 96(1), 459–464.
- [19] J. Liu and Y. He, *New periodic solitary wave solutions for the $(3+1)$ -dimensional generalized shallow water equation*, Nonlinear Dyn., 2017, 90(1), 363–369.
- [20] J. Liu, Y. Tian and Z. Zeng, *New exact periodic solitary-wave solutions for the new $(3+1)$ -dimensional generalized Kadomtsev-Petviashvili equation in multi-temperature electron plasmas*, AIP. Adv., 2017, 7, 105013.
- [21] J. Li and Y. Zhou, *Exact Solutions in Invariant Manifolds of Some Higher-Order Models Describing Nonlinear Waves*, Qual. Theor. Dyn. Syst., 2019, 18(1), 183–199.
- [22] J. Li, G. Chen and S. Deng, *Smooth Exact Traveling Wave Solutions Determined by Singular Nonlinear Traveling Wave Systems: Two Models*, Int. Bifurcat. Chaos., 2019, 29(4), 1950047.

- [23] J. Li and G. Chen, *More on Bifurcations and Dynamics of Traveling Wave Solutions for a Higher-Order Shallow Water Wave Equation*, Internat. J. Bifur. Chaos Appl. Sci. Engrg., 2019, 29(1), 1950014.
- [24] Z. Lan, *Periodic, breather and rogue wave solutions for a generalized $(3 + 1)$ -dimensional variable-coefficient B-type Kadomtsev-Petviashvili equation in fluid dynamics*, Appl. Math. Lett., 2019, 94, 126–132.
- [25] Z. Lan, W. Hu and B. Guo, *General propagation lattice Boltzmann model for a variable-coefficient compound KdV-Burgers equation*, Appl. Math. Model., 2019, 73, 695–714.
- [26] Z. Lan, *Dark solitonic interactions for the $(3+1)$ -dimensional coupled nonlinear Schrödinger equations in nonlinear optical fibers*, Opt. Laser Technol., 2019, 113, 462–466.
- [27] Z. Lan and J. Su, *Solitary and rogue waves with controllable backgrounds for the non-autonomous generalized AB system*, Nonlinear Dyn., 2019, 96, 2535–2546.
- [28] Z. Lan and B. Gao, *Lax pair, infinitely many conservation laws and solitons for a $(2 + 1)$ -dimensional Heisenberg ferromagnetic spin chain equation with time-dependent coefficients*, Appl. Math. Lett., 2018, 79, 6–12.
- [29] Z. Lan, *Rogue wave solutions for a coupled nonlinear Schrödinger equation in the birefringent optical fiber*, Appl. Math. Lett., 2019, 98, 128–134.
- [30] Z. Lan, *Multi-soliton solutions for a $(2 + 1)$ -dimensional variable-coefficient nonlinear Schrödinger equation*, Appl. Math. Lett., 2018, 86, 243–248.
- [31] Y. Li and J. Liu, *Multiple periodic-soliton solutions of the $(3 + 1)$ -dimensional generalised shallow water equation*, Pramana., 2018, 90, 71.
- [32] Y. Li and J. Liu, *New periodic solitary wave solutions for the new $(2 + 1)$ -dimensional Korteweg-de Vries equation*, Nonlinear Dyn., 2018, 91(1), 497–504.
- [33] W. Ma and Y. Zhou, *Lump solutions to nonlinear partial differential equations via Hirota bilinear forms*, J. Differ. Equations., 2018, 264, 2633–2659.
- [34] W. Ma, *Riemann-Hilbert problems and N -soliton solutions for a coupled mKdV system*, J. Geom. Phys., 2018, 132, 45–54.
- [35] W. Ma and A. Abdeljabbar, *A bilinear bäcklund transformation of a $(3 + 1)$ -dimensional generalized KP equation*, Appl. Math. Lett., 2012, 25(10), 1500–1504.
- [36] M. S. Osman, B. Ghanbari and J. A. T. Machado, *New complex waves in nonlinear optics based on the complex Ginzburg-Landau equation with Kerr law nonlinearity*, Eur. Phys. J. Plus., 2019, 134, 20.
- [37] H. M. Srivastava, H. Günerhan and B. Ghanbari, *Exact traveling wave solutions for resonance nonlinear Schrödinger equation with intermodal dispersions and the Kerr law nonlinearity*, Math. Method. Appl. Sci., 2019, <https://doi.org/10.1002/mma.5827>.
- [38] A. M. Wazwaz and S. A. El-Tantawy, *A new $(3 + 1)$ -dimensional generalized kadomtsev-petviashvili equation*, Nonlinear Dyn., 2016, 84(2), 1107–1112.
- [39] G. Xu and A. M. Wazwaz, *Characteristics of integrability, bidirectional solitons and localized solutions for a $(3 + 1)$ -dimensional generalized breaking soliton equation*, Nonlinear Dyn., 2019, 96, 1989–2000.