DYNAMICAL BEHAVIOR OF THE GENERALIZED COMPLEX LORENZ CHAOTIC SYSTEM*

Fuchen Zhang^{1*#}, Fei Xu²

- 1. Chongqing Key Laboratory of Social Economy and Applied Statistics, School of Mathematics and Statistics, Chongqing Technology and Business University, Chongqing 400067, China;
- 2. Department of Mathematics, Wilfrid Laurier University, Waterloo, Ontario, Canada

Abstract

The purpose of this paper is to investigate the boundedness and global attractivity of the complex Lorenz system:

$$\dot{x} = \alpha (y-x), \quad \dot{y} = \gamma x - cy - dxz, \quad \dot{z} = -\beta z + \frac{1}{2} (\overline{x}y + x\overline{y}),$$

where $\alpha, \beta, \gamma, c, d$ are real parameters, x and y are complex variables, z is a real

variable, an overbar denotes complex conjugate variable and dots represent derivatives with respect to time. This system arises in many important applications in laser physics and rotating fluids dynamics. It is very interesting that we find that this system exhibits chaos phenomenon for the given parameters. Using generalized Lyapunov-like functions, we prove the existence of the ultimate bound set and the globally exponentially attractive set in this generalized complex Lorenz system. The rate of the trajectories is also obtained. Numerical simulations show the effectiveness and correctness of the conclusions. Finally, we present an application of our results that obtained in this paper.

Key words Complex Lorenz chaotic system; chaotic attractor; Lyapunov exponent;

Lyapunov dimension; global attractivity

1. Introduction

In 1963, Edward Lorenz [1] introduced the real Lorenz chaotic system:

$$\begin{cases} \dot{x} = \alpha (y - x), \\ \dot{y} = \gamma x - y - xz, \\ \dot{z} = -\beta z + xy, \end{cases}$$
(1)

where α, β, γ are real parameters of the Lorenz system as stated in several papers [1-3]. The Lorenz system can describe the thermal convection in fluids [1-3]. The

^{*#}Corresponding author: Fuchen Zhang. E-mail:zhangfuchen1983@163.com. Fei Xu: fxu.feixu@gmail.com

Lorenz chaotic system has inspired many researchers to study new chaotic systems and chaotic phenomena [4-12]. Since then, many methods have been proposed to study chaotic behaviors of chaotic systems [13-32].

A chaotic system is a nonlinear deterministic system that displays complex and unpredictable behaviors. Since the pioneering work by A.C. Fowler et al. [33], complex chaotic systems have become an interesting field of research over the last few decades [34-40]. The complex Lorenz system is as follows:

$$\dot{x} = \alpha (y-x), \quad \dot{y} = \gamma x - y - xz, \quad \dot{z} = -\beta z + \frac{1}{2} (\overline{x}y + x\overline{y}),$$
 (2)

where x and y are complex variables, z is a real variable and α, β, γ are real parameters, an overbar denotes complex conjugate variable and dots represent derivatives with respect to time. Variables x, y, z in system (2) are related

respectively to electric field, the atomic polarization amplitudes and the population inversion in a ring laser system of two-level atoms, for more details, see [11, 33]. It is reported in the literature [34] that the complex Lorenz system (2) is often used to describe and simulate the physics of detuned lasers. The complex Lorenz model applies to the description of detuned single mode, homogeneously broadened lasers when a certain constraint on the parameters is observed [34]. The complex Lorenz system also has many important applications in physics, for example, in laser physics and rotating fluids dynamics [35-38]. Nonlinear dynamical behaviors of the complex Lorenz system, such as bifurcation, limit cycle, analytic solution, the stability of equilibrium point, synchronous behavior, geometric structure, have been studied in [33-40].

Boundedness is an important concept in the study of chaotic dynamical systems which can be applied to analyze the Lyapunov dimension of chaotic attractors [10,14], chaos control and chaos synchronization [25, 41]. The bounds of the Lorenz system were studied by Leonov et al. in [18,19]. Inspired by Leonov' idea, Liao et al. have proposed the concept of the global exponential attractive set of a chaotic system and have obtained the global exponential attractive sets of the Lorenz system [41]. It is reported in the literatures [32, 42] that how to get the bounds of the Chen system and the Lu system is considered as an open problem. Bounds of the Chen system and the Lu system have been addressed in [29, 30].

The rest of this paper is organized as follows. The new generalized complex Lorenz chaotic system is proposed in Section 2. In Section 3, we will study chaotic behaviour of the five-dimensional Lorenz system (4). In Section 4, we will study the ultimate boundedness of the five-dimensional Lorenz system (4). In Section 5, we will study global attractivity of the five-dimensional Lorenz system (4). In Section 6, we will give conclusion remarks.

2. Mathematical model

According to the complex Lorenz system (2), we propose a generalized complex Lorenz system as follows

$$\begin{cases} \dot{x} = \alpha (y - x), \\ \dot{y} = \gamma x - cy - dxz, \\ \dot{z} = -\beta z + \frac{1}{2} (\overline{x}y + x\overline{y}), \end{cases}$$
(3)

where $x = u_1 + iu_2$, $y = u_3 + iu_4$ are complex variables, $z = u_5$ is a real state variable, $\alpha, \beta, \gamma, c, d$ are real parameters, an overbar denotes complex conjugate variable, $i^2 = -1$ and dots represent derivatives with respect to time. Variables x, y, z of system (3) are related respectively to electric field, the atomic polarization amplitudes and the population inversion in a ring laser system of two-level atoms, for more details, see [11, 33]. System (3) has many important applications in laser physics and rotating fluids dynamics [33-40]. The real version of (3) is described by

$$\begin{cases} \dot{u}_{1} = \alpha \left(u_{3} - u_{1} \right), \\ \dot{u}_{2} = \alpha \left(u_{4} - u_{2} \right), \\ \dot{u}_{3} = \gamma u_{1} - c u_{3} - d u_{1} u_{5}, \\ \dot{u}_{4} = \gamma u_{2} - c u_{4} - d u_{2} u_{5}, \\ \dot{u}_{5} = u_{1} u_{3} + u_{2} u_{4} - \beta u_{5}, \end{cases}$$

$$(4)$$

where $\alpha > 0, \beta > 0, c > 0, d > 0, \gamma \in R$ are real parameters of system (4).

3. Chaos Phenomenon

When the parameters α =0.0046, β =0.0008, γ =0.03, c = 0.001,d = 0.009, we have calculated the Lyapunov exponents of system (4) as $\lambda_1 = 0.0314$, $\lambda_2 = 0.0075$, $\lambda_3 = -0.0058$, $\lambda_4 = -0.0136$, $\lambda_5 = -0.0136$ by using the algorithm [13]. The Lyapunov exponents of system (4) is shown in Fig. 1.

The Lyapunov dimension of system (4) is given by [12, 13]

$$D_L = j + \frac{\sum_{i=1}^{j} \lambda_i}{\left|\lambda_{j+1}\right|},\tag{5}$$

such that j is the largest integer that guarantees the inequality $\sum_{i=1}^{j} \lambda_i > 0$. According

to the above formula (5), the Lyapunov dimension of system (4) is calculated as

$$D_L = 4 + \frac{\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4}{|\lambda_5|} = 4.6171.$$

The Lyapunov dimension of system (4) is a fractional number which ensures the presence of a strange attractor.

Since the largest the Lyapunov exponents of system (4) is $\lambda_1 = 0.0314 > 0$ and the Lyapunov dimension of system (4) is a fractional number, so the system (4) shows chaotic behaviour for parameters $\alpha = 0.0046$, $\beta = 0.0008$, $\gamma = 0.03$, c = 0.001, d = 0.009.



Fig. 1 Lyapunov exponents of system (4).

Remark 1:

The Lyapunov exponents of the system (4) are $\lambda_1 = -6.1757$, $\lambda_2 = -6.2097$, $\lambda_3 = -6.3292$, $\lambda_4 = -7.5025$, $\lambda_5 = -7.4897$ when $\alpha = 14$, $\beta = 3.7$, $\gamma = 35$, c = 1, d = 1. Since all Lyapunov exponents of system (4) are negative, the system (4) is not chaotic when the parameters $\alpha = 14$, $\beta = 3.7$, $\gamma = 35$, c = 1, d = 1.

In the following part, we will study the boundedness and global attractivity of the five-dimensional Lorenz system (4).

4. Boundedness

In this section, we will study the boundedness of the five-dimensional Lorenz system (4). Firstly, let us introduce the following Lemma 1 and Lemma 2 that will be

used in the following section.

Lemma 1. Define

$$\Gamma_{1} = \left\{ \left(x_{1}, x_{2}, y_{1}, y_{2}, z \right) \middle| \frac{x_{1}^{2}}{a^{2}} + \frac{x_{2}^{2}}{b^{2}} + \frac{\left(z-c\right)^{2}}{c^{2}} + \frac{y_{1}^{2}}{d^{2}} + \frac{y_{2}^{2}}{e^{2}} = 1 \right\}$$

and

$$H_1(x_1, x_2, y_1, y_2, z) = x_1^2 + x_2^2 + y_1^2 + y_2^2 + (z - 2c)^2, (x_1, x_2, y_1, y_2, z) \in \Gamma_1.$$

Then, we get

$$\max_{(x_1, x_2, y_1, y_2, z) \in \Gamma_1} H_1(x_1, x_2, y_1, y_2, z) = \begin{cases} \frac{a^4}{a^2 - c^2}, \ a \ge b, a \ge d, a \ge e, a \ge \sqrt{2}c, \\ \frac{b^4}{b^2 - c^2}, \ b > a, b \ge d, b > e, b \ge \sqrt{2}c, \\ \frac{d^4}{d^2 - c^2}, \ d > a, d > b, d \ge e, d \ge \sqrt{2}c, \\ \frac{e^4}{e^2 - c^2}, \ e > a, e \ge b, e > d, e \ge \sqrt{2}c, \\ 4c^2, \qquad a < \sqrt{2}c, b < \sqrt{2}c, d < \sqrt{2}c, e < \sqrt{2}c. \end{cases}$$

Proof: It can be easily proved by the Lagrange multiplier method. **Lemma 2.** Define

$$\Gamma_{2} = \left\{ \left(x, y, z\right) \middle| \frac{x^{2}}{a^{2}} + \frac{y^{2}}{b^{2}} + \frac{\left(z - c\right)^{2}}{c^{2}} = 1 \right\},$$
(6)

and

$$H_2(x, y, z) = x^2 + y^2 + (z - 2c)^2, (x, y, z) \in \Gamma_2$$

Then, we get

$$\max_{(x,y,z)\in\Gamma_{2}}H_{2}(x,y,z) = \begin{cases} \frac{a^{4}}{a^{2}-c^{2}}, & a \ge b, a \ge \sqrt{2}c, \\ \frac{b^{4}}{b^{2}-c^{2}}, & b > a, b \ge \sqrt{2}c, \\ 4c^{2}, & a < \sqrt{2}c, b < \sqrt{2}c. \end{cases}$$

Proof: It can be easily proved by the Lagrange multiplier method.

By Lemma 1 and Lemma 2, we can get the ultimate bound and positively invariant set of the five-dimensional Lorenz system (4).

Theorem 1. For any parameters $\alpha > 0, \beta > 0, c > 0, d > 0, \gamma \in R$, the following set with

two parameters λ and m

$$\Omega_{\lambda,m} = \left\{ U \left| mu_1^2 + mu_2^2 + \lambda u_3^2 + \lambda u_4^2 + \lambda d \left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d} \right)^2 \le R_{\lambda,m}^2 \right\}$$
(7)

is the ultimate bound set and positively invariant set of the five-dimensional Lorenz system (4), where

$$U = (u_1, u_2, u_3, u_4, u_5),$$

$$R_{\lambda,m}^2 = \begin{cases} \frac{\beta^2 (\lambda \gamma + \alpha m)^2}{4\alpha d (\beta - \alpha) \lambda}, c \ge \alpha, \beta \ge 2\alpha, \\ \frac{\beta^2 (\lambda \gamma + \alpha m)^2}{4cd (\beta - c) \lambda}, \beta \ge 2c, \alpha > c, \\ \frac{(\lambda \gamma + \alpha m)^2}{\lambda d}, \beta < 2\alpha, \beta < 2c. \end{cases}$$

Proof: Construct the Lyapunov-like function

$$V_{\lambda,m}(U) = mu_1^2 + mu_2^2 + \lambda u_3^2 + \lambda u_4^2 + \lambda d \left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d} \right)^2, \qquad (8)$$

where $\forall \lambda > 0, \forall m > 0, U = (u_1, u_2, u_3, u_4, u_5)$. The derivative of $V_{\lambda,m}(U)$ along the trajectory of (4) is

$$\begin{aligned} \frac{dV_{\lambda,m}(U)}{dt} \bigg|_{(4)} &= 2mu_1 \frac{du_1}{dt} + 2mu_2 \frac{du_2}{dt} + 2\lambda u_3 \frac{du_3}{dt} + 2\lambda u_4 \frac{du_4}{dt} + 2\lambda d \left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d} \right) \frac{du_5}{dt}, \\ &= 2\alpha mu_1 \left(u_3 - u_1 \right) + 2\alpha mu_2 \left(u_4 - u_2 \right) + 2\lambda u_3 \left(\gamma u_1 - cu_3 - du_1 u_5 \right) \\ &+ 2\lambda u_4 \left(\gamma u_2 - cu_4 - du_2 u_5 \right) + 2\lambda d \left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d} \right) \left(u_1 u_3 + u_2 u_4 - \beta u_5 \right), \\ &= -2\alpha mu_1^2 - 2\alpha mu_2^2 - 2\lambda cu_3^2 - 2\lambda cu_4^2 - 2\lambda d \beta u_5^2 + 2\beta \left(\lambda \gamma + \alpha m \right) u_5, \\ &= -2\alpha mu_1^2 - 2\alpha mu_2^2 - 2\lambda cu_3^2 - 2\lambda cu_4^2 - 2\lambda d \beta \left(u_5 - \frac{\lambda \gamma + \alpha m}{2\lambda d} \right)^2 + \frac{\beta \left(\lambda \gamma + \alpha m \right)^2}{2\lambda d}. \end{aligned}$$

Let

$$\Gamma = \left\{ U \Big| \alpha m u_1^2 + \alpha m u_2^2 + \lambda c u_3^2 + \lambda c u_4^2 + \lambda d \beta \left(u_5 - \frac{\lambda \gamma + \alpha m}{2\lambda d} \right)^2 = \frac{\beta \left(\lambda \gamma + \alpha m\right)^2}{4\lambda d} \right\}, \quad (9)$$

then Γ is an ellipsoid in \mathbb{R}^5 for $\alpha > 0, \beta > 0, c > 0, d > 0, \gamma \in \mathbb{R}$. Outside Γ , $\dot{V}_{\lambda,m}(U) < 0$, while inside $\Gamma, \dot{V}_{\lambda,m}(U) > 0$. Since $V_{\lambda,m}(U)$ is a generalized positively definite and radially unbounded continuous function and Γ is a bounded close set, then the maximum value $\max_{U \in \Gamma} V_{\lambda,m}(U) = R_{\lambda,m}^2$ of the function $V_{\lambda,m}(U)$ exists. Obviously, $\left\{ U | V_{\lambda,m}(U) \le \max_{U \in \Gamma} V_{\lambda,m}(U) = R_{\lambda,m}^2 \right\}$ contains the solutions of system (4). In order to get the maximum value $\max_{U \in \Gamma} V_{\lambda,m}(U) = R_{\lambda,m}^2$, we have to solve the following optimization problem:

$$\begin{cases} \max V_{\lambda,m}(U) = \max \left\{ mu_1^2 + mu_2^2 + \lambda u_3^2 + \lambda u_4^2 + \lambda d \left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d} \right)^2 \right\}, \\ s.t.\alpha mu_1^2 + \alpha mu_2^2 + \lambda c u_3^2 + \lambda c u_4^2 + \lambda d \beta \left(u_5 - \frac{\lambda \gamma + \alpha m}{2\lambda d} \right)^2 = \frac{\beta \left(\lambda \gamma + \alpha m\right)^2}{4\lambda d}. \end{cases}$$
(10)

In order to use Lemma 1 to solve problem (10), let us take $\sqrt{m}u_1 = x_1, \sqrt{m}u_2 = x_2$,

 $\sqrt{\lambda}u_3 = y_1, \sqrt{\lambda}u_4 = y_2, \sqrt{\lambda}du_5 = z$ as new variables, then optimization problem (10) transforms into:

$$\begin{cases} \max V_{\lambda,m}(U) = \max \left\{ x_1^2 + x_2^2 + y_1^2 + y_2^2 + \left(z - \frac{\lambda \gamma + \alpha m}{\sqrt{d\lambda}}\right)^2 \right\},\\ s.t.\frac{x_1^2}{\frac{\beta(\lambda \gamma + \alpha m)^2}{4\lambda \alpha d}} + \frac{x_2^2}{\frac{\beta(\lambda \gamma + \alpha m)^2}{4\lambda c d}} + \frac{y_1^2}{\frac{\beta(\lambda \gamma + \alpha m)^2}{4\lambda c d}} + \frac{y_2^2}{\frac{\beta(\lambda \gamma + \alpha m)^2}{4\lambda c d}} + \frac{\left(z - \frac{\lambda \gamma + \alpha m}{2\sqrt{d\lambda}}\right)^2}{\frac{(\lambda \gamma + \alpha m)^2}{4\lambda d}} = 1.\end{cases}$$

We can easily get the optimal solution of the above optimization problem by Lemma 1,

$$\max_{U \in \Gamma} V_{\lambda,m} (U) = R_{\lambda,m}^{2} = \begin{cases} \frac{\beta^{2} (\lambda \gamma + \alpha m)^{2}}{4\alpha d (\beta - \alpha) \lambda}, c \geq \alpha, \beta \geq 2\alpha, \\ \frac{\beta^{2} (\lambda \gamma + \alpha m)^{2}}{4cd (\beta - c) \lambda}, \beta \geq 2c, \alpha > c, \\ \frac{(\lambda \gamma + \alpha m)^{2}}{\lambda d}, \beta < 2\alpha, \beta < 2c. \end{cases}$$

It is easy to show that (7) is the ultimate bound set and positively invariant set of system (4).

This completes the proof.

Remark 2:

i) Let us take m = 1 in Theorem 1, then we can get that

$$\Omega_{\lambda,1} = \left\{ \left(u_1, u_2, u_3, u_4, u_5\right) \middle| u_1^2 + u_2^2 + \lambda u_3^2 + \lambda u_4^2 + \lambda d \left(u_5 - \frac{\lambda \gamma + \alpha}{\lambda d}\right)^2 \le l_\lambda^2 \right\},\$$

is the ultimate bound set and positively invariant set of the complex Lorenz system (4), where

$$l_{\lambda}^{2} = \begin{cases} \frac{\beta^{2} (\lambda \gamma + \alpha)^{2}}{4\alpha d (\beta - \alpha) \lambda}, c \geq \alpha, \beta \geq 2\alpha, \\ \frac{\beta^{2} (\lambda \gamma + \alpha)^{2}}{4cd (\beta - c) \lambda}, \beta \geq 2c, \alpha > c, \\ \frac{(\lambda \gamma + \alpha)^{2}}{\lambda d}, \beta < 2\alpha, \beta < 2c. \end{cases}$$

ii) Let us take $\lambda = 1$ in Theorem 1, then we can get that

$$\Omega_{1,m} = \left\{ \left(u_1, u_2, u_3, u_4, u_5\right) \middle| mu_1^2 + mu_2^2 + u_3^2 + u_4^2 + d\left(u_5 - \frac{\gamma + \alpha m}{d}\right)^2 \le L_m^2 \right\}$$

is the ultimate bound set and positively invariant set of the complex Lorenz system (4), where

$$L_{m}^{2} = \begin{cases} \frac{\beta^{2} (\gamma + \alpha m)^{2}}{4\alpha d (\beta - \alpha)}, c \geq \alpha, \beta \geq 2\alpha, \\ \frac{\beta^{2} (\gamma + \alpha m)^{2}}{4cd (\beta - c)}, \beta \geq 2c, \alpha > c, \\ \frac{(\gamma + \alpha m)^{2}}{d}, \beta < 2\alpha, \beta < 2c. \end{cases}$$

iii) Let us take $\lambda = 1, m = 1$ in Theorem 1, then we can get that

$$\Omega_{1,1} = \left\{ \left(u_1, u_2, u_3, u_4, u_5\right) \middle| u_1^2 + u_2^2 + u_3^2 + u_4^2 + d\left(u_5 - \frac{\gamma + \alpha}{d}\right)^2 \le r^2 \right\},\$$

is the ultimate bound set and positively invariant set of the complex Lorenz system (4), where

$$r^{2} = \begin{cases} \frac{\beta^{2} (\gamma + \alpha)^{2}}{4\alpha d (\beta - \alpha)}, c \geq \alpha, \beta \geq 2\alpha, \\ \frac{\beta^{2} (\gamma + \alpha)^{2}}{4cd (\beta - c)}, \beta \geq 2c, \alpha > c, \\ \frac{(\gamma + \alpha)^{2}}{d}, \beta < 2\alpha, \beta < 2c. \end{cases}$$

Let us take $\alpha = 14, \beta = 3.7, \gamma = 35, c = 1, d = 1$, then we can obtain that

$$\Omega_{1,1} = \left\{ \left(u_1, u_2, u_3, u_4, u_5 \right) \middle| u_1^2 + u_2^2 + u_3^2 + u_4^2 + \left(u_5 - 49 \right)^2 \le 55.2^2 \right\}$$

is the ultimate bound set and positively invariant set of the complex Lorenz system (4).

Fig. 2. shows the projection of $\Omega_{1,1}$ into the (u_2, u_3, u_4) space. Fig. 3. shows the projection of $\Omega_{1,1}$ into the (u_3, u_4, u_5) space. Projection of $\Omega_{1,1}$ onto the (u_3, u_5) plane is shown in Figure 4. Projection of $\Omega_{1,1}$ onto the (u_4, u_5) plane is shown in Figure 5.



Fig. 2 Projection of $\Omega_{1,1}$ into the (u_2, u_3, u_4) space.



Fig. 3 Projection of $\Omega_{1,1}$ into the (u_3, u_4, u_5) space.







Fig. 5 Projection of $\Omega_{1,1}$ onto the (u_4, u_5) plane.

Theorem 2. For any parameters $\alpha > 0, \beta > 0, c > 0, d > 0, \gamma \in R$, the following set

$$\Lambda = \left\{ \left(u_1, u_2, u_3, u_4, u_5\right) \middle| u_3^2 + u_4^2 + d\left(u_5 - \frac{\gamma}{d}\right)^2 \le r_0^2, u_1^2 + u_2^2 \le r_0^2 \right\}$$
(11)

is the bounds for the five-dimensional Lorenz system (4), where

$$r_{0}^{2} = \begin{cases} \frac{\beta^{2}\gamma^{2}}{4cd(\beta-c)}, \beta \geq 2c, \\ \frac{\gamma^{2}}{d}, \qquad \beta < 2c. \end{cases}$$

Proof: Construct the Lyapunov-like function

$$V_1(U) = u_3^2 + u_4^2 + d\left(u_5 - \frac{\gamma}{d}\right)^2.$$

The derivative of $V_1(U)$ along the trajectory of (4) is

$$\frac{dV_1}{dt} = 2u_3 \frac{du_3}{dt} + 2u_4 \frac{du_4}{dt} + 2d\left(u_5 - \frac{\gamma}{d}\right) \frac{du_5}{dt},$$

= $2u_3 \left(\gamma u_1 - cu_3 - du_1 u_5\right) + 2u_4 \left(\gamma u_2 - cu_4 - du_2 u_5\right)$
+ $2d\left(u_5 - \frac{\gamma}{d}\right) \left(-\beta u_5 + u_1 u_3 + u_2 u_4\right)$
= $-2cu_3^2 - 2cu_4^2 - 2d\beta u_5^2 + 2\gamma\beta u_5.$

Let

$$\Gamma_3 = \left\{ \left(u_3, u_4, u_5\right) \middle| cu_3^2 + cu_4^2 + d\beta \left(u_5 - \frac{\gamma}{2d}\right)^2 = \frac{\beta\gamma^2}{4d} \right\},\$$

then Γ_3 is an ellipsoid in R^3 for $\alpha > 0, \beta > 0, c > 0, d > 0, \gamma \in R$. Outside Γ_3 , $\dot{V}_1(U) < 0$, while inside Γ_3 , $\dot{V}_1(U) > 0$. Since $V_1(U)$ is a generalized positively definite and radially unbounded continuous function and Γ_3 is a bounded close set, then the maximum value $\max_{U \in \Gamma_3} V_1(U) = r_0^2$ of the function $V_1(U)$ exists.

In order to get the maximum value $\max_{U \in \Gamma_3} V_1(U) = r_0^2$, we have to solve the following optimization problem:

$$\begin{cases} \max V_{1}(U) = \max \left\{ u_{3}^{2} + u_{4}^{2} + d \left(u_{5} - \frac{\gamma}{d} \right)^{2} \right\}, \\ s.t.cu_{3}^{2} + cu_{4}^{2} + d \beta \left(u_{5} - \frac{\gamma}{2d} \right)^{2} = \frac{\beta \gamma^{2}}{4d}. \end{cases}$$
(12)

The following optimization problem is equivalent to

$$\begin{cases} \max V_{1}(U) = \max \left\{ u_{3}^{2} + u_{4}^{2} + d\left(u_{5} - \frac{\gamma}{d}\right)^{2} \right\}, \\ s.t. \frac{u_{3}^{2}}{\frac{\beta\gamma^{2}}{4dc}} + \frac{u_{4}^{2}}{\frac{\beta\gamma^{2}}{4dc}} + \frac{d\left(u_{5} - \frac{\gamma}{2d}\right)^{2}}{\frac{\gamma^{2}}{4d}} = 1. \end{cases}$$
(13)

In order to use Lemma 2 to solve problem (13), let us take $u_3 = z_1, u_4 = z_2, \sqrt{d}u_5 = z_3$ as new variables, then optimization problem (13) transforms into:

$$\begin{cases} \max V_{1}(U) = \max \left\{ z_{1}^{2} + z_{2}^{2} + \left(z_{3} - \frac{\gamma}{\sqrt{d}} \right)^{2} \right\} \\ s.t. \frac{z_{1}^{2}}{\frac{\beta\gamma^{2}}{4dc}} + \frac{z_{2}^{2}}{\frac{\beta\gamma^{2}}{4dc}} + \frac{\left(z_{3} - \frac{\gamma}{2\sqrt{d}} \right)^{2}}{\frac{\gamma^{2}}{4d}} = 1. \end{cases}$$

We can easily get the optimal solution of the above optimization problem by Lemma 2,

$$\max_{U\in\Gamma_{3}}V_{1}(U)=r_{0}^{2}=\begin{cases}\frac{\beta^{2}\gamma^{2}}{4cd(\beta-c)}, \beta\geq 2c,\\ \frac{\gamma^{2}}{d}, \qquad \beta<2c.\end{cases}$$

Construct the Lyapunov-like function

$$V_2(U) = u_1^2 + u_2^2.$$

The derivative of $V_2(U)$ along the trajectory of (4) is

$$\begin{aligned} \frac{dV_2}{dt} &= 2u_1 \frac{du_1}{dt} + 2u_2 \frac{du_2}{dt}, \\ &= 2\alpha u_1 (u_3 - u_1) + 2\alpha u_2 (u_4 - u_2), \\ &= -2\alpha u_1^2 - 2\alpha u_2^2 + 2\alpha u_1 u_3 + 2\alpha u_2 u_4, \\ &= -2\alpha u_1^2 - 2\alpha u_2^2 + \alpha (2u_1 u_3 + 2u_2 u_4), \\ &\leq -2\alpha u_1^2 - 2\alpha u_2^2 + \alpha (u_1^2 + u_3^2 + u_2^2 + u_4^2), \\ &= -\alpha u_1^2 - \alpha u_2^2 + \alpha (u_3^2 + u_4^2), \\ &\leq -\alpha u_1^2 - \alpha u_2^2 + \alpha r_0^2, \\ &= -\alpha [V_2(U) - r_0^2]. \end{aligned}$$

Thus, we have

$$V_{2}(U(t))-r_{0}^{2} \leq [V_{2}(U(t_{0}))-r_{0}^{2}]e^{-\alpha(t-t_{0})}.$$

So,

$$\overline{\lim_{t\to+\infty}}V_2(U(t)) \leq r_0^2.$$

This completes the proof.

5. Global exponential attractive domain

Though Theorem 1 and Theorem 2 point out that the solution of the system (4) is ultimately bounded, they do not give the rate of the trajectories going from the exterior of the trapping region into the interior trapping region. The rate of the trajectories going from the exterior of the trapping region into the interior trapping region of system (4) is given in the following Theorem 3.

Theorem 3. For any $\alpha > 0, \beta > 0, c > 0, d > 0, \gamma \in R$, with

$$V_{\lambda,m}(U) = mu_1^2 + mu_2^2 + \lambda u_3^2 + \lambda u_4^2 + \lambda d \left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d} \right)^2$$
$$\eta = \min(\alpha, c, \beta) > 0, L_{\lambda,m} = \frac{\beta (\lambda \gamma + \alpha m)^2}{\lambda d \eta}.$$

When $V_{\lambda,m}(U(t)) > L_{\lambda,m}, V_{\lambda,m}(U(t_0)) > L_{\lambda,m}$, we can get an exponential inequality of system (4), given by

$$V_{\lambda,m}\left(U\left(t\right)\right) - L_{\lambda,m} \leq \left[V_{\lambda,m}\left(U\left(t_{0}\right)\right) - L_{\lambda,m}\right]e^{-\eta(t-t_{0})}$$

Hence, the set

$$\Delta_{\lambda,m} = \left\{ U \mid V_{\lambda,m} \left(U \right) \le L_{\lambda,m} \right\}$$

$$= \left\{ \left(u_1, u_2, u_3, u_4, u_5 \right) \mid mu_1^2 + mu_2^2 + \lambda u_3^2 + \lambda u_4^2 + \lambda d \left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d} \right)^2 \le \frac{\beta \left(\lambda \gamma + \alpha m\right)^2}{\lambda d \eta} \right\},$$

$$(14)$$

is the global exponential attractive set of the five-dimensional Lorenz system (4). **Proof:** Define the Lyapunov-like function

$$V_{\lambda,m}(U) = mu_1^2 + mu_2^2 + \lambda u_3^2 + \lambda u_4^2 + \lambda d \left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d} \right)^2,$$

where $\lambda > 0, m > 0, U = (u_1, u_2, u_3, u_4, u_5).$

When $V_{\lambda,m}(U(t)) > L_{\lambda,m}, V_{\lambda,m}(U(t_0)) > L_{\lambda,m}$, the derivative of $V_{\lambda,m}(U)$ along the trajectory of (4) is

$$\begin{aligned} \frac{dV_{\lambda,m}(U)}{dt} \bigg|_{(4)} &= 2mu_1 \frac{du_1}{dt} + 2mu_2 \frac{du_2}{dt} + 2\lambda u_3 \frac{du_3}{dt} + 2\lambda u_4 \frac{du_4}{dt} + 2\lambda d\left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d}\right) \frac{du_5}{dt}, \\ &= 2\alpha mu_1 \left(u_3 - u_1\right) + 2\alpha mu_2 \left(u_4 - u_2\right) + 2\lambda u_3 \left(\gamma u_1 - cu_3 - du_1 u_5\right) \\ &+ 2\lambda u_4 \left(\gamma u_2 - cu_4 - du_2 u_5\right) + 2\lambda d\left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d}\right) \left(u_1 u_3 + u_2 u_4 - \beta u_5\right), \\ &= -2\alpha mu_1^2 - 2\alpha mu_2^2 - 2\lambda cu_3^2 - 2\lambda cu_4^2 - 2\lambda d\beta u_5^2 + 2\beta \left(\lambda \gamma + \alpha m\right) u_5, \\ &\leq -\alpha mu_1^2 - \alpha mu_2^2 - \lambda cu_3^2 - \lambda cu_4^2 - \lambda d\beta \left(u_5 - \frac{\lambda \gamma + \alpha m}{\lambda d}\right)^2 + \frac{\beta \left(\lambda \gamma + \alpha m\right)^2}{\lambda d}, \\ &\leq -\eta V_{\lambda,m} \left(U\right) + \frac{\beta \left(\lambda \gamma + \alpha m\right)^2}{\lambda d} < 0. \end{aligned}$$

That is equivalent to say,

$$\frac{dV_{\lambda,m}(U)}{dt}\Big|_{(4)} \leq -\eta \left(V_{\lambda,m}(U) - \frac{\beta(\lambda\gamma + \alpha m)^2}{\lambda d\eta}\right),\tag{15}$$

From (15), we can get

$$V_{\lambda,m}(U(t)) \leq V_{\lambda,m}(U_0)e^{-\eta(t-t_0)} + \int_{t_0}^t e^{-\eta(t-\tau)} \frac{\beta(\lambda\gamma + \alpha m)^2}{\lambda d\eta} d\tau,$$

= $V_{\lambda,m}(U_0)e^{-\eta(t-t_0)} + L_{\lambda,m}(1-e^{-\eta(t-t_0)}).$

We have the following exponential inequality

$$V_{\lambda,m}\left(U\left(t\right)\right) - L_{\lambda,m} \leq \left[V_{\lambda,m}\left(U_{0}\right) - L_{\lambda,m}\right]e^{-\eta\left(t-t_{0}\right)}$$

Taking limit on both sides of the above inequality as $t \rightarrow +\infty$ results in

$$\overline{\lim_{t \to +\infty}} V_{\lambda,m} \left(U(t) \right) \leq L_{\lambda,m}.$$

Namely, the set $\Delta_{\lambda,m}$ is the global exponential attractive set of system (4).

This completes the proof.

Remark 3:

i) Let us take $\lambda = 1, m = 0$, then we can get that the following set

$$\Psi_{1,0} = \left\{ \left(u_3, u_4, u_5 \right) \mid u_3^2 + u_4^2 + \left(u_5 - \frac{\gamma}{d} \right)^2 \le \delta^2 \right\},\tag{16}$$

is the global exponential attractive set of system (4), where

$$\eta = \min(c,\beta) > 0, \delta^2 = \frac{\beta \gamma^2}{d\eta}.$$

The proved method is similar to Theorem 3.

In the following part, we will present an application of the results that obtained in this paper. We will apply above results to show that the equilibrium point O(0,0,0,0,0) of the system (4) is the globally exponentially stable when $\alpha > 0, \beta > 0, c > 0, d > 0$,

$$\gamma < 0.$$

Theorem 4. If real parameters $\alpha > 0$, $\beta > 0$, c > 0, d > 0, $\gamma < 0$, then the equilibrium point O(0,0,0,0,0) of system (4) is the globally exponentially stable.

Proof. When $\alpha > 0, \beta > 0, c > 0, d > 0, \gamma < 0$, let us choose $m = -\gamma, \lambda = \alpha$ in Theorem

3. Then, we can get $L_{\lambda,m} = L_{\alpha,-\gamma} = 0$ according to Theorem 3. And the exponential inequality in Theorem 3 becomes

$$\begin{bmatrix} -\gamma u_{1}^{2}(t) - \gamma u_{2}^{2}(t) + \alpha u_{3}^{2}(t) + \alpha u_{4}^{2}(t) + \alpha d u_{5}^{2}(t) \end{bmatrix} \leq \begin{bmatrix} -\gamma u_{1}^{2}(t_{0}) - \gamma u_{2}^{2}(t_{0}) + \alpha u_{3}^{2}(t_{0}) + \alpha u_{4}^{2}(t_{0}) + \alpha d u_{5}^{2}(t_{0}) \end{bmatrix} e^{-\eta(t-t_{0})}$$
(17)

where $\eta = \min(\alpha, c, \beta) > 0$. The above inequality (17) shows that the equilibrium

point O(0,0,0,0) of system (4) is globally exponentially stable.

Thus, the proof is complete.

Remark 4:

The results of this paper can also be used for chaos synchronization, chaos control and the estimation of the Hausdorff dimension of attractors. The applications of the boundedness of chaotic systems in chaos control and chaos synchronization can be referred to the papers [12, 25]. The applications of the boundedness of chaotic systems in the estimation of the Hausdorff dimension of attractors can be referred to the papers [9,14, 21].

6. Conclusions

In this paper, a new generalized complex Lorenz system was proposed and studied by using the theory of chaotic systems. Boundedness and the global exponential attractive set of the complex Lorenz system are obtained. The corresponding boundedness is numerically verified by the computer. Numerical simulations are presented to show the effectiveness of the theoretical research results. Finally, the theoretical results obtained in this paper are used to study the globally exponential stability of the equilibrium point of system (4).

Acknowledgments

This work is supported by the Natural Science Foundation of Chongqing Municipality (Grant No. CSTB2022NSCQ-MSX1548), the Scientific and Technological Research Program of Chongqing Municipal Education Commission (Nos. KJQN202100813, KJQN201800818, KJCX2020037), Scientific and Technological Research Program of Chongqing Technology and Business University (Grant Nos: 1952012, 1952026, 1951075), the Program for Chongqing Key Laboratory of Social Economy and

Applied Statistics (Grant No: ZDPTTD201909). The authors wish to thank the editors and reviewers for their conscientious reading of this paper and their numerous comments for improvement which were extremely useful and helpful in modifying this paper.

References

- [1] E.N. Lorenz, Deterministic non-periodic flow, J. Atmos. Sci., 1963, 20, 130–141.
- [2] H. K. Chen, Synchronization of two different chaotic systems: a new system and each of the dynamical systems Lorenz, Chen and Lü, Chaos, Solitons & Fractals, 2005, 25, 1049-1056.
- [3] H. Richter, Controlling the Lorenz system: combining global and local schemes, Chaos, Solitons & Fractals, 2001, 12(13), 2375-2380.
- [4] G. Chen and T. Ueta, Yet another chaotic attractor, Int. J. Bifurc. Chaos Appl. Sci. Eng., 1999, 9, 1465-1466.
- [5] O. Rossler, An equation for hyperchaos, Phys. Lett. A., 1979, 2–3, 155–157.
- [6] J, Lü and G. Chen, A new chaotic attractor coined, Int. J. Bifurc. Chaos, 2002, 12, 659–661.
- [7] N. Kuznetsov, T. Mokaev, V. Ponomarenko, E. Seleznev, N. Stankevich, L. Chua, Hidden attractors in Chua circuit: mathematical theory meets physical experiments, Nonlinear Dyn., 2023, 111(6), 5859-5887.
- [8] L. Chua, M. Komuro, T. Matsumoto, IEEE Trans. Circuits Syst., 1986, 33(11), 1072-1118.
- [9] N. Kuznetsov, G. Leonov, T. Mokaev, A. Prasad and M. Shrimali, Finite-time Lyapunov dimension and hidden attractor of the Rabinovich system, Nonlinear Dyn., 2018, 92(2), 267–285.
- [10] N. Kuznetsov, T. Mokaev, O. Kuznetsova, et al. The Lorenz system: hidden boundary of practical stability and the Lyapunov dimension, Nonlin. Dyn., 2020, 102, 713–732.
- [11] E.E. Mahmoud, Dynamics and synchronization of new hyperchaotic complex Lorenz system, Mathematical and Computer Modelling, 2012, 55(7-8), 1951 -1962.

- [12] G.M. Mahmoud, M.A. Al-Kashif, A.A. Farghaly, Chaotic and hyperchaotic attractors of a complex nonlinear system, J. Phys. A: Math. Theor., 2008, 41(5), 055104.
- [13] P. Frederickson, J. Kaplan, E. Yorke and J. Yorke, The Lyapunov dimension of strange attractors, J. Differ. Equ., 1983, 49(2), 185–207.
- [14] G. Leonov and V. Boichenko, Lyapunov's direct method in the estimation of the Hausdorff dimension of attractors, Acta Appl. Math., 1992, 26(1), 1–60.
- [15] G. Leonov and N. Kuznetsov, Hidden attractors in dynamical systems. From hidden oscillations in Hilbert-Kolmogorov, Aizerman, and Kalman problems to hidden chaotic attractor in Chua circuits, Int. J. Bifurc. Chaos Appl. Sci. Eng., 2013, 23, 1330002.
- [16] G. Leonov, N. Kuznetsov and T. Mokaev, Homoclinic orbits, and self-excited and hidden attractors in a Lorenz-like system describing convective fluid motion, Eur. Phys. J. Spec. Top., 2015, 224(8), 1421–1458.
- [17] G. Leonov, N. Kuznetsov and V. Vagaitsev, Hidden attractor in smooth Chua systems, Physica D., 2012, 41(18), 1482–1486.
- [18] G. Leonov, Bounds for attractors and the existence of homoclinic orbits in the Lorenz system, J. Appl. Math. Mech., 2001, 65(1), 19–32.
- [19] G. Leonov, A. Bunin and N. Koksch, Attractor localization of the Lorenz system,Z. Angew. Math. Mech., 1978, 67, 649–656.
- [20] L. Stenflo, Generalized Lorenz equations for acoustic-gravity waves in the atmosphere, Phys. Scr., 1996, 3, 83–84.
- [21] N.V. Kuznetsov, The Lyapunov dimension and its estimation via the Leonov method, Phys. Lett. A, 2016, 380(25-26), 2142-2149.
- [22] N.V. Kuznetsov, Theory of hidden oscillations and stability of control systems, J. Comput. Syst. Sci. Int., 2020, 59, 647-668.
- [23] G. Leonov and N. Kuznetsov, Hidden attractors in dynamical systems. From hidden oscillations in Hilbert–Kolmogorov, Aizerman, and Kalman problems to hidden chaotic attractor in Chua circuits, Int. J. Bifurc. Chaos Appl. Sci. Eng., 2013, 23, 1330002.
- [24] G. Leonov, Lyapunov dimension formulas for Henon and Lorenz attractors, St. Petersburg Math. J., 2001, 13, 1–12.
- [25] F. Zhang, Y. Li and C. Mu, Bounds of solutions of a kind of hyper-chaotic

systems and application, J. Math. Res. Appl., 2013, 33 (3), 345–352.

- [26] F. Zhang, R. Chen, X. Wang, X. Chen, C. Mu and X. Liao, Dynamics of a new 5D hyperchaotic system of Lorenz type, Int. J. Bifurc. Chaos Appl. Sci. Eng., 2018, 28(3), 1850036.
- [27] F. Zhang, X. Liao, G. Zhang and C. Mu, Dynamical analysis of the generalized Lorenz systems, J. Dyn. Control Syst., 2017, 23(2), 349–362.
- [28] F. Zhang, X. Liao, G. Zhang, C. Mu, P. Zhou and M. Xiao, Dynamical behaviors of a generalized Lorenz family, Discrete Contin. Dyn. Syst., Ser. B., 2017, 22(10) 3707–3720.
- [29] F. Zhang, X. Liao and G. Zhang, On the global boundedness of the Lu system, Appl. Math. Comput., 2016, 284, 332–339.
- [30] F. Zhang, X. Liao, C. Mu, G. Zhang and Y. Chen, On global boundedness of the Chen system, Discrete Contin. Dyn. Syst., Ser. B., 2017, 22(4), 1673–1681.
- [31] F. Zhang, X. Liao and G. Zhang, Some new results for the generalized Lorenz system, Qual. Theory Dyn. Syst., 2017, 16(3), 749–759.
- [32] F. Zhang, C. Mu and X. Li, On the boundness of some solutions of the Lu system, Int. J. Bifurc. Chaos Appl. Sci. Eng., 2012, 22, 1250015.
- [33] A.C. Fowler, J.D. Gibbon and M.J. McGuinness, The complex Lorenz equations, Physica D. 1982, 4, 139–163.
- [34] A. Rauh, L. Hannibal, N.B. Abraham, Global stability properties of the complex Lorenz model, Physica D: Nonlinear Phenomena, 1996, 99(1), 45–58.
- [35] G.M. Mahmoud, M.A. Al-Kashif, S.A. Aly, Basic properties and chaotic synchronization of complex Lorenz system, International Journal of Modern Physics C, 2007, 18(02), 253–265.
- [36] G.M. Mahmoud, M.E. Ahmed, E.E. Mahmoud, Analysis of hyperchaotic complex Lorenz systems, International Journal of Modern Physics C, 2008, 19(10), 1477–1494.
- [37] A.C. Fowler, J.D. Gibbon and M.J. McGuinness, The real and complex Lorenz equations and their relevance to physical systems, Physica D: Nonlinear Phenomena, 1983, 7(1-3), 126-134.
- [38] J.D. Gibbon and M.J. McGuinness, The real and complex Lorenz equations in rotating fluids and lasers, Physica D: Nonlinear Phenomena, 1982, 5(1), 108-122.
- [39] S. Liu and P. Liu, Adaptive anti-synchronization of chaotic complex nonlinear systems with unknown parameters, Nonlinear Analysis: Real World Applications, 2011, 12, 3046–3055.
- [40] F. Zhang, S. Liu and W. Yu, Characteristics of time-delay complex Lorenz chaotic system and its self-synchronization of time delay, Acta Phys. Sin., 2013, 62(22), 220505.
- [41] X. Liao, Y. Fu, S. Xie and P. Yu, Globally exponentially attractive sets of the

family of Lorenz systems, Sci. China, Ser. F., 2008, 51(3), 283–292.

[42] W. Qin and G. Chen, On the boundedness of solutions of the Chen system, J. Math. Anal. Appl., 2007, 329(11), 445–451.