

# UNIVERSAL APPROACH TO THE TAKESAKI-TAKAI $\gamma$ -DUALITY

M.I. Yaremenko<sup>1,†</sup>

**Abstract** In this article, we generalize and simplify the proof of the Takesaki-Takai  $\gamma$ -duality theorem. Assume a morphism  $\omega : G \rightarrow \text{Aut}(A)$  is a projective representation of the locally compact Abelian group  $G$  in  $\text{Aut}(A)$ , mapping  $\gamma : G \rightarrow G$  is continuous, and  $(A, G, \omega)$  is a dynamic system then there exists isomorphism

$$\Upsilon : \text{Env}_{\hat{\omega}}^{\gamma} \left( L^1 \left( \hat{G}, \text{Env}_{\omega}^{\gamma} \left( L^1(G, A) \right) \right) \right) \rightarrow A \otimes LK(L^2(G))$$

which is the equivariant for the double dual action

$$\hat{\omega} : G \rightarrow \text{Aut} \left( \text{Env}_{\hat{\omega}}^{\gamma} \left( L^1 \left( \hat{G}, \text{Env}_{\omega}^{\gamma} \left( L^1(G, A) \right) \right) \right) \right).$$

These results deepen our understanding of the representation theory and are especially interesting given their possible applications to problems of the quantum theory.

**Keywords** Takai Duality,  $\gamma$ -duality, Wigner function,  $C^*$ -algebra, Pontryagin duality, induced representation, cross product.

**MSC(2010)** 32K05, 46A16, 46A22, 46B25, 46B50, 46E15.

## 1. Introduction

Let  $G$  be a locally compact group, let  $C_c(G)$  be a space of real-valued function with compact support.

**Definition 1.1.** A Radon measure on a locally compact group  $G$  is called a linear form  $\mu$  on  $C_c(G)$  such that for any compact set  $K \subset G$  restriction of the linear form  $\mu$  to subspace  $C_c(K) \subset C_c(G)$  functions of  $C_c(G)$  which support contains in  $K$ , is continuous in the topology of uniform convergence. The value  $\mu(\psi)$  of the Radon measure  $\mu$  on the continuous function  $\psi \in C_c(G)$  with compact support is called a Radon integral of the function  $\psi$ .

As a consequence of the definition, we have that for any compact subset  $K \subset G$  there exists a constant  $\tilde{c}(K)$  dependent on  $K$  such that the equality

$$|\mu(\psi)| \leq \tilde{c} \|\psi\|_{C_c(G)}$$

---

<sup>†</sup>The corresponding author.

<sup>1</sup>Department of PDE, Technical University of Ukraine, PDE, 37, Prospect Beresteiskyi (former Peremohy), Kyiv, Ukraine, 03056,  
Email: [math.kiev@gmail.com](mailto:math.kiev@gmail.com) (M.I. Yaremenko)

holds for all  $\psi \in C_C(G)$ .

Let  $C_C^+(G)$  be a set of all finite positive continuous functions with compact supports. We denote by  $\wp_+(G)$  the set of all lower semicontinuous positive functions i.e., all functions  $\psi$  such that at every point  $g_0$  of its domain satisfy the following condition

$$\liminf_{g \rightarrow g_0, g \in G} \psi(g) = \psi(g_0).$$

**Definition 1.2.** Let  $\mu$  be positive Radon measure on  $G$ , then the upper integral  $\mu^*(\psi)$  of a function  $\psi \in \wp_+(G)$  is defined by

$$\mu^*(\psi) = \sup_{\varphi \in C_C^+(G), \varphi \leq \psi} \mu(\varphi).$$

The upper integral of an arbitrary positive function  $\psi : G \rightarrow R^+$  is defined by

$$\mu^*(\psi) = \inf_{\varphi \in \wp_+(G), \varphi \geq \psi} \mu^*(\varphi).$$

**Definition 1.3.** The outer measure  $\mu^*(E)$  of an arbitrary subset  $E \subset G$  is an upper integral  $\mu^*(1_E)$  of the characteristic function  $1_E$  of  $E$ .

The set  $M(G)$  of all Radon measures  $\mu$  on the locally compact space  $G$  is the space of all linear forms on the vector space  $C_C(G)$  and thus  $M(G)$  is a topological space with the \*-weak or so-called wide topology of the weak convergence. If  $G$  is a compact group then the wide topology coincides with the classical weak topology.

Wide topology in  $M(G)$  can be defined by seminorms  $\mu \mapsto \sup_{1 \leq i \leq k} |\mu(\psi_i)|$ , where  $\{\psi_i\}_{1 \leq i \leq k} \subset C_C(G)$  is an arbitrary finite sequence of functions of  $C_C(G)$ .

The dual group  $\hat{G}$  consists of all homomorphisms (characters) from  $G$  to the circle group with natural measure  $\hat{\mu}(\chi) = \int \chi(g) d\mu(g)$ ,  $\chi \in \hat{G}$ .

The Fourier transform of a function  $\psi \in L^1(G)$  is given by

$$\hat{\psi}(\chi) = \int_G \psi(g) \overline{\chi(g)} d\mu(g).$$

Let  $A$  be a  $C^*$ -algebra then we call a triplet  $(A, G, \omega)$  a dynamical system where  $\omega : G \rightarrow \text{Aut}(A)$  is a strongly continuous representation, and let  $H$  be a Hilbert space then a triplet  $(H, \pi, \rho)$  is called a covariant representation of  $(A, G, \omega)$ .

The Takai duality theory is a generalization of the Takesai duality theorem for the Neumann algebras, which are unital \*-algebras of bounded operators on Hilbert spaces that are closed in the weak operator topology. The classical Takai duality theorem can be formulated as follows: let  $(A, \omega)$  be an action of an Abelian group  $G$  then there exists an isomorphism  $\Upsilon$  from the iterated product  $(A \times_{\omega} G) \times_{\hat{\omega}} \hat{G}$  to the maximal product  $A \otimes LK(L^2(G))$ .

Considerable interest in  $C^*$ -algebras is justified by many applications to the problems of quantum mechanics for instance so-called von Neumann algebras. Some applications of  $C^*$ -algebras to quantum physics are described in [5, 12]. B. Abadie [1] considers the Cuntz-Krieger-Pimsner algebras that be a generalization of the crossed product by the set of integer numbers and Toeplitz and Cuntz-Krieger algebras. In [2, 3], the Cuntz-Pimsner covariance condition is considered as a nondegeneracy condition for representations of cross algebras and a groupoid model for the Cuntz-Pimsner algebra is constructed; in [10], the author considers the  $C^*$ -envelope of a tensor algebra as the corresponding Cuntz-Krieger  $C^*$ -algebra.

We will consider the cross product of  $C^*$ -algebra  $A \times_{\omega} G$  as the universal enveloping  $C^*$ -algebra  $Env_{\omega}(L^1(G, A))$  of the Banach algebra completed in the universal norm. The covariant representation  $(H, \pi, \rho)$  can be unequivocally characterized by morphism  $(\rho \circ \pi) : L^p(G, A) \rightarrow LB(H, H)$ . This approach can be applied to generalize this theory to include the pseudo-differential operators for general quantization. Thus, we could define a binary operation as

$$\begin{aligned} (\Psi_1 \odot_{\gamma} \Psi_2)(g) &= \\ &= \int_G \omega\left(\gamma(g)^{-1} \gamma(h)\right) \Psi_1(h) \omega\left(\gamma(g)^{-1} h \gamma(h^{-1}g), \Psi_2(h^{-1}g)\right) d\mu(h) \end{aligned}$$

and  $\Psi_1^{\odot_{\gamma}}(g) = \omega\left(\gamma(g)^{-1} h \gamma(g^{-1}), (\Psi_1(g^{-1}))^*\right)$  where  $\gamma : G \rightarrow G$  is a continuous function. So, we could define a  $\gamma$ -quantization for  $\gamma : G \rightarrow G$ , and corresponding pseudo-differential operators, and recover the Weyl-Wigner theory; the next logical step in generalization is to consider  $p$ -Schatten classes. For further reading consider a list of references [1-14] and the most recent [15-18].

## 2. The $C^*$ -algebra

Let  $A$  be a  $C^*$ -algebra. Let  $G$  be a locally compact group equipped with Haar measure  $\mu$ . Let for each  $g \in G$  we define a  $C^*$ -algebra isomorphism  $\omega(g) : A \rightarrow A$ , for each fixed  $\psi \in A$  the morphism  $\omega(g, \psi)$  is a continuous mapping  $\omega(\cdot, \psi) : G \rightarrow A$  and satisfies the semigroup condition  $\omega(g, \psi) \circ \omega(h, \psi) = \omega(gh, \psi)$  for all  $g, h \in G$ , a such defined morphism will be denoted  $\omega : G \rightarrow Aut(A)$ . The triplet  $(A, G, \omega)$  is called a dynamical system.

**Definition 2.1.** *Let  $H$  be a separable Hilbert space,  $\pi : G \rightarrow U(H)$  be a continuous unitary representation,  $\rho : A \rightarrow LB(H)$  be a  $*$ -representation, then the covariant representation is a set  $(H, \pi, \rho)$  under the condition  $\pi(g) \rho(\psi) \pi(g)^* = \rho(\omega(g, \psi))$  for all  $g \in G$  and  $\psi \in A$ . Often, the triplet  $(H, \pi, \rho)$  is abbreviated to duplet  $(\pi, \rho)$ .*

Let  $L^p(G, A)$  be a Banach  $*$ -algebra of  $A$ -valued function on  $G$ , with the norm given by

$$\|\Psi\|_{L^p}^p = \int_G \|\Psi(g)\|_A^p d\mu(g),$$

we assume  $p = 1$  and the multiplication operation  $\odot : L^p(G, A) \times L^p(G, A) \rightarrow L^p(G, A)$  is defined by

$$(\Psi_1 \odot \Psi_2)(g) = \int_G \Psi_1(h) \omega(\Psi_2(h^{-1}g)) d\mu(h)$$

and

$$\Psi_1^{\odot}(g) = \omega\left(g, (\Psi_1(g^{-1}))^*\right)$$

for any pair  $\Psi_1, \Psi_2 \in L^p(G, A)$ .

The universal enveloping  $C^*$ -algebra  $Env(L^p(G, A))$  of the Banach  $*$ -algebra  $L^p(G, A)$  is constructed as follows. First, we construct the free tensor algebra

$$\begin{aligned} T(L^p(G, A)) &= G \oplus L^p(G, A) \oplus (L^p(G, A) \otimes L^p(G, A)) \oplus \\ &\oplus (L^p(G, A) \otimes L^p(G, A) \otimes L^p(G, A)) \dots \end{aligned}$$

where  $\oplus$  is the direct sum and  $\otimes$  is the tensor product. Second, the multiplication operation  $\odot : L^p(G, A) \times L^p(G, A) \rightarrow L^p(G, A)$  is bilinear and the tensor product is bilinear so the natural lift is accomplished in such a way as to preserve multiplication as a homomorphism. Third, the universal enveloping algebra  $Env_\omega(L^p(G, A))$  is a quotient space  $Env_\omega(L^p(G, A)) = T(L^p(G, A)) / \sim$ , where the equivalence relation is  $\Psi_1 \otimes \Psi_2 - \Psi_2 \otimes \Psi_1 = \Psi_1 \odot \Psi_2$ . The set  $I$  of all elements generated by elements given by  $\Psi_1 \otimes \Psi_2 - \Psi_2 \otimes \Psi_1 - \Psi_1 \odot \Psi_2$  is a two-side ideal so  $I$  lies in the kernel of the quotient map, so we have the short **exact** sequence

$$0 \rightarrow I \rightarrow T(L^p(G, A)) \rightarrow T(L^p(G, A)) / I \rightarrow 0$$

since the sequence is exact, the kernel of the map coincides with the image of the mapping before. In this interpretation, the universal enveloping  $C^*$ -algebra  $Env_\omega(L^p(G, A))$  is defined as  $Env_\omega(L^p(G, A)) = T(L^p(G, A)) / I$ .

The universal norm is given as

$$\|\Psi\|_{U_n} = \sup_{\Pi} \|\Pi(\Psi)\|_{LB(H)},$$

where mapping  $\Pi$  is a representation of  $L^p(G, A)$  in  $LB(H, H)$ .

The integral transformation  $(\rho \times \pi) : L^p(G, A) \rightarrow LB(H, H)$  defined by

$$(\rho \times \pi)(\Psi) = \int_G \rho(\Psi(g)) \pi(g) d\mu(g)$$

extends to mapping  $(\rho \times \pi) : Env_\omega(L^p(G, A)) \rightarrow LB(H, H)$  due to the universality of enveloping  $C^*$ -algebra.

### 3. The Takesaki-Takai duality

Let morphism  $\omega : G \rightarrow Aut(A)$  be a projective representation of the locally compact Abelian group  $G$  in  $Aut(A)$ . We denote a  $C^*$ -algebra of compact operators on a separable Hilbert space  $H$  by  $LK(H)$ . The morphism  $\omega : G \rightarrow Aut(A)$  is called an action of the group  $G$ . Let a triplet  $(A, G, \omega)$  be a dynamical system. We obtain the dual action as the homomorphism

$$\hat{\omega} : \hat{G} \rightarrow Aut(Env_\omega(L^1(G, A))),$$

then the triplet  $(Env_\omega(L^1(G, A)), \hat{G}, \hat{\omega})$  is called the dual dynamic system.

**Theorem 3.1. (Variant of the Takai duality).** *Let  $G$  be a locally compact Abelian group and let  $(A, G, \omega)$  be the dynamic system. Then,  $Env_{\hat{\omega}}(L^1(\hat{G}, Env_\omega(L^1(G, A))))$  isomorphically equals  $A \otimes LK(L^2(G))$ , so there exists such isomorphism  $\Upsilon : Env_{\hat{\omega}}(L^1(\hat{G}, Env_\omega(L^1(G, A)))) \rightarrow A \otimes LK(L^2(G))$  which is equivariant for the double dual action  $\hat{\hat{\omega}} : G \rightarrow Aut(Env_{\hat{\omega}}(L^1(\hat{G}, Env_\omega(L^1(G, A))))$  and equivariant for  $\omega \otimes Ad(\zeta) : G \rightarrow Aut(A \otimes LK(L^2(G)))$ .*

**Proof.** The statement of the Takai duality theorem will be proven if we show that there is a sequence of the following isomorphisms:

$$\begin{aligned}
 & \text{Env}_{\hat{\omega}} \left( L^1 \left( \hat{G}, \text{Env}_{\omega} \left( L^1(G, A) \right) \right) \right) \xrightarrow{\Upsilon_1} \\
 & \xrightarrow{\Upsilon_1} \text{Env}_{\hat{\lambda}^{-1} \otimes \omega} \left( L^1 \left( G, \text{Env}_{\text{Id}} \left( L^1 \left( \hat{G}, A \right) \right) \right) \right), \\
 & \text{Env}_{\hat{\lambda}^{-1} \otimes \omega} \left( L^1 \left( G, \text{Env}_{\text{Id}} \left( L^1 \left( \hat{G}, A \right) \right) \right) \right) \xrightarrow{\Upsilon_2} \text{Env}_{\lambda \otimes \omega} \left( L^1(G, C_0(G, A)) \right), \\
 & \text{Env}_{\lambda \otimes \omega} \left( L^1(G, C_0(G, A)) \right) \xrightarrow{\Upsilon_3} \text{Env}_{\lambda \otimes \text{Id}} \left( L^1(G, C_0(G, A)) \right), \\
 & \text{Env}_{\lambda \otimes \text{Id}} \left( L^1(G, C_0(G, A)) \right) \xrightarrow{\Upsilon_4} \text{Env}_{\lambda} \left( L^1(G \otimes A, C_0(G)) \right), \\
 & \text{Env}_{\lambda} \left( L^1(G \otimes A, C_0(G)) \right) \xrightarrow{\Upsilon_5} LK(L^2(G)) \otimes A,
 \end{aligned}$$

so that the isomorphism in question can be presented as  $\Upsilon = \Upsilon_5 \circ \Upsilon_4 \circ \Upsilon_3 \circ \Upsilon_2 \circ \Upsilon_1$ , where  $\lambda$  is left translation.

Let  $K$  be compact, by construction, the set  $C_C(K \times H, A)$  is a dense subspace of  $\text{Env}_{\beta} \left( L^1(H, \text{Env}_{\alpha} \left( L^1(K, A) \right)) \right)$ . Since the topology of  $C_C(K, A)$  is induced by the topology of  $L^1$ - norm, we presume  $C_C(K, A) \subset \text{Env}_{\alpha} \left( L^1(K, A) \right)$  is invariant under homomorphism  $\beta$  and there is  $f(h, g) \in C_C(H \times K, A)$  such that  $f(h, g) = \beta(\ell_f(h))(g)$  where  $\ell_f \in C_C(H, \text{Env}_{\alpha} \left( L^1(K, A) \right))$ .

The proof will follow from the next statements.

**Statement 1.** The isomorphism

$$\begin{aligned}
 \Upsilon_1 & : \text{Env}_{\hat{\omega}} \left( L^1 \left( \hat{G}, \text{Env}_{\omega} \left( L^1(G, A) \right) \right) \right) \rightarrow \\
 & \rightarrow \text{Env}_{\hat{\lambda}^{-1} \otimes \omega} \left( L^1 \left( G, \text{Env}_{\text{Id}} \left( L^1 \left( \hat{G}, A \right) \right) \right) \right)
 \end{aligned}$$

maps dense subalgebras

$$\Upsilon_1 : C_C \left( \hat{G} \times G, A \right) \xrightarrow{\text{onto}} C_C \left( G \times \hat{G}, A \right)$$

so that  $\Upsilon_1(f)(g, \chi) = \chi(g) f(\chi, g)$  for all  $(g, \chi) \in G \times \hat{G}$  and  $f \in C_C \left( \hat{G} \times G, A \right)$ .

**Statement 2.** Let  $G$  be an Abelian group and let  $C_C(A, G, \omega)$  be a dynamical system then the mapping  $\Upsilon_2 : C_C \left( G \times \hat{G}, A \right) \rightarrow C_C \left( G, C_0(G, A) \right)$  is given by  $\Upsilon_2(f)(g, h) = \int_{\hat{G}} f(g, \chi) \overline{\chi(h)} d\hat{\mu}(\chi)$ , the mapping

$$\Upsilon_2 : \text{Env}_{\hat{\lambda}^{-1} \otimes \omega} \left( L^1 \left( G, \text{Env}_{\text{Id}} \left( L^1 \left( \hat{G}, A \right) \right) \right) \right) \rightarrow \text{Env}_{\lambda \otimes \omega} \left( L^1(G, C_0(G, A)) \right)$$

is an isomorphism.

**Statement 3.** Let  $G$  be an Abelian group and let  $(A, G, \omega)$  be a dynamical system then there exists an isomorphism

$$\Upsilon_3 : \text{Env}_{\lambda \otimes \omega} \left( L^1(G, C_0(G, A)) \right) \rightarrow \text{Env}_{\lambda \otimes \text{Id}} \left( L^1(G, C_0(G, A)) \right)$$

such that equality  $\Upsilon_3(f)(g, h) = \omega^{-1}(h, f(g, h))$  holds for all  $f \in C_C \left( G, C_0(G, A) \right)$ .

**Statement 4.** Let  $G$  be a locally compact group and let  $(A, G, \omega)$  be a dynamical system then there exists an isomorphism  $\tilde{\Upsilon}_5 = \Upsilon_5 \Upsilon_4$  such that

$$\tilde{\Upsilon}_5 : Env_{\lambda \otimes \omega} (L^1(G, C_0(G, A))) \rightarrow LK(L^2(G)) \otimes A.$$

Proof statement 1. In order to prove statement 1, we must show that  $\|\Upsilon_1(f)\| = \|f\|$ .

Let  $f_1, f_2 \in C_C(H \times K, A)$  then

$$((h, g) \mapsto \ell_{f_1}(h) * \alpha(h, \ell_{f_1}(h^{-1}s))(g)) \in C_C(H \times K, A),$$

so that  $\ell_{f_1} * \ell_{f_2} \in C_C(K, A) \subset Env_{\alpha}(L^1(K, A))$  and we have

$$\begin{aligned} & (\ell_{f_1} * \ell_{f_2})(s)(g) = \\ &= \int_H \int_K \ell_{f_1}(h, t) \omega(t, \alpha(h, (\ell_{f_2}(h^{-1}s)), t^{-1}g)) d\mu_K(t) d\mu_H(h). \end{aligned}$$

Thus, we obtain those equalities

$$\begin{aligned} & (\ell_{f_1} * \ell_{f_2})(\chi)(g) = \\ &= \int_{\hat{G}} \int_G \ell_{f_1}(\zeta, t) \overline{\zeta(t^{-1}g)} \omega(t, \alpha(\ell_{f_2}(\bar{\zeta}\chi), t^{-1}g)) d\mu(t) d\hat{\mu}(\zeta) \end{aligned}$$

and dual

$$\begin{aligned} & (\ell_{\tilde{f}_1} * \ell_{\tilde{f}_2})(g)(\chi) = \\ &= \int_G \int_{\hat{G}} \ell_{\tilde{f}_1}(t, \zeta) \bar{\zeta}(t) \chi(t) \omega(t, \alpha(\ell_{\tilde{f}_2}(t^{-1}g), \bar{\zeta}\chi)) d\hat{\mu}(\zeta) d\mu(t) \end{aligned}$$

hold for all  $f_1, f_2 \in C_C(\hat{G} \times G, A)$  and for all  $\tilde{f}_1, \tilde{f}_2 \in C_C(G \times \hat{G}, A) \subset Env_{\lambda \otimes \omega}(L^1(G, Env_{Id}(L^1(G, A))))$ . Then, we have a homomorphism

$$\Upsilon_1 : C_C(\hat{G} \times G, A) \xrightarrow{onto} C_C(G \times \hat{G}, A).$$

We write the equalities

$$\begin{aligned} & \ell^*_{\Upsilon_1(f)}(g)(\chi) = \\ &= (\hat{\lambda}^{-1} \otimes \omega)(g)(\ell_{\Upsilon_1(f)}(g^{-1}))(\chi) = \chi(g) \omega(g, \ell_{\Upsilon_1(f)}(g^{-1}))^*(\chi) = \\ &= \chi(g) \omega(g, \Upsilon_1(f)(g^{-1}, \bar{\chi}))^* = \omega(g, f(\bar{\chi}, g^{-1}))^* = \omega(g, \ell_f(\bar{\chi})(g^{-1}))^* \\ &= \chi(g) \ell^*_f(\bar{\chi})(g) = \Upsilon_1(\ell^*_f)(\chi)(g). \end{aligned}$$

In general, every continuous in the inductive topology  $*$ -homomorphism is bounded in the topology of the universal norm thus this  $*$ -homomorphism extends to a representation on  $Env_{\omega}(L^1(G, A))$ .

Let  $U : G \rightarrow U(H)$  be a unitary representation and  $(U, \rho)$  be a covariant representation of the dynamic system  $(Env_{Id}(L^1(\hat{G}, A)), G, \hat{\lambda}^{-1} \otimes \omega)$ , and  $(V, \pi)$  be a covariant representation of  $(A, \hat{G}, Id)$ , then we denote

$$\Lambda := (U, \rho)(f) = \int_G \rho(f(g)) U(g) d\mu(g)$$

so that  $\Lambda = (U, \rho) : \text{Env}_{\hat{\lambda}^{-1} \otimes \omega} \left( L^1 \left( G, \text{Env}_{Id} \left( L^1 \left( \hat{G}, A \right) \right) \right) \right) \rightarrow LB(H)$ , and

$$\rho := (V, \pi)(f) = \int_{\hat{G}} \pi(f(\hat{g})) V(\hat{g}) d\hat{\mu}(\hat{g}).$$

Next, we have  $\hat{\lambda}^{-1}(g) \circ \phi(\bar{\chi}\zeta) = \chi(g) \phi(\bar{\chi}\zeta) \circ \hat{\lambda}^{-1}(g)$ . Let us take  $a \in A$ ,  $\psi \in C_C(G)$  and  $\phi \in C_C(\hat{G})$  so that all linear combinations  $a \otimes \phi \otimes \psi$  constitute a dense subset of  $C_C(\hat{G} \times G, A)$ , so that

$$\begin{aligned} U(g) V(\chi) \Lambda(a \otimes \phi \otimes \psi) &= U(g) V(\chi) \pi(a) V(\phi) U(\psi) = \\ &= \pi(\omega(g, a)) V(\hat{\lambda}^{-1}(g) \circ \phi(\bar{\chi}\zeta)) U(g) U(\psi) = \\ &= \chi(g) V(\chi, \pi(\omega(g, a))) V(\hat{\lambda}^{-1}(g, \phi)) U(g) U(\psi) = \\ &= \chi(g) V(\chi) U(g) \Lambda(a \otimes \phi \otimes \psi) \end{aligned}$$

and we obtain  $U(g) V(\chi) = \chi(g) V(\chi) U(g)$ . Next, we write

$$\begin{aligned} U(g) \pi(b) \pi(a) V(\phi) U(\psi) &= \\ &= \pi(\omega(g, ba)) V(\hat{\lambda}^{-1}(g, \phi)) U(g) U(\psi) = \\ &= \pi(\omega(g, b)) U(g) \pi(a) V(\phi) U(\psi). \end{aligned}$$

We compute

$$\begin{aligned} \Lambda(\Upsilon_1(f)) &= \int_G \int_{\hat{G}} \pi(\Upsilon_1(f)(\chi, g)) V(\chi) U(g) d\hat{\mu}(\chi) d\mu(g) = \\ &= \int_G \int_{\hat{G}} \pi(f(\chi, g)) \chi(g) V(\chi) U(g) d\hat{\mu}(\chi) d\mu(g) = \\ &= \int_{\hat{G}} \int_G \pi(f(\chi, g)) U(g) V(\chi) d\mu(g) d\hat{\mu}(\chi) \end{aligned}$$

so, we obtain  $\|\Upsilon_1(f)\| \leq \|f\|$ , the similarly, we obtain  $\|f\| \leq \|\Upsilon_1(f)\|$  and  $\Upsilon_1 : \text{Env}_{\hat{\omega}} \left( L^1 \left( \hat{G}, \text{Env}_{\omega} \left( L^1(G, A) \right) \right) \right) \rightarrow \text{Env}_{\hat{\lambda}^{-1} \otimes \omega} \left( L^1 \left( G, \text{Env}_{Id} \left( L^1 \left( \hat{G}, A \right) \right) \right) \right)$  is an isomorphism, statement 1 is proven.

Proof statement 2. The isomorphism  $\text{Env}_{Id} \left( L^1 \left( \hat{G}, A \right) \right) \rightarrow C_0(G, A)$  given by  $\langle \psi \bar{\chi} \rangle_{\hat{G}}$  can be constructed as an extension of the mapping  $a \otimes \phi \mapsto a \otimes \hat{\phi}$  that is defined on the span of bases as  $A \otimes C^*(\hat{G}) \cong \text{Env}_{Id} \left( L^1 \left( \hat{G}, A \right) \right) \rightarrow C_0(G, A) \cong C_0(G) \otimes A$ . The mapping  $\Upsilon_2 := \langle \psi \bar{\chi} \rangle_{\hat{G}} \otimes Id$  is equivariant isomorphism since

$$\begin{aligned} (\lambda \otimes \omega)(g) \int_{\hat{G}} \psi(\chi) \overline{\chi(g)} d\hat{\mu}(\chi) &= \\ &= \int_{\hat{G}} (\lambda^{-1} \otimes \omega)(g) \psi(\chi) \overline{\chi(g)} d\hat{\mu}(\chi), \end{aligned}$$

statement 2 is proven.

Proof of statement 3. Since  $\omega^{-1}(h, \varphi(h))$  is an isomorphism  $C_0(G, A) \rightarrow C_0(G, A)$ , statement 3 follows from

$$\begin{aligned} \omega^{-1}(g, (\lambda \otimes \omega)(g, \varphi))(h) &= \omega^{-1}(h, \omega(g, \varphi(g^{-1}h))) = \omega^{-1}(g^{-1}h, \varphi(g^{-1}h)) = \\ &= (\lambda \otimes \omega)(g) \omega^{-1}(h, \varphi(h)), \end{aligned}$$

so that  $\Upsilon_3(f)(g, h) = \omega^{-1}(h, f(g, h))$ .

Proof of statement 4. Let  $\Delta$  be a modular function on  $G$ , namely,  $\Delta : G \rightarrow R_+$  is a continuous homomorphism and the equality

$$\Delta(g) \int_G \psi(hg) d\mu(h) = \int_G \psi(h) d\mu(h)$$

holds for all  $\psi \in C_C(G)$ . Next, we must show that  $Env_\lambda(L^1(G, C_0(G))) \cong LK(L^2(G))$ . The  $Env_\lambda(L^1(G, C_0(G)))$  is simple. We define a natural covariant representation  $(M, l)$  of  $(C_0(G), G, \lambda)$  as  $M(\psi)\varphi(g) = \psi(g)\varphi(g)$  where  $l : G \rightarrow U(L^2(G))$  is the left-regular representation and  $M$  operator of pointwise multiplication. Let  $k \in C_C(G \times G)$  then  $\Delta(h^{-1}g)k(g, h^{-1}g) = \psi_k(h, g)$ ,  $\psi_k \in C_C(G \times G)$  so that

$$\begin{aligned} & \int_G \langle M(\psi_k(g, \cdot))l(g)\varphi_1, \varphi_2 \rangle_{L^2} d\mu(g) = \\ & = \int_G \int_G \psi_k(g, h)\varphi_1(g^{-1}h)\overline{\varphi_2(h)} d\mu(g)d\mu(h) = \\ & = \int_G \int_G \Delta(g^{-1})k(h, g^{-1}h)\varphi_1(g^{-1}h)\overline{\varphi_2(h)} d\mu(g)d\mu(h) = \\ & = \int_G \int_G k(h, g)\varphi_1(g)\overline{\varphi_2(h)} d\mu(g)d\mu(h). \end{aligned}$$

The kernel  $k \in C_C(G \times G) \subset L^2(G \times G)$  defines a compact Hilbert-Schmidt operator. Since  $C_C(G)$  is dense in  $L^2(G)$  we have  $LK(L^2(G))$  belongs to the image of a compact Hilbert-Schmidt operator with kernel  $k$  mapping  $Env_\lambda(L^1(G, C_0(G)))$ . Assume  $\psi \in C_C(G \times G)$  we denote  $k(h, g) = \Delta(g^{-1})\psi(hg^{-1}, h)$  so  $\psi_k = \psi$ , and  $Env_\lambda(L^1(G, C_0(G))) \cong LK(L^2(G))$  follows from the density of  $C_C(G \times G)$  in  $Env_\lambda(L^1(G, C_0(G)))$ .

So, since

$$\begin{aligned} & \Delta(t)^{\frac{1}{2}} \int_G \psi(g, ht)\varphi(g^{-1}ht) d\mu(g) = \\ & = \int_G (\rho \otimes Id)(t, \psi)(g, h)\tau(t)\varphi(g^{-1}h) d\mu(g) = \end{aligned}$$

the mapping given by integration  $\int_G \psi(g, h)(g^{-1}h) d\mu(g)$  defines an equivariant isomorphism

$$(Env_\lambda(L^1(G, C_0(G))), G, \rho \otimes Id) \rightarrow (LK(L^2(G)), G, Ad(\tau)),$$

where  $\rho$  is a right translation of the group  $G$  on itself.

Thus, we obtain the existence of the equivariant isomorphism

$$\tilde{\Upsilon}_5 : Env_{\lambda \otimes Id}(L^1(G, C_0(G, A))) \rightarrow LK(L^2(G)) \otimes A,$$

statement 5 is proven so proof of the variant of the Takai duality theorem is completed.

## 4. The general cross product $C^*$ -algebra

Let  $\gamma : G \rightarrow G$  be a continuous mapping, we define an enveloping  $C^*$ -algebra  $Env_\omega^\gamma(L^p(G, A))$  as  $T(L^p(G, A))/I$  where mapping  $I$  is the two-sided ideal generated by elements

$$\Psi_1 \otimes \Psi_2 - \Psi_2 \otimes \Psi_1 - \Psi_1 \circ_\gamma \Psi_2,$$



where a binary operation  $\odot_\gamma$  is defined by

$$\begin{aligned} & (\Psi_1 \odot_\gamma \Psi_2)(g) = \\ & = \int_G \omega\left(\gamma(g)^{-1} \gamma(h)\right) \Psi_1(h) \omega\left(\gamma(g)^{-1} h \gamma(h^{-1}g)\right) \Psi_2(h^{-1}g) d\mu(h). \end{aligned}$$

Thus, we generalized the **Takai duality theory** on  $\gamma$ -case as follows.

**Theorem 4.1.** ( *$\gamma$ -variant of the Takai duality*). *Let  $G$  be a locally compact Abelian group, let  $\gamma : G \rightarrow G$  be a continuous mapping, and let  $(A, G, \omega)$  be the dynamic system. Then,  $\text{Env}_{\hat{\omega}}^\gamma\left(L^1\left(\hat{G}, \text{Env}_\omega^\gamma\left(L^1(G, A)\right)\right)\right)$  isomorphically equals  $A \otimes LK\left(L^2(G)\right)$ , so there exists such isomorphism*

$$\Upsilon : \text{Env}_{\hat{\omega}}^\gamma\left(L^1\left(\hat{G}, \text{Env}_\omega^\gamma\left(L^1(G, A)\right)\right)\right) \rightarrow A \otimes LK\left(L^2(G)\right),$$

which is equivariant for the double dual action

$$\hat{\omega} : G \rightarrow \text{Aut}\left(\text{Env}_{\hat{\omega}}^\gamma\left(L^1\left(\hat{G}, \text{Env}_\omega^\gamma\left(L^1(G, A)\right)\right)\right)\right).$$

The proof is similar to the previous theorem.

## 5. Conclusions

This paper dedicated to dynamical systems and  $C^*$ -algebras. We establish that the enveloping  $C^*$ -algebra  $\text{Env}_{\hat{\omega}}^\gamma\left(L^1\left(\hat{G}, \text{Env}_\omega^\gamma\left(L^1(G, A)\right)\right)\right)$  with a pointwise convergence topology is isomorphically identical to maximal product  $A \otimes LK\left(L^2(G)\right)$ . In our future works, we will generalize this statement to include the classes of non-abelian groups  $G$  and wide class functions  $\gamma : G \rightarrow G$ , we also plan to develop a new approach to its application to symmetry in quantum mechanics.

## References

- [1] B. Abadie, *Takai duality for crossed products by Hilbert  $C^*$ -bimodules*, J. Operator Theory 64 (2010), 19–34.
- [2] S. Albandik and R. Meyer, *Product systems over Ore monodies*, Doc. Math. 20 (2015) 1331–1402.
- [3] A. Alldridge, C. Max, M. R. Zirnbauer, *Bulk-Boundary Correspondence for Disordered Free-Fermion Topological Phases*, Commun. Math. Phys. 377, 1761–1821 (2020).
- [4] E. Bedos, S. Kaliszewski, J. Quigg, and D. Robertson, *A new look at crossed product correspondences and associated  $C^*$ -algebras*, J. Math. Anal. Appl. 426 (2015), 1080–1098.
- [5] N. Dupuis, L. Canet, A. Eichhorn, W. Metzner, J. Pawłowski, M. Tissier, and N. Wschebor, *The nonperturbative functional renormalization group and its applications*, Physics Reports 910, 1–114 (2021).
- [6] V. Deaconu, *Group actions on graphs and  $C^*$ -correspondences*, Houston J. Math. 44 (2018), 147–168.

- [7] V. Deaconu, A. Kumjian, and J. Quigg, *Group actions on topological graphs*, Ergodic Theory Dynam. Systems 32 (2012),1527–1566.
- [8] A. Carey, G. C. Thiang, *The Fermi gerbe of Weyl semimetals*, Letters Math. Phys. 111, 1-16 (2021).
- [9] S. Kaliszewski, J. Quigg and D. Robertson, *Coactions on Cuntz-Pimsner algebras*, Math. Scand. 116 (2015), 222–249.
- [10] E. Katsoulis, *Non-selfadjoint operator algebras: dynamics, classification, and  $C^*$ -envelopes*, Recent advances in operator theory and operator algebras, 27–81, CRC Press, Boca Raton, FL, (2018).
- [11] E. Katsoulis,  *$C^*$ -envelopes and the Hao-Ng Isomorphism for discrete groups*, International Mathematics Research Notices, Volume 2017, Issue 18 (2017), 5751–5768.
- [12] I. Raeburn, *Dynamical systems and operator algebras. In National Symposium on Functional Analysis, Optimization and Applications*, pages 109–119. Australian National University, Mathematical Sciences Institute, (1999).
- [13] S. Sundar,  *$C^*$ -algebras associated to topological Ore semigroups*, Munster J. of Math. 9 (2016), no. 1, 155–185.
- [14] M.I. Yaremenko, *Calderon-Zygmund Operators and Singular Integrals*, Applied Mathematics & Information Sciences: Vol. 15: Iss. 1, Article 13, (2021).
- [15] F.F. Miao, G.L. Wang, W. Yuan, *Product Systems of  $C^*$ -correspondences and Baaĵ—Skandalis Duality*, Acta Mathematica Sinica, English Series 39, no. 2 (2023), 240-256.
- [16] A. Dor-On, E. Katsoulis, K. Laca,  *$C^*$ -envelopes for operator algebras with a coaction and couniversal  $C^*$ -algebras for product systems*, Adv. Math., 400, 108286, 40 pp. (2022).
- [17] G. Szabo, *Equivariant property (SI) revisited*, Anal. PDE 14 (2021), no. 4, 1199–1232.
- [18] A. McKee, R. Pourshahami, *Amenable and inner amenable actions and approximation properties for crossed products by locally compact groups*, Canad. Math. Bull. 65 (2) R. (2022), 381–399.