

# The solvability of some kinds of singular integral equations of convolution type with variable integral limits

Wenwen Zhang, Yanxin Lei, Pingrun Li \*

(School of Mathematical Sciences, Qufu Normal University, Qufu 273165, China)

**Abstract** In this paper, we discuss several classes of convolution type singular integral equations with variable integral limits in class  $H_1^*$ . By means of the theory of complex analysis, Fourier analysis and integral transforms, we can transform singular integral equations with variable integral limits into the Riemann boundary value problems with discontinuous coefficients. Under the solvability conditions, the existence and uniqueness of the general solutions can be obtained. Further, we analyze the asymptotic properties of the solutions at the nodes. Our work improves the Noether theory of singular integral equations and boundary value problems, and develops the knowledge architecture of complex analysis.

**Keywords** Singular integral equations; variable integral limits; Riemann boundary value problems; Wiener-Hopf type; dual type; Fourier integral transform

**MR(2010) Subject Classification**

## 1 Introduction

Singular integral equations (SIEs) are closely related to the classical theory of boundary value problems for analytic functions, which has a wide range of applications in many fields, such as quantum mechanics, asymptotic analysis, statistical physics and orthogonal polynomial theory. Many scholars have systematically researched convolution type SIEs and formed a rigorous theoretical system.

Gakhov [1-3] studied the general solutions of Riemann boundary value problems. Litvinchuk [4] obtained the explicit solutions of SIEs in more general cases, and further developed the Noether theory of the general SIEs. Muskhelishvili [5] investigated the conditions of solvability for SIEs with convolution kernels and discontinuous coefficients. Lu [6,7] considered the explicit solutions and the solvability theory of convolution SIEs with constant coefficients, and obtained some worthwhile results. Du and Shen [8] further dealt with the integral equations of convolution type with variable coefficients. Subsequently, Li and Ren [9-12] developed the theory of solvability and asymptotic theory for singular integral equations with the mixture of convolution kernel and singular integral kernel in the case of non-normal type.

---

<sup>1</sup>\* Corresponding author at: School of Mathematical Science, Qufu Normal University, Qufu, 273165, P.R.China.

<sup>2</sup>E-mail address: lipingrun@163.com (P. Li)

Based on the above work, in this paper, we will deal with SIEs of convolution type with variable integral limits, and we investigate the asymptotic properties and Noether solvability theory of solutions for such SIEs with variable integral limits under the solvability conditions. The main aim of this paper is to solve the following three classes SIEs with convolution kernel and variable integral limits.

(1) SIEs with two convolution kernels

$$A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_0^\tau \psi(t)h(\tau-t)dt + \frac{C}{\sqrt{2\pi}} \int_\tau^0 \psi(t)k(\tau-t)dt + \frac{D}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau), \quad \tau \in \mathbb{R}. \quad (1.1)$$

(2) SIEs of Wiener-Hopf with convolution kernels

$$A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_0^\tau \psi(t)h(\tau-t)dt + \frac{C}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau), \quad \tau \in \mathbb{R}^+. \quad (1.2)$$

(3) Dual SIEs with convolution kernels

$$\begin{cases} A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_0^\tau \psi(t)h_1(\tau-t)dt + \frac{C}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau), & \tau \in \mathbb{R}^+; \\ A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_\tau^0 \psi(t)h_2(\tau-t)dt + \frac{C}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau), & \tau \in \mathbb{R}^-. \end{cases} \quad (1.3)$$

In Eqs. (1.1)-(1.3),  $A, B, C, D$  are constants. The known functions  $h, k, n, h_j \in H_1, j = 1, 2$ . We require the unknown function  $\psi \in H_1$ . The notations mentioned above can be referred to Section 2.

It is well known that SIEs with variable integral limits, an important class of equations in physics, is closely related to the Riemann boundary value problems. By the Sokhotski-Plemelj formula and the principle of analytic continuation, we transform Eqs. (1.1)-(1.3) into linear Riemann-Hilbert problems with discontinuous coefficients, and prove the existence and uniqueness of analytic solutions given by integral-form. We propose a novel approach different from the one used in the classical Riemann-Hilbert problems. In view of the theory of classical boundary value problems for analytic functions, we study the properties of solutions at nodes, and obtain the solvability conditions and asymptotic properties of the general solutions. Therefore, this paper develops the theory of Noether solvability of SIEs and boundary value problems, and extends the ones in [13-18].

This paper is arranged as follows. In Section 2, we give properties of the function classes  $H_1^*(H_2^*, H_3^*), H_1(H_2, H_3)$ , and the relation between Cauchy type integrals and Fourier transform. In Sections 3-5, by using the boundary value theory, complex analysis and the system of linear algebra, the explicit solutions and asymptotic properties are obtained under the conditions of solvability, and the properties of solutions at nodes are further analyzed.

## 2 Preliminaries

$\mathbb{R}, \mathbb{C}$  denote the sets of real and complex numbers respectively,  $\bar{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$ . As usual,  $C(\mathbb{R}), C(\bar{\mathbb{R}})$  denote the sets of continuous functions on  $\mathbb{R}$  and  $\bar{\mathbb{R}}$  respectively. For  $1 \leq q < \infty$ ,

the space of Lebesgue integrable functions  $L^q(\mathbb{R}) = \left\{ \Phi \mid \|\Phi\|_q < \infty \right\}$  with the standard norm

$$\|\Phi\|_q = \left( \int_{\mathbb{R}} |\Phi(s)|^q ds \right)^{\frac{1}{q}},$$

where  $\mathbb{R}^- = (-\infty, 0)$ ,  $\mathbb{R}^+ = (0, \infty)$ ,  $\mathbb{R} = (-\infty, \infty)$ .

Moreover, we respectively denote by  $H(\mathbb{R})$ ,  $\hat{H}(\overline{\mathbb{R}})$  as the spaces of Hölder continuous functions on  $\mathbb{R}$  and  $\overline{\mathbb{R}}$ .

In the following, we give some necessary preliminary knowledge and notations.

**Definition 2.1.** Let  $\psi \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ , denote the Fourier integral transform  $\mathcal{W}$  and inverse transform  $\mathcal{W}^{-1}$  as

$$\mathcal{W}[\psi(\tau)] = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{is\tau} \psi(\tau) d\tau, \quad \mathcal{W}^{-1}[\Phi(s)] = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-is\tau} \Phi(s) ds. \quad (2.1)$$

For simplicity, (2.1) can be denoted as

$$\mathcal{W}[\psi(\tau)] = \Phi(s), \quad \mathcal{W}^{-1}[\Phi(s)] = \psi(\tau). \quad (2.2)$$

From (2.1) and (2.2), we know that

$$\mathcal{W} : L^1(\mathbb{R}) \cap L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}). \quad (2.3)$$

Since  $L^1(\mathbb{R}) \cap L^2(\mathbb{R})$  is dense in  $L^2(\mathbb{R})$ , from the Planchere theorem [19], the operator  $\mathcal{W}$  can be uniquely extended to a self-mapping

$$\mathcal{W} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}), \quad (2.4)$$

and

$$\|\Phi\|_2 = \|\psi\|_2. \quad (2.5)$$

We introduce the concepts of several classes  $H_1, H_2, H_3$  and  $H_1^*, H_2^*, H_3^*$ .

**Definition 2.2.** If  $\Phi \in L^2(\mathbb{R}) \cap \hat{H}(\overline{\mathbb{R}})$ , we say  $\Phi \in H_1^*$ . If  $\Phi \in H_1^*$ , then  $\mathcal{W}^{-1}\Phi = \psi \in H_1$ .

**Definition 2.3.** If (1)  $\Phi \in \hat{H}(\overline{\mathbb{R}})$ ; (2) For  $\ell > \frac{1}{2}$ ,  $s \in N_\infty$ ,  $\Phi(s) = O(|s|^{-\ell})$ , we say  $\Phi \in H_2^{*\ell}$  or  $H_2^*$ , where  $N_\infty = \{s \in \mathbb{R} \mid |s| > \delta^{-1}, \forall \delta > 0\}$ .

For  $\Phi \in H_2^{*\ell}$  or  $H_2^*$ , we say that  $\mathcal{W}^{-1}\Phi = \psi \in H_2^\ell$  or  $H_2$ .

**Definition 2.4.** If (1)  $\Phi \in \hat{H}(\overline{\mathbb{R}}) \cap H^\ell(N_\infty)$ ,  $\ell > \frac{1}{2}$ ; (2)  $\Phi(\infty) = 0$ , then  $\Phi \in H_3^{*\ell}$  or  $H_3^*$ , and  $\psi \in H_3^\ell$  or  $H_3$ .

For  $s \in N_\infty$ , we put

$$h_1 = \{\Phi \mid \Phi(s) = O(|s|^{-\ell})\}, \quad \underline{h_2 = H^\ell(N_\infty)}, \quad h_3 = \{\Phi \mid \Phi(\infty) = 0\}.$$

If  $\Phi \in h_2 \cap h_3$ , we have

$$|\Phi(s_1) - \Phi(s_2)| \leq a \left| \frac{1}{s_1} - \frac{1}{s_2} \right|^\ell, \quad s_1, s_2 \in N_\infty, \quad (2.6)$$

where  $a \in \mathbb{R}^+$ ,  $\ell \in (0, 1]$ . In (2.6), we let  $s_2 \rightarrow \infty$ , since  $\lim_{s_2 \rightarrow \infty} \Phi(s_2) = 0$ , thus we have

$$|\Phi(s_1)| \leq a \left| \frac{1}{s_1} \right|^\ell,$$

which implies  $\Phi \in h_1$ . Further, we get

$$\int_{\mathbb{R}} |\Phi(s)|^2 ds \leq a^2 \int_{\mathbb{R}} |s|^{-2\ell} ds. \quad (2.7)$$

When  $\ell > \frac{1}{2}$ , it is easily seen that

$$\|\Phi\|_2 < \infty, \quad i.e., \quad \Phi \in L^2(\mathbb{R}).$$

From Definitions 2.2, 2.3 and 2.4,  $H_2^* = \hat{H} \cap h_1$ ,  $H_3^* = \hat{H} \cap h_2 \cap h_3$ , hence  $H_3^* \subset H_2^* \subset H_1^* \subset H \cap L^2$ , further,  $H_3 \subset H_2 \subset H_1$ .

**Definition 2.5.** Let  $\psi, g \in L^2(\mathbb{R})$ , then their convolution is defined by

$$\psi * g(\tau) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \psi(t)g(\tau - t)dt. \quad (2.8)$$

Obviously, we have

$$\psi * g(\tau) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \psi(t)g(\tau - t)dt = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} g(t)\psi(\tau - t)dt = g * \psi(\tau), \quad (2.9)$$

this implies, the convolution is commutative. From the Hölder inequality, we know that  $\psi * g \in L^2(\mathbb{R})$ . By the convolution theorem [5,20], we have

$$\mathcal{W}[\psi * g(\tau)] = \underline{\mathcal{W}[\psi(\tau)] \cdot \mathcal{W}[g(\tau)]} = \Phi(s)G(s). \quad (2.10)$$

**Definition 2.6.** Denote the Cauchy principal integral operator  $\mathcal{V}$  as follow

$$\mathcal{V}\psi(\tau) = P.V. \frac{1}{\pi i} \int_{\mathbb{R}} \frac{\psi(\tau)}{\tau - t} d\tau = \lim_{\substack{\varepsilon \rightarrow +0 \\ X \rightarrow +\infty}} \frac{1}{\pi i} \int_{[-X, X] \setminus (t-\varepsilon, t+\varepsilon)} \frac{\psi(\tau)}{\tau - t} d\tau. \quad (2.11)$$

The operator  $\mathcal{V}$  is a self-mapping under the modified notion of Cauchy principal value integrals. Moreover, in view of the Poincaré-Bertrand formula and the Riesz theorem [11,21], we know that  $\mathcal{V}$  is an involution in  $L^2(\mathbb{R})$ , namely,  $\mathcal{V}^2 = \mathcal{I}$ , where  $\mathcal{I}$  is a unit operator.

**Definition 2.7.** We denote the operators  $\mathcal{A}, \mathcal{K}$  as follows

$$\mathcal{A}\psi(\tau) = \psi(-\tau), \quad \mathcal{K}\psi(\tau) = \psi(\tau)\text{sgn}(\tau).$$

Obviously, we have  $\mathcal{A}^2 = \mathcal{K}^2 = \mathcal{I}$ .

**Lemma 2.1.** For the function  $\psi(\tau) = \mathcal{W}^{-1}[\Phi(s)]$ , we can get

$$\mathcal{W}\left[\frac{1}{\pi i} \int_{\mathbb{R}} \frac{\psi(\tau)}{\tau - t} d\tau\right] = -\Phi(s)\text{sgn}(s), \quad s \in \mathbb{R}. \quad (2.12)$$

**Proof.** Denote Cauchy type integral  $\psi(z)$  as

$$\psi(z) = \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-z} dt, \quad z \in \mathbb{C} \setminus \mathbb{R}. \quad (2.13)$$

Substituting (2.2) into (2.13), it follows that

$$\psi(z) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \Phi(s) ds \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{e^{-ist}}{t-z} dt.$$

From the generalized residue theorem [12], for  $z \in \mathbb{C}^+$ , we have

$$\psi^+(z) = \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-z} dt = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}^-} \Phi(s) e^{-isz} ds, \quad (2.14)$$

and for  $z \in \mathbb{C}^-$ , we have

$$\psi^-(z) = \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-z} dt = \frac{-1}{\sqrt{2\pi}} \int_{\mathbb{R}^+} \Phi(s) e^{-isz} ds. \quad (2.15)$$

By the Sokhotski-Plemelj formula [5], we obtain

$$\mathcal{V}\psi(\tau) = \psi^+(t) + \psi^-(t) = \frac{-1}{\sqrt{2\pi}} \int_{\mathbb{R}} \Phi(s) \text{sgn}(s) e^{-ist} ds = \mathcal{W}^{-1}[\mathcal{K}\Phi(s)],$$

where

$$\psi^+(t) = \lim_{\substack{z \rightarrow t \\ z \in \mathbb{C}^+}} \psi^+(z), \quad \psi^-(t) = \lim_{\substack{z \rightarrow t \\ z \in \mathbb{C}^-}} \psi^-(z). \quad \square$$

**Lemma 2.2.** For the operators  $\mathcal{W}, \mathcal{A}, \mathcal{V}, \mathcal{K}$ , we have

$$(1) \mathcal{W}\mathcal{K}\mathcal{V} + \mathcal{V}\mathcal{K}\mathcal{W} = 0; \quad (2) \mathcal{V}\mathcal{W}\mathcal{A} - \mathcal{A}\mathcal{V}\mathcal{W} = 0.$$

**Proof.** (1) Note that, we may write (2.12) in the following form

$$\mathcal{W}\mathcal{V} = -\mathcal{K}\mathcal{W}.$$

Similarly, we can verify that

$$\mathcal{W}\mathcal{K} = \mathcal{V}\mathcal{W}, \quad (2.16)$$

hence

$$\mathcal{W}\mathcal{K}\mathcal{V} = \mathcal{V}\mathcal{W}\mathcal{V} = -\mathcal{V}\mathcal{K}\mathcal{W}.$$

(2) From (2.1) and (2.2), we know that

$$\mathcal{W}[\Phi(s)] = \underline{\mathcal{W}^{-1}[\Phi(-s)]} = \psi(-\tau), \quad \mathcal{W}^{-1}[\psi(\tau)] = \mathcal{W}[\psi(-\tau)] = \Phi(-s),$$

which implies

$$\mathcal{W}^{-1} = \mathcal{A}\mathcal{W}, \quad \mathcal{A}^{-1} = \mathcal{W}^2, \quad \underline{\mathcal{K}\mathcal{A} - \mathcal{A}\mathcal{K} = 0}, \quad (2.17)$$

it gives rise to

$$\mathcal{A}\mathcal{V}\mathcal{W} = \mathcal{A}\mathcal{W}\mathcal{K} = \mathcal{A}\mathcal{W}^3\mathcal{A}\mathcal{K} = \mathcal{W}\mathcal{K}\mathcal{A} = \underline{\mathcal{V}\mathcal{W}\mathcal{A}}. \quad \square$$

**Lemma 2.3.** If  $\psi \in H_1$ , and  $\mathcal{W}\psi(0) = 0$ , we have  $\mathcal{W}\mathcal{V}\psi \in H_1^*$ , further  $\mathcal{V}\psi \in H_1$ .

**Proof.** By assumption, we have  $\mathcal{W}\psi = \Phi \in H_1^*$ . Notice that

$$\lim_{s \rightarrow \infty} \Phi(s) = 0,$$

and  $\Phi(0) = 0$ , one has  $\mathcal{K}\Phi \in \hat{H}$  and

$$\int_{\mathbb{R}} |\Phi(s) \text{sgn}s|^2 ds < \infty, \quad \text{i.e., } \mathcal{K}\Phi \in L^2(\mathbb{R}),$$

thus we get  $\mathcal{W}\mathcal{V}\psi \in H_1^*$ , further,  $\mathcal{V}\psi \in H_1$ .  $\square$

**Lemma 2.4.** (see [22-24]). If  $\psi, g \in H_1(H_2, H_3)$ , then  $\psi * g \in H_1(H_2, H_3)$ , thus we have  $\Phi G \in H_1^*(H_2^*, H_3^*)$ ; if  $\psi \in H_1, g \in H_2(H_3)$ , then  $\psi * g \in H_2(H_3)$ , further,  $\Phi G \in H_2^*(H_3^*)$ .

**Lemma 2.5.** Let  $\Phi \in \hat{H}$ , we denote the Cauchy type integral as follows

$$\Phi(z) = \int_{\mathbb{R}} \frac{\Phi(s)}{s - z} ds, \quad z \notin \mathbb{R}, \quad (2.18)$$

then we can get

$$\mathcal{W}[\psi^+(\tau)] = \Phi^+(s), \quad \mathcal{W}[\psi^-(\tau)] = \Phi^-(s), \quad \Phi^+(s) - \Phi^-(s) = \Phi(s), \quad (2.19)$$

where

$$\psi^+(\tau) = \begin{cases} \psi(\tau), & \tau \geq 0; \\ 0, & \tau < 0, \end{cases} \quad \psi^-(\tau) = \begin{cases} 0, & \tau \geq 0; \\ -\psi(\tau), & \tau < 0. \end{cases}$$

**Proof.** From (2.16) and Sokhotski-Plemelj formula, we have

$$\Phi^+(s) = \frac{1}{2} \mathcal{W}[\psi(\tau)] + \frac{1}{2} \mathcal{W}[\mathcal{K}\psi(\tau)] = \frac{1}{2\sqrt{2\pi}} \int_{\mathbb{R}} \psi(\tau)(1 + \text{sgn}(\tau))e^{is\tau} d\tau = \mathcal{W}[\psi^+(\tau)],$$

for the negative boundary value, we have

$$\Phi^-(s) = \frac{-1}{2} \mathcal{W}[\psi(\tau)] + \frac{1}{2} \mathcal{W}[\mathcal{K}\psi(\tau)] = \frac{1}{2\sqrt{2\pi}} \int_{\mathbb{R}} \psi(\tau)(\text{sgn}(\tau) - 1)e^{is\tau} d\tau = \mathcal{W}[\psi^-(\tau)].$$

Further,

$$\Phi^+(s) - \Phi^-(s) = \mathcal{W}[\psi^+(\tau)] - \mathcal{W}[\psi^-(\tau)] = \mathcal{W}[\psi(\tau)] = \Phi(s). \quad \square$$

From Lemma 2.5, we know that the positive and negative boundary values  $\Phi^\pm(s)$  of  $\Phi(z)$  are the single sided Fourier integral transforms of  $\psi^\pm(\tau)$ , respectively.

### 3 SIEs with two convolution kernels

We solve the following SIEs with two convolution kernels and variable integral limits

$$A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_0^\tau \psi(t)h(\tau - t)dt + \frac{C}{\sqrt{2\pi}} \int_\tau^0 \psi(t)k(\tau - t)dt + \frac{D}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t - \tau} dt = n(\tau), \quad \tau \in \mathbb{R}, \quad (3.1)$$

where  $A, B, C, D$  are constants, and  $D \neq 0$ . The functions  $h, k, n \in H_1$ , and the unknown function  $\psi \in H_1$ .

Expanding  $t$  to  $t \in \mathbb{R}$ , then Eq. (3.1) can be transformed into

$$A\psi(\tau) + \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \psi(t) h_1(\tau-t) \operatorname{sgn}(t) dt + \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \psi(t) k_1(\tau-t) dt + \frac{D}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau), \quad \tau \in \mathbb{R}, \quad (3.2)$$

where

$$h_1(\tau) = \frac{B}{2} h^+(\tau) + \frac{C}{2} k^-(\tau), \quad k_1(\tau) = \frac{B}{2} h^+(\tau) - \frac{C}{2} k^-(\tau),$$

in which

$$h^\pm(\tau) = \frac{\operatorname{sgn}(\tau) \pm 1}{2} h(\tau), \quad k^\pm(\tau) = \frac{\operatorname{sgn}(\tau) \pm 1}{2} k(\tau),$$

that is to say,

$$h_1(\tau) = \begin{cases} \frac{B}{2} h(\tau), & \tau \geq 0; \\ -\frac{C}{2} k(\tau), & \tau < 0, \end{cases} \quad k_1(\tau) = \begin{cases} \frac{B}{2} h(\tau), & \tau \geq 0; \\ \frac{C}{2} k(\tau), & \tau < 0. \end{cases}$$

Applying the Fourier transforms to both sides of (3.2), then we have

$$A\Phi(s) + \frac{H(s)}{\pi i} \int_{\mathbb{R}} \frac{\Phi(t)}{t-s} dt + K(s)\Phi(s) - D\operatorname{sgn}(s)\Phi(s) = N(s), \quad (3.3)$$

where

$$\mathcal{W}[h_1(\tau)] = H(s), \quad \mathcal{W}[k_1(\tau)] = K(s), \quad \mathcal{W}[n(\tau)] = N(s), \quad \mathcal{W}[\psi(\tau)] = \Phi(s).$$

Denote the Cauchy type integral

$$\Phi(z) = \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{\Phi(t)}{t-z} dt, \quad z \in \mathbb{C} \setminus \mathbb{R}, \quad (3.4)$$

by the Sokhotski-Plemelj formula, we can get

$$\Phi^+(s) - \Phi^-(s) = \Phi(s), \quad \Phi^+(s) + \Phi^-(s) = \frac{1}{\pi i} \int_{\mathbb{R}} \frac{\Phi(t)}{t-s} dt. \quad (3.5)$$

Substituting (3.5) into (3.3), then we shall solve the following Riemann problem in place of (3.3)

$$\Phi^+(s) = J(s)\Phi^-(s) + N_0(s), \quad s \in \mathbb{R}, \quad (3.6)$$

where

$$J(s) = \frac{A - H(s) + K(s) - D\operatorname{sgn}(s)}{A + H(s) + K(s) - D\operatorname{sgn}(s)}, \quad N_0(s) = \frac{N(s)}{A + H(s) + K(s) - D\operatorname{sgn}(s)}.$$

Moreover, we can write  $J(s), N_0(s)$  in the forms

$$J(s) = \begin{cases} \frac{A-H(0)+K(0)}{A+H(0)+K(0)}, & s = 0; \\ 1 - \frac{2H(s)}{A+H(s)+K(s)-D}, & s \in \mathbb{R}^+; \\ 1 - \frac{2H(s)}{A+H(s)+K(s)+D}, & s \in \mathbb{R}^-, \end{cases} \quad N_0(s) = \begin{cases} \frac{N(0)}{A+H(0)+K(0)}, & s = 0; \\ \frac{N(s)}{A+H(s)+K(s)-D}, & s \in \mathbb{R}^+; \\ \frac{N(s)}{A+H(s)+K(s)+D}, & s \in \mathbb{R}^-. \end{cases}$$

Since  $H, K, N \in H_1^*$ , then we know that

$$\lim_{s \rightarrow \infty} H(s) = \lim_{s \rightarrow \infty} K(s) = \lim_{s \rightarrow \infty} N(s) = 0,$$

which implies  $\lim_{s \rightarrow \infty} J(s) = 1$ . In this case,  $s = \infty$  is not a node of (3.6).

Let  $J_1(s) = A + H(s) + K(s) - D \operatorname{sgn}(s)$  have some zero-points  $d_1, d_2, \dots, d_m$  with the orders  $\alpha_1, \alpha_2, \dots, \alpha_m$  respectively; let  $J_2(s) = A - H(s) + K(s) - D \operatorname{sgn}(s)$  have some zero-points  $e_1, e_2, \dots, e_n$  with the orders  $\beta_1, \beta_2, \dots, \beta_n$  respectively, where  $d_i \neq e_j$ ,  $\alpha_i, \beta_j$  ( $i = 1, 2, \dots, m; j = 1, 2, \dots, n$ ) are non-negative integers. Let

$$E_1(s) = \prod_{i=1}^m (s - d_i)^{\alpha_i}, \quad E_2(s) = \prod_{j=1}^n (s - e_j)^{\beta_j},$$

and

$$\alpha_1 + \alpha_2 + \dots + \alpha_m = M_1, \quad \beta_1 + \beta_2 + \dots + \beta_n = M_2.$$

Next, we only consider the case of non-normal type, that is,  $M_1 > 0, M_2 > 0$ .

Without loss of generality, we take any fixed points  $z_0 = a + ib \in \mathbb{C}^+$ , and  $z_0^* = a - ib \in \mathbb{C}^-$ , and rewrite (3.6) as

$$\Phi^+(s) = \frac{E_2(s)(s - z_0^*)^{M_1}}{E_1(s)(s - z_0)^{M_2}} P(s) \Phi^-(s) + N_0(s), \quad s \in \mathbb{R}, \quad (3.7)$$

where

$$J(s) = \frac{E_2(s)(s - z_0^*)^{M_1}}{E_1(s)(s - z_0)^{M_2}} P(s), \quad P(s) \neq 0,$$

and  $P \in H$ . Note that, the solution  $\Phi(s)$  of (3.6) is continuous along  $\mathbb{R}$ , and  $\lim_{s \rightarrow \infty} \Phi(s) = 0$ .

Denote

$$\gamma_0 = \lambda_0 + i\eta_0, \quad \lambda_0 = \frac{1}{2\pi} \arg \frac{P(-0)}{P(+0)}, \quad \eta_0 = -\frac{1}{2\pi} \ln \left| \frac{P(-0)}{P(+0)} \right|, \quad (3.8)$$

and we say  $\kappa$  the index of (3.7) which satisfies  $\kappa = [\lambda_0]$ . Let

$$\lambda = \lambda_0 - \kappa, \quad \gamma = \gamma_0 - \kappa,$$

hence  $\gamma = \lambda + i\eta_0, \lambda \in [0, 1)$ .

In order to prove the solvability of (3.6), we first introduce the sectionally holomorphic function

$$X(z) = \begin{cases} (z - z_0^*)^{-\kappa} e^{\Gamma_0(z)}, & \operatorname{Im} z > 0; \\ (z - z_0)^{-\kappa} e^{\Gamma_0(z)}, & \operatorname{Im} z < 0, \end{cases} \quad (3.9)$$

in which we have put

$$\Gamma_0(z) = \frac{1}{2\pi i} \int_{\mathbb{R}} \ln P_0(s) \frac{ds}{s - z}, \quad z \in \mathbb{C} \setminus \mathbb{R}, \quad (3.10)$$

and

$$P_0(s) = \left( \frac{s - z_0^*}{s - z_0} \right)^{\kappa} P(s), \quad P(s) = \frac{X^+(s)}{X^-(s)}.$$



Obviously,

$$\kappa \ln \frac{s - z_0^*}{s - z_0} = \ln P_0(s) - \ln P(s), \quad (3.11)$$

where  $\ln P_0 \in \hat{H}$ , and we take a continuous single-valued branch of  $\ln \frac{s - z_0^*}{s - z_0}$  which satisfies

$$\ln \frac{\pm\infty - z_0^*}{\pm\infty - z_0} = 0, \quad \text{or} \quad \ln \frac{\pm 0 - z_0^*}{\pm 0 - z_0} = \pm\pi i.$$

Therefore, we rewrite (3.7) in the form

$$\Phi^+(s) = \frac{E_2(s)(s - z_0^*)^{M_1} X^+(s)}{E_1(s)(s - z_0)^{M_2} X^-(s)} \Phi^-(s) + N_0(s), \quad (3.12)$$

for convenience, we deal with (3.12) in the problem  $R_{-1}$ , that is,  $\Phi(\infty) = 0$ .

For homogeneous problem of (3.12), we consider the function

$$\Omega(z) = \begin{cases} \frac{\Phi(z)}{X(z)E_2(z)}(z - z_0^*)^{-M_1}, & \text{Im}z > 0; \\ \frac{\Phi(z)}{X(z)E_1(z)}(z - z_0)^{-M_2}, & \text{Im}z < 0. \end{cases} \quad (3.13)$$

From the principle of analytic continuation, we know that  $\Omega(z)$  is analytic in  $\mathbb{C}$ . Moreover, by the generalized Liouville theorem, we have

$$\Omega(z) = U_{\kappa - M_1 - M_2 - 1}(z) = \sum_{t=0}^{\kappa - M_1 - M_2 - 1} C_{\kappa - M_1 - M_2 - 1 - t} z^t,$$

where  $C_0, \dots, C_{\kappa - M_1 - M_2 - 1}$  are arbitrary complex constants. Therefore, the analytic solution of (3.12) is given by

$$\Phi_1(z) = \begin{cases} E_2(z)(z - z_0^*)^{M_1} X(z) U_{\kappa - M_1 - M_2 - 1}(z), & \text{Im}z > 0; \\ E_1(z)(z - z_0)^{M_2} X(z) U_{\kappa - M_1 - M_2 - 1}(z), & \text{Im}z < 0. \end{cases} \quad (3.14)$$

When  $\kappa \leq M_1 + M_2$ , we have  $\Omega(z) = 0$ , thus  $\Phi_1(z) \equiv 0$ .

To solve the non-homogeneous problem (3.12), we introduce the following function

$$Y(z) = \begin{cases} X_0(z), & \text{Im}z > 0; \\ (z - z_0^*)^\kappa (z - z_0)^{-\kappa} X_0(z), & \text{Im}z < 0, \end{cases} \quad (3.15)$$

where  $X_0(z) = e^{\Gamma_0(z)}$ , then (3.12) is written as

$$\frac{\Phi^+(s)E_1(s)}{Y^+(s)(s - z_0^*)^{M_1}} = \frac{\Phi^-(s)E_2(s)}{Y^-(s)(s - z_0)^{M_2}} + \frac{N_0(s)E_1(s)}{Y^+(s)(s - z_0^*)^{M_1}}. \quad (3.16)$$

Define

$$\Psi(z) = \frac{1}{2\pi i} \int_{\mathbb{R}} \Psi^*(s) \frac{ds}{s - z}, \quad z \in \mathbb{C} \setminus \mathbb{R}, \quad (3.17)$$

where

$$\Psi^*(s) = \frac{N_0(s)E_1(s)}{Y^+(s)(s - z_0^*)^{M_1}}, \quad \Psi^* \in \hat{H}.$$

Via the generalized Liouville theorem, we know that

$$\Psi^+(s) - \Psi^-(s) = \Psi^*(s).$$

Again define

$$W(z) = \begin{cases} \frac{\Phi(z)E_1(z)}{Y(z)(z-z_0^*)^{M_1}} - \Psi(z), & \text{Im}z > 0; \\ \frac{\Phi(z)E_2(z)}{Y(z)(z-z_0)^{M_2}} - \Psi(z), & \text{Im}z < 0, \end{cases} \quad (3.18)$$

obviously,  $W(z)$  is analytic in  $\mathbb{C} \setminus \{z_0^*\}$ , thus we can obtain the solution of (3.12) with singularities at  $d_i$  and  $e_j$ . Hence, we shall construct a Hermite interpolation polynomial

$$\Omega_\varrho(z) = b_0 z^\varrho + \cdots + b_{\varrho-1} z + b_\varrho, \quad (3.19)$$

where  $\varrho = M_1 + M_2 - 1$ , and  $b_1, \dots, b_\varrho \in \mathbb{C}$ . Note that,  $\Omega_\varrho(z)$  is uniquely determined, and it has zero-points of the orders  $\alpha_i (i = 1, 2, \dots, m)$ ,  $\beta_j (j = 1, 2, \dots, n)$  at  $d_i, e_j$ , respectively, which implies

$$\begin{aligned} [\Psi(z)(z - z_0^*)^\kappa]^{(L_1)} \Big|_{z=d_i} &= \Omega_\varrho^{(L_1)}(z) \Big|_{z=d_i}; \\ [\Psi(z)(z - z_0^*)^\kappa]^{(L_2)} \Big|_{z=e_j} &= \Omega_\varrho^{(L_2)}(z) \Big|_{z=e_j}, \end{aligned}$$

where

$$L_1 = 1, 2, \dots, \alpha_i - 1, \quad i = 1, 2, \dots, m; \quad L_2 = 1, 2, \dots, \beta_j - 1, \quad j = 1, 2, \dots, n.$$

By the Riemann boundary value theory, (3.12) has the particular solution

$$\Phi_2(z) = \begin{cases} \frac{Y(z)(z-z_0^*)^{M_1}}{E_1(z)(z-z_0^*)^\kappa} [\Psi(z)(z - z_0^*)^\kappa - \Omega_\varrho(z)], & \text{Im}z > 0; \\ \frac{Y(z)(z-z_0)^{M_2}}{E_2(z)(z-z_0^*)^\kappa} [\Psi(z)(z - z_0^*)^\kappa - \Omega_\varrho(z)], & \text{Im}z < 0, \end{cases} \quad (3.20)$$

where

$$(z - z_0^*)^\kappa X(z) = Y(z), \quad z \in \mathbb{C} \setminus \mathbb{R}.$$

In view of linearity, we obtain the general solution of (3.12) as follows

$$\Phi(z) = \begin{cases} X(z)(z - z_0^*)^{M_1} E_1^{-1}(z) F(z), & \text{Im}z > 0; \\ X(z)(z - z_0^*)^{M_2} E_2^{-1}(z) F(z), & \text{Im}z < 0, \end{cases} \quad (3.21)$$

where

$$F(z) = \Psi(z)(z - z_0^*)^\kappa - \Omega_\varrho(z) + E_1(z)E_2(z)U_{\kappa-M_1-M_2-1}(z).$$

Obviously,  $\Phi_2(z)$  has the singularity at  $z = z_0^*$  when  $\kappa < 0$ . Due to  $\Phi \in H_1^*$ , we must have

$$\int_{\mathbb{R}} \Psi^*(s) \frac{ds}{(s - z_0^*)^\nu} = 0, \quad \nu = 1, 2, \dots, |\kappa|. \quad (3.22)$$

Taking the boundary values of  $Y(z)$ , then we obtain

$$Y^+(s) = \sqrt{P_0(s)} X_0(s), \quad Y^-(s) = \frac{X_0(s)}{\sqrt{P_0(s)}}, \quad (3.23)$$

hence we have

$$\Phi^+(s) = \frac{N_0(s)}{2} + \frac{X_0(s)\sqrt{P_0(s)}}{E_1(s)(s-z_0^*)^{\kappa-M_1}}F_*(s),$$

and

$$\Phi^-(s) = -\frac{N_0(s)E_1(s)(s-z_0^*)^{M_2-M_1}}{2E_2(s)P_0(s)} + \frac{X_0(s)(s-z_0^*)^{M_2-\kappa}}{\sqrt{P_0(s)}E_2(s)}F_*(s),$$

where

$$F_*(s) = F(s) - \Omega_\rho(s) + E_1(s)E_2(s)U_{\kappa-M_1-M_2-1}(s).$$

By (2.20) we get the following closed-form solution

$$\begin{aligned} \Phi(s) &= \frac{N_0(s)}{2} + \frac{N_0(s)(s-z_0^*)^{M_2-M_1}E_1(s)}{2P_0(s)E_2(s)} \\ &+ \frac{X_0(s)F_*(s)}{(s-z_0^*)^\kappa\sqrt{P_0(s)}} \left( \frac{P_0(s)}{E_1(s)}(s-z_0^*)^{M_1} - \frac{(s-z_0^*)^{M_2}}{E_2(s)} \right). \end{aligned} \quad (3.24)$$

Next, we consider the properties of the solutions at  $s = 0$ .

Suppose that  $s = 0$  is an ordinary node, that is,  $\lambda \in (0, 1)$ , then  $\gamma \neq 0$ . Since

$$\lim_{s \rightarrow 0^+} \Phi(s) = \lim_{s \rightarrow 0^-} \Phi(s) = \Phi(0),$$

then by (3.23) we can prove that

$$Y^+(s) = s^\gamma \sqrt{P_0(s)} e^{\Gamma_*(s)}, \quad Y^-(s) = \frac{s^\gamma e^{\Gamma_*(s)}}{\sqrt{P_0(s)}}, \quad (3.25)$$

where  $\Gamma_*(s) = \Gamma_0(s) - \gamma \ln s$ , and  $\Gamma_* \in H$ . From (3.9) and (3.10), we have

$$\ln P_0(\pm 0) = \ln P(\pm 0) + \kappa \ln \frac{\pm 0 - z_0^*}{\pm 0 - z_0} = \ln P(\pm 0) \pm \kappa \pi i, \quad (3.26)$$

again from (3.9) and (3.26), we obtain

$$\sqrt{P_0(+0)} = e^{\frac{\kappa}{2}\pi i} \sqrt{P(+0)} = e^{\kappa \pi i} e^{\frac{1}{2} \ln \frac{P(+0)}{P(-0)}} e^{-\frac{\kappa \pi i}{2}} e^{\frac{1}{2} \ln P(-0)} = e^{-\gamma \pi i} \sqrt{P_0(-0)}. \quad (3.27)$$

According to [5-7,13,25], when  $s > 0$ , it is clear that

$$\Psi(s) = \frac{e^{-\gamma \pi i} s^{-\gamma}}{2i e^{\Gamma_*(s)}} \Lambda_0 + \Psi_0(s), \quad (3.28)$$

where

$$\Lambda_0 = e^{\gamma \pi i} \Lambda(+0) \cot \gamma \pi - \Lambda(-0) \csc \gamma \pi, \quad \Lambda(s) = \frac{N_0(s)E_1(s)}{\sqrt{P_0(s)}(s-z_0^*)^{M_1}},$$

and  $\Psi_0(s)|s|^{\lambda^*} \in H$ ,  $\lambda^* \in (0, \lambda)$ . Therefore, substituting (3.25)-(3.28) into  $\Phi^+(s)$ , one has

$$\Phi^+(+0) = \frac{N_0(+0) \csc \gamma \pi}{2i e^{2\gamma \pi i}} \left[ e^{3\gamma \pi i} - \frac{N_0(-0)}{N_0(+0)} \right]. \quad (3.29)$$

As in [4,13,26], when  $s < 0$ , we have

$$\Psi(s) = \frac{e^{\gamma\pi i} s^{-\gamma}}{2ie^{\Gamma_*(s)}} \Lambda_1 + \Psi_0(s), \quad (3.30)$$

where

$$\Lambda_1 = -\cot \gamma\pi e^{-\gamma\pi i} \Lambda(-0) + \Lambda(+0) \csc \gamma\pi,$$

it gives rise to

$$\Phi^+(-0) = \frac{N_0(+0) \csc \gamma\pi}{2ie^{\gamma\pi i}} \left[ e^{3\gamma\pi i} - \frac{N_0(-0)}{N_0(+0)} \right]. \quad (3.31)$$

By comparing (3.29) with (3.31), and  $e^{\gamma\pi i} \neq 1$ , we know that  $\Phi^+(+0) = \Phi^+(-0)$  if and only if

$$e^{3\gamma\pi i} N_0(+0) = N_0(-0). \quad (3.32)$$

Since  $\Phi(s)$  is continuous at  $s = 0$ , then we have

$$\Phi^-(0) = \Phi^+(0) = 0,$$

and  $\Phi(0) = 0$ . Further,  $N_0(0) = 0$ , and so

$$N(0) = 0. \quad (3.33)$$

If (3.33) is valid, for  $s \in N_0 = \{|s| < \epsilon, \forall \epsilon > 0\}$ , we have  $\Phi^\pm \in H_1^*$ .

Suppose that  $s = 0$  is a special node, then  $\lambda = 0$ ,  $\gamma = i\eta_0$ . We have  $\Psi_0 \in H$  near  $s = 0$ . If  $\eta_0 \neq 0$ , from [5-7,22], (3.29) and (3.31) are still fulfilled, and  $\Psi_0(\pm 0)$  exist and may not be equal. Return to (3.29) and (3.31), we should modify  $\Phi^+(\pm 0)$  to

$$\Phi^+(+0) = \frac{N_0(+0) \csc \gamma\pi}{2ie^{2\gamma\pi i}} \left[ e^{3\gamma\pi i} - \frac{N_0(-0)}{N_0(+0)} \right] + \sigma(0) \sqrt{P_0(+0)} \lim_{s \rightarrow +0} s^{i\eta_0} [\Psi_0(s) - \Lambda], \quad (3.34)$$

and

$$\Phi^+(-0) = \frac{N_0(+0) \csc \gamma\pi}{2ie^{\gamma\pi i}} \left[ e^{3\gamma\pi i} - \frac{N_0(-0)}{N_0(+0)} \right] + \sigma(0) \sqrt{P_0(-0)} \lim_{s \rightarrow -0} s^{i\eta_0} [\Psi_0(s) - \Lambda], \quad (3.35)$$

where

$$\sigma(s) = \frac{e^{\Gamma_*(s)}}{E_1(s)} (s - z_0^*)^{M_1},$$

and when  $\kappa > \varrho + 1$ ,

$$\Lambda = (-1)^\kappa \frac{b_\varrho - E_1(0)E_2(0)C_{\kappa-M_1-M_2-1}}{(z_0^*)^\kappa}.$$

Since  $\Phi^+(+0) = \Phi^+(-0)$ , and  $\Psi_0(s)$  is continuous at  $s = 0$ , then we have

$$\Psi_0(\pm 0) = \Lambda,$$

in this case, (3.32) holds, and so (3.33) holds. On the other hand, if (3.33) is valid, thus  $\Phi(s)$  is continuous at  $s = 0$ . When  $\kappa > \varrho + 1$ , we have the condition of solvability  $F(0) = 0$ , that is to say,

$$b_\varrho = E_1(0)E_2(0)C_{\kappa-M_1-M_2-1} + (-z_0^*)^\kappa \Psi(0); \quad (3.36)$$

when  $\kappa \leq \varrho + 1$ , the constant  $b_\varrho$  takes the value

$$b_\varrho = \frac{(-z_0^*)^\kappa}{2\pi i} \int_{\mathbb{R}} \frac{\Psi^*(s)}{s} ds. \quad (3.37)$$

If  $\eta_0 = 0$ , then  $\gamma = 0$ , therefore  $J(+0) = J(-0)$ . We can also prove that  $\Phi(0) = 0$  if and only if (3.33) is valid. Under the condition (3.33), we have  $\Phi \in H$ , and  $\Phi(0) = 0$ . Moreover, when  $\kappa < \varrho + 1$ , we have the following conditions of solvability

$$b_0 = \cdots = b_{\varrho-\kappa} = 0, \quad (3.38)$$

then  $\Omega_\varrho(z)$  is a polynomial with the degree  $\kappa - 1$ . When  $\kappa = 1$ ,  $\Omega_\varrho(z) = b_\varrho$ . In this case, we require that

$$\Psi(s)(s - z_0^*)|_{s=d_i} = b_\varrho; \quad \Psi(s)(s - z_0^*)|_{s=e_j} = b_\varrho, \quad (3.39)$$

and

$$\int_{\mathbb{R}} \Psi^*(s)(s - d_i)^{-v_1-1} ds = 0; \quad \int_{\mathbb{R}} \Psi^*(s)(s - e_j)^{-v_2-1} ds = 0, \quad (3.40)$$

are satisfied, where

$$v_1 = 1, 2, \dots, \alpha_i - 1, \quad i = 1, 2, \dots, m; \quad v_2 = 1, 2, \dots, \beta_j - 1, \quad j = 1, 2, \dots, n.$$

When  $\kappa \leq 0$ ,  $\Omega_\varrho(z) = 0$ . Moreover, when  $\kappa < 0$ , in order that  $\Phi_2 \in H_1^*$ , we need (3.24) and the solvability conditions as follows

$$\int_{\mathbb{R}} \frac{\Psi^*(s)}{s - d_i} ds = 0; \quad \int_{\mathbb{R}} \frac{\Psi^*(s)}{s - e_j} ds = 0. \quad (3.41)$$

Combining (3.40) and (3.41), when  $\kappa < 0$ , we require that the following (3.42) holds

$$\int_{\mathbb{R}} \Psi^*(s)(s - d_i)^{-v_1} ds = 0; \quad \int_{\mathbb{R}} \Psi^*(s)(s - e_j)^{-v_2} ds = 0, \quad (3.42)$$

where

$$v_1 = 1, 2, \dots, \alpha_i, \quad i = 1, 2, \dots, m; \quad v_2 = 1, 2, \dots, \beta_j, \quad j = 1, 2, \dots, n.$$

Under the assumptions and solvability conditions, (3.1) has the solution

$$\psi(\tau) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \Phi(s) e^{-is\tau} ds, \quad (3.43)$$

where  $\Phi(s)$  is given by (3.24).

The results about the solutions of (3.1) are formulated in the following theorem.

**Theorem 3.1.** Under the case of non-normal type, the necessary condition of solvability for problem (3.6) is (3.33).

(1) If  $s = 0$  is an ordinary node, (3.33) holds. When  $\kappa > \varrho + 1$ , Eq. (3.1) has  $\kappa - \varrho - 1$  linearly independent solutions. When  $\kappa = \varrho + 1$ , (3.1) has a unique solution. When  $\kappa < \varrho + 1$ , (3.1) is solvable if (3.38) holds. When  $\kappa = 1$ , (3.39) and (3.40) hold, and when  $\kappa < 0$ , (3.22) and (3.42) hold.

(2) If  $s = 0$  is a special node, (3.1) is solvable if (3.33) and (3.36) are valid. The solutions of (3.1) belong to the class  $H_1$ .

**Remark 3.1.** Suppose that  $J_1(s), J_2(s)$  have common zero-points  $j_1, \dots, j_{k'}$  with the orders  $\delta_1, \dots, \delta_{k'}$  respectively, where  $j_p (1 \leq p \leq k')$  are different from  $d_i (1 \leq i \leq m)$  and  $e_j (1 \leq j \leq n)$ , the additional solvability condition should be fulfilled

$$N^{(l)}(j_p) = 0, \quad l = 1, \dots, \delta_p - 1.$$

**Remark 3.2.** SIE with a convolution kernel and variable integral limits is a special case of Eq. (3.1), i.e.,  $h(\tau) = 0$  or  $k(\tau) = 0$ . In this case, the process of analysis is not fundamentally different from (3.1), and will be omitted.

## 4 SIEs of Wiener-Hopf with convolution kernels

Next, we consider SIEs of Wiener-Hopf with convolution kernels and variable integral limits

$$A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_0^\tau \psi(t)h(\tau-t)dt + \frac{C}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau), \quad \tau \in \mathbb{R}^+, \quad (4.1)$$

where  $A, B, C$  are constants, and  $C \neq 0$ . The known functions  $h, n \in H_3^\ell$ ,  $\ell \in (\frac{1}{2}, 1)$ , and the unknown function  $\psi \in H_1$ . Extending  $\tau$  to  $\tau \in \mathbb{R}$  by adding  $\psi^-(\tau)$  to the right side of (4.1), we have

$$A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_{\mathbb{R}} \psi^+(t)h^+(\tau-t)dt + \frac{C}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau) + \psi^-(\tau), \quad \tau \in \mathbb{R}. \quad (4.2)$$

By applying the operator  $\mathcal{W}$  to (4.2), we get

$$\Phi^+(s) = J(s)\Phi^-(s) + N_0(s), \quad s \in \mathbb{R}, \quad (4.3)$$

in which

$$J(s) = \frac{1}{A + BH^+(s) - C\text{sgn}(s)}, \quad N_0(s) = \frac{N(s)}{A + BH^+(s) - C\text{sgn}(s)},$$

and

$$\mathcal{W}[h^+(\tau)] = H^+(s), \quad \mathcal{W}[n(\tau)] = N(s), \quad \mathcal{W}[\psi^\pm(\tau)] = \Phi^\pm(s), \quad N_0(s) = J(s)N(s).$$

It follows from  $H, N \in h_3$  that

$$\lim_{s \rightarrow +\infty} J(s) = \frac{1}{A - C}, \quad \lim_{s \rightarrow -\infty} J(s) = \frac{1}{A + C}, \quad \lim_{s \rightarrow \infty} N_0(s) = 0, \quad (4.4)$$

hence  $s = 0, \infty$  are nodes of problem (4.3).

Now, we give the definitions of  $\gamma_\infty$  and  $\gamma_0$ . Define

$$\gamma_\infty = \lambda_\infty + i\eta_\infty = \frac{1}{2\pi i} \ln \frac{J(-\infty)}{J(+\infty)} = \frac{1}{2\pi i} \ln \frac{A - C}{A + C}, \quad (4.5)$$

where  $\ln J(s)$  takes a continuous branch such that  $\lambda_\infty \in [0, 1)$ . From [5-7], we know that  $\gamma_\infty \neq 0$  since  $C \neq 0$ . Again denote

$$\gamma_0 = \lambda_0 + i\eta_0 = \frac{1}{2\pi i} \ln \frac{J(-0)}{J(+0)} = \frac{1}{2\pi i} \ln \frac{A + BH^+(0) - C}{A + BH^+(0) + C}. \quad (4.6)$$

We call  $\kappa$  the index of (4.3) which satisfies  $\lambda_0 - 1 < \kappa \leq \lambda_0$ . Suppose that  $J^{-1}(s)$  has some zero-points  $d_i (1 \leq i \leq m)$  with the orders  $\alpha_i (1 \leq i \leq m)$  respectively, then we put

$$E(s) = \sum_{i=1}^m (s - d_i)^{\alpha_i}, \quad M = \alpha_1 + \cdots + \alpha_m. \quad (4.7)$$

Here, we still only discuss the case  $M > 0$ . Rewrite (4.3) as

$$\Phi^+(s) = \frac{X^+(s)(s - z_0^*)^M}{X^-(s)E(s)} \Phi^-(s) + N_0(s). \quad (4.8)$$

By the principle of analytic continuation and the generalized Liouville theorem, (4.3) has the analytic solution

$$\Phi_1(z) = \begin{cases} X(z)(z - z_0^*)^M P_{\kappa-M-1}(z), & \text{Im}z > 0; \\ X(z)E(z)P_{\kappa-M-1}(z), & \text{Im}z < 0. \end{cases} \quad (4.9)$$

Next, we consider the non-homogeneous problem (4.8). Put

$$\Phi_2(z) = \begin{cases} \frac{Y(z)}{E(z)(z - z_0^*)^{\kappa-M}} [\Psi(z)(z - z_0^*)^\kappa - \Omega_\rho(z)], & \text{Im}z > 0; \\ \frac{Y(z)}{(z - z_0^*)^\kappa} [\Psi(z)(z - z_0^*)^\kappa - \Omega_\rho(z)], & \text{Im}z < 0, \end{cases} \quad (4.10)$$

where

$$\Omega_\rho(z) = b_\rho + b_{\rho-1}z + \cdots + b_0 z^\rho \quad (4.11)$$

has zero-points of the orders  $\alpha_i$  at  $d_i$ , and  $\rho = M - 1$ , this implies

$$\Omega_\rho^{(L)}(z) = [\Psi(z)(z - z_0^*)^\kappa]^{(L)}, \quad 1 \leq L \leq \alpha_i - 1, \quad 1 \leq i \leq m. \quad (4.12)$$

Using the theory of classical boundary value, we can verify that (4.10) is the particular solution of (4.3). Therefore, (4.3) has the general solution

$$\Phi(z) = \begin{cases} \frac{X(z)(z - z_0^*)^M}{E(z)} [\Psi(z)(z - z_0^*)^\kappa - \Omega_\rho(z) + E(z)P_{\kappa-M-1}(z)], & \text{Im}z > 0; \\ X(z) [\Psi(z)(z - z_0^*)^\kappa - \Omega_\rho(z) + E(z)P_{\kappa-M-1}(z)], & \text{Im}z < 0. \end{cases} \quad (4.13)$$

Further process is the same as section 3 and will be omitted.

Now, we discuss the properties of solutions at  $s = \infty$ . By [5,13,27], we have, near  $s \in N_\infty$ ,

$$Y(s)s^{\lambda_\infty} \in h_2.$$

Notice that, in (4.13), we shall write  $P_{\kappa-M}(z)$  instead of  $P_{\kappa-M-1}(z)$ .

Suppose that  $s = \infty$  is an ordinary node, then  $\lambda_\infty \in (0, 1)$ , further  $\gamma_\infty \neq 0$ . Consider the positive boundary value  $\Phi^+(s)$  as follow

$$\Phi^+(s) = Y^+(s)[\Psi(s)Q_1(s) + (s - z_0^*)^{M-\kappa}Q_2(s)] + \frac{N_0(s)}{2}, \quad (4.14)$$

where

$$Q_1(s) = (s - z_0^*)^M E^{-1}(s), \quad Q_2(s) = P_{\kappa-M}(s) - \Omega_\rho(s)E^{-1}(s). \quad (4.15)$$

We can see that  $Q_1(s)$  is bounded near  $s \in N_\infty$ . Moreover, when  $\kappa \geq M$ , one has

$$Q_2(s) = O(|s - z_0^*|^{\kappa-M}), \quad Y^+(s)Q_2(s)(s - z_0^*)^{M-\kappa} = O(|s|^{-\lambda_\infty}). \quad (4.16)$$

When  $\frac{1}{2} < \lambda_\infty < \ell < 1$ , we have  $|\Psi^*(s)| \leq r_0$  near  $s \in N_\infty$ , then

$$|\Psi(s)| \leq \frac{r_0}{2\pi} \int_{\mathbb{R}} \left| \frac{1}{s} \right| \left| 1 - \frac{t}{s} \right|^{-1} |dt| = \frac{r_0}{2\pi} \int_{\mathbb{R}} \left| \sum_{n=0}^{\infty} \frac{t^n}{s^{n+1}} \right| |dt| \leq r_0^*,$$

where  $r_0, r_0^*$  take constants. Further, we obtain

$$Y^+(s)\Psi(s) = O(|s|^{-\lambda_\infty}), \quad (4.17)$$

and by (4.15)-(4.17), when  $\kappa \geq M$ , it gives rise to

$$\Phi^+(s) = O(|s|^{-\lambda_\infty}). \quad (4.18)$$

When  $\kappa < M$ ,  $P_{\kappa-M}(z) = 0$ , then we require the solvability conditions as follows

$$b_j = 0, \quad j = 0, \dots, \rho - \kappa - 1. \quad (4.19)$$

In this case,  $\Omega_\rho(z) = \sum_{j=0}^{\kappa} b_{\rho-j} z^j$ . Moreover, when  $\kappa > 0$ , (4.18) is still valid; when  $\kappa = 0$ ,  $\Omega_\rho(z) = b_\rho$ , one must have

$$\Psi(d_i)(d_i - z_0^*) = b_\rho, \quad 1 \leq i \leq m; \quad (4.20)$$

when  $\kappa < 0$ , we need that (3.22) and the following (4.21) hold

$$\int_{\mathbb{R}} \Psi^*(s)(s - d_i)^{-v} ds = 0, \quad (4.21)$$

where  $1 \leq v \leq \alpha_i, 1 \leq i \leq m$ .

When  $\frac{1}{2} < \ell \leq \lambda_\infty < 1$ , there exists  $\epsilon > 0$  such that  $\lambda_\infty - \epsilon > \frac{1}{2}$ . Therefore, we have

$$Y^+(s)\Psi(s) = O(|s|^{-\lambda_\infty + \epsilon}), \quad s \in N_\infty. \quad (4.22)$$

Similarly, we can get

$$\Phi^+(s) = O(|s|^{-\lambda_\infty + \epsilon}), \quad s \in N_\infty. \quad (4.23)$$

Combining (4.18) and (4.23), when  $\lambda_\infty > \frac{1}{2}$ , it turns out that

$$\Phi^+(s) = O(|s|^{-r}), \quad s \in N_\infty, \quad (4.24)$$



where  $r > \min \{\ell, \lambda_\infty - \epsilon\}$ , thus  $r > \frac{1}{2}$ . Moreover,  $\Phi^+ \in H_2^*$ .

When  $0 < \lambda_\infty \leq \frac{1}{2}$ , by [13,24] we know that

$$Y^+(s)\Psi(s) = O(|s|^{-\ell}).$$

When  $\kappa \geq M$ , in order that  $\Phi^+ \in L^2(\mathbb{R})$ , (4.18) needs to be satisfied. When  $\kappa < M$ , (4.19) holds. Further, when  $\kappa \geq 0$ , we must have

$$b_{\rho-\kappa} = 0; \tag{4.25}$$

when  $\kappa < 0$ , we require that (4.19) and (4.21) are satisfied.

Under the assumptions and conditions of solvability, from [7,13,21], it is easily known that  $\Phi^+ \in L^2(\mathbb{R})$ .

Suppose that  $s = \infty$  is a special node, then  $\lambda_\infty = 0$ . It follows from (4.4) that  $\gamma_\infty \neq 0$ , thus we can transform it to the case  $\lambda_\infty < \frac{1}{2}$ . Further discussions are the same as above.

In conclusion, we state the following result.

**Theorem 4.1.** Under the case of non-normal type, (3.36) is a necessary condition of solvability for Eq. (4.3).

(1) If  $\lambda_\infty > \frac{1}{2}$ , in (4.13), we write  $P_{\kappa-M}(z)$  in place of  $P_{\kappa-M-1}(z)$ . When  $\kappa > M$ , (4.1) has  $\kappa - M$  linearly independent solutions. When  $\kappa = M$ , (4.3) has the unique solution. When  $\kappa < M$ , (4.19) and (4.25) hold, in this case, (4.3) has the general solution. When  $\kappa > 0$ , (4.18) holds; when  $\kappa = 0$ , (4.20) holds; when  $\kappa < 0$ , (3.22) and (4.21) hold.

(2) If  $\lambda_\infty \leq \frac{1}{2}$ , when  $\kappa \geq M$ , (4.18) holds. When  $\kappa < M$ , (4.19) holds. When  $\kappa \geq 0$ , (4.25) holds, and when  $\kappa < 0$ , (4.19) and (4.21) hold.

Under the conditions of solvability, (4.1) has the general solutions

$$\psi^+(\tau) = \mathcal{W}^{-1}[\Phi^+(s)], \tag{4.26}$$

where  $\Phi^+(s)$  is given by (4.14).

## 5 Dual SIEs with convolution kernels

Now we deal with dual SIEs with convolution kernels and variable integral limits

$$\begin{cases} A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_0^\tau \psi(t)h_1(\tau-t)dt + \frac{C}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau), & \tau \in \mathbb{R}^+; \\ A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_\tau^0 \psi(t)h_2(\tau-t)dt + \frac{C}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau), & \tau \in \mathbb{R}^-, \end{cases} \tag{5.1}$$

where  $A, B, C$  are constants, and  $C \neq 0$ . The functions  $h_1, h_2, n \in H_1$ , and the unknown function  $\psi \in H_1$ . Combining the two equations in (5.1) to the following (5.2):

$$\begin{aligned} & A\psi(\tau) + \frac{B}{\sqrt{2\pi}} \int_{\mathbb{R}} \psi(t)h(\tau-t)dt + \frac{B}{\sqrt{2\pi}} \int_{\mathbb{R}} \psi(t)k(\tau-t)\text{sgn}(t)dt \\ & + \frac{C}{\pi i} \int_{\mathbb{R}} \frac{\psi(t)}{t-\tau} dt = n(\tau), \quad \tau \in \mathbb{R}, \end{aligned} \tag{5.2}$$

where

$$h(\tau) = \frac{1}{2}[h_1^+(\tau) - h_2^-(\tau)], \quad k(\tau) = \frac{1}{2}[h_1^+(\tau) + h_2^-(\tau)],$$

then  $h, k \in H_1$ . Obviously,

$$h(\tau) = \begin{cases} \frac{1}{2}h_1(\tau), & \tau \geq 0; \\ \frac{1}{2}h_2(\tau), & \tau < 0, \end{cases} \quad k(\tau) = \begin{cases} \frac{1}{2}h_1(\tau), & \tau \geq 0; \\ -\frac{1}{2}h_2(\tau), & \tau < 0. \end{cases}$$

We apply the operator  $\mathcal{W}$  to (5.2) and obtain

$$A\Phi(s) + BH(s)\Phi(s) + \frac{BK(s)}{\pi i} \int_{\mathbb{R}} \frac{\Phi(t)}{t-s} dt - C\Phi(s)\text{sgn}(s) = N(s), \quad s \in \mathbb{R}, \quad (5.3)$$

where

$$\mathcal{W}[h(\tau)] = H(s), \quad \mathcal{W}[k(\tau)] = K(s), \quad \mathcal{W}[n(\tau)] = N(s), \quad \mathcal{W}[\psi(\tau)] = \Phi(s),$$

and  $H, K, N, \Phi \in H_1^*$ . We shall solve the following (5.4) instead of (5.3)

$$\Phi^+(s) = J(s)\Phi^-(s) + N_0(s), \quad (5.4)$$

where

$$J(s) = \frac{A + B[H(s) - K(s)] - C\text{sgn}(s)}{A + B[H(s) + K(s)] - C\text{sgn}(s)}, \quad N_0(s) = \frac{N(s)}{A + B[H(s) + K(s)] - C\text{sgn}(s)}.$$

From the analysis above, it is easy to prove that (5.4) has a unique node  $s = 0$ . Further process is similar to section 3 and will be omitted.

## 6 Conclusions

In this paper, we mainly focus on three classes SIEs with variable integral limits in the case of non-normal type. By means of the theory of complex variable functions and classical boundary value problems, Eqs. (1.1)-(1.3) are transformed into the linear Riemann problems with discontinuous property in class  $H_1^*$ , further, we obtain the general solution given by integral-form. Moreover, the novel method in this paper can effectively solve such equations mentioned in [28-33], and we may also prove the existence and stability of solutions for Eqs. (1.1)-(1.3) in Clifford analysis (see [34,35]).

Recently, many scholars have studied SIEs with convolution kernels and non-linear SIEs in multidimensional and hyper-singular fields, and gained a lot of excellent results. Based on our results in this paper, we may consider the solvability theory of Eqs. (1.1)-(1.3) in these areas. More precise details will be omitted now.

## Acknowledgment

This work is supported financially by the National Natural Science Foundation of China (11971015).

## Availability of data and materials

Our manuscript has no associated data.

## Competing interests

The authors declare that they have no competing interests.

## References

- [1]F. D. Gakhov, The boundary value problem of analysis function[M]. Moscow: Nauka Press, 1977.
- [2]F. D. Gakhov, Boundary Value Problems, New York, 1990.
- [3]F. D. Gakhov, U. I. Chersky, Integral Equations of Convolution type, Nauka, Moscow, 1987.
- [4]G. S. Litvinchuk, Solvability Theory of Boundary Value Problems and Singular Integral Equations with Shift, Kluwer Academic Publisers, London, 2004.
- [5]N. I. Muskhelishvilli, Singular Integral Equations, NauKa, Moscow, 2002.
- [6]J. K. Lu, Boundary Value Problems for Analytic Functions, Singapore, World Sci., 2004.
- [7]J. K. Lu, On method of solution for some classes of singular integral equations with convolution. Chin. Ann. of Math., 8B(1)(1987), 97-108.
- [8]H. Du, J. H. Shen, Reproducing kernel method of solving singular integral equation with cosecant kernel, J. Math. Comput. 384(1)(2008), 308-314.
- [9]P. R. Li, Singular integral equations of convolution type with reflection and translation shifts, Numer. Func. Anal. Opt., 40 (9)(2019), 1023-1038.
- [10]P. R. Li, G. B. Ren, Some classes of equations of discrete type with harmonic singular operator and convolution, Appl. Math. Comput., 284(2016), 185-194.
- [11]P. R. Li, G. B. Ren, Solvability of singular integro-differential equations via Riemann-Hilbert problem, J. Differ. Equ., 265(2018), 5455-5471.
- [12]P. R. Li, Generalized boundary value problems for analytic functions with convolutions and its applications, Math. Meth. Appl. Sci., 42(2019), 2631-2643.
- [13]R. V. Duduchava, Integral equations of convolution type with discontinuous coefficients, Math. Nachr., 79(1977), 75-78.
- [14]C. X. Miao, J. Y. Zhang, J. Q. Zheng, Scattering theory for the radial  $\dot{H}^{\frac{1}{2}}$ -critical wave equation with a cubic convolution, J. Differ. Equ., 259(2015), 7199-7237.
- [15]J. K. Lu, Some classes boundary value problems and singular integral equations with a transformation. Advances in Mathematics, 23(5)(1994), 424-431.
- [16]R. V. Duduchava, General singular integral equations and fundamental problems of the plane theory of elasticity, Trudy Tbiliss. Math. Inst. Razmadze, 82(1986), 45-89.

- [17]J. K. Lu, On solution of a kind of Riemann boundary value problem with square roots, *Acta Math. Sci.*, 22B(2)(2002), 145-512.
- [18]S. Ma, Traveling waves for non-local delayed diffusion equations via auxiliary equations, *J. Differ. Equ.*, 237(2007), 259-277.
- [19]L. Hörmander, *The Analysis of Linear Partial Differential Operators. I*, Reprint of the second (1990) edition, Springer-Verlag, Berlin, 2003.
- [20]M. C. De-Bonis, C. Laurita, Numerical solution of systems of Cauchy singular integral equations with constant coefficients, *Appl. Math. Comput.*, 219(2012), 1391-1410.
- [21]P. R. Li, Solvability theory of convolution singular integral equations via Riemann-Hilbert approach, *J. Comput. Appl. Math.*, 370(2)(2020), 112601.
- [22]P. R. Li, Existence of analytic solutions for some classes of singular integral equations of non-normal type with convolution kernel, *Acta Applicandae Mathematicae*, 181(1)(2022).
- [23]S. G. Zhong, The growth in time of higher Sobolev norms of solutions to Schrödinger equations on compact Riemannian manifolds, *J. Differ. Equ.*, 245(2)(2008), 359-376.
- [24]P. R. Li, Existence of solutions for dual singular integral equations with convolution kernels in case of non-normal type. *J. Appl. Anal. Comput.*, 10(6)(2020), 2756-2766.
- [25]P. Wójcik, M. A. Sheshko, S. M. Sheshko, Application of Faber polynomials to the approximate solution of singular integral equations with the Cauchy kernel, *Differential Equations*, 49(2)(2013), 198-209.
- [26]Z. H. Du, J. Y. Du, On direct method of solution for a class of singular integral equations, *Acta Math. Sci.*, 27(2007), 301-307.
- [27]P. R. Li, Non-normal type singular integral-differential equations by Riemann-Hilbert approach, *J. Math. Anal. Appl.*, 483(2)(2020), 123643.
- [28]B. Guo, N. Liu, Y. Wang, Long-time asymptotics for the Hirota equation on the half-line, *Nonlinear Anal.*, 174(2018), 118-140.
- [29]J. Colliander, M. Keel, G. Staffilani, Transfer of energy to high frequencies in the cubic defocusing nonlinear Schrödinger equation, *Invent. Math.*, 181(1)(2010), 39-113.
- [30]N. M. Tuan, N. T. Thu-Huyen, The solvability and explicit solutions of two integral equations via generalized convolutions, *J. Math. Anal. Appl.*, 369(2010), 712-718.
- [31]Y. Jiang, Y. Xu, A fast Fourier-Galerkin methods for solving singular boundary integral equations: numerical integration and precondition, *J. Comput. Appl. Math.*, 234(2010), 2792-2807.
- [32]H. Khosravi, R. Allahyari, A. Haghighi, Existence of solutions of functional integral equations of convolution type using a new construction of a measure of noncompactness on  $L^p(\mathbb{R}_+)$ , *Appl. Math. Comput.*, 260(2015), 140-147.
- [33]Y. F. Wang, J. Y. Du, On Riemann boundary value problem for polyanalytic functions on the real axis, *Acta. Math. Sci.*, 24B(4)(2004), 663-671.
- [34]Y. F. Gong, L. T. Leong, T. Qiao, Two integral operators in Clifford analysis, *J. Math. Anal. Appl.*, 354(2009), 435-444.

- [35]Z. Blocki, Suita conjecture and the Ohsawa-Takegoshi extension theorem. *Invent. Math.*, 193(2013), 149-158.
- [36]D. Kinzebulatov, K. R. Madou, Stochastic equations with time-dependent singular drift, *J. Differ. Equ.*, 337(2022), 255-293.
- [37]I. Zamanpour, R. Ezzati, Operational matrix method for solving fractional weakly singular 2D partial Volterra integral equations, *J. Comput. Appl. Math.*, 419(2023), 114704.
- [38]T. K. Wang, S. J. Liu, Z. Y. Zhang, Singular expansions and collocation methods for generalized Abel integral equations, *J. Comput. Appl. Math.*, 429(2023), 115240.
- [39]Y. P. Chen, Z. H. Guo, Generalized singular integral with rough kernel and approximation of surface quasi-geostrophic equation, *J. Differ. Equ.*, 346(2023), 205-228.
- [40]Z. J. Du, J. Li, Geometric singular perturbation analysis to Camassa-Holm Kuramoto-Sivashinsky equation, *J. Differ. Equ.*, 306(2022), 418-438.
- [41]D. S. Kong, S. H. Xiang, H. Y. Wu, An efficient numerical method for Volterra integral equation of the second kind with a weakly singular kernel, *J. Comput. Appl. Math.*, 427(2023), 115101.
- [42]R. Katani, S. McKee, Numerical solution of two-dimensional weakly singular Volterra integral equations with non-smooth solutions, *J. Comput. Appl. Math.*, 402(2022), 113779.