

SOLVABILITY FOR COUPLED IMPULSIVE FRACTIONAL PROBLEMS OF THE KIRCHHOFF TYPE WITH $P&Q$ -LAPLACIAN

Yi Wang¹ and Lixin Tian^{1,2,†}

Abstract In this paper, we investigate the existence and multiplicity of nontrivial solutions for a class of $p&q$ -Laplacian Kirchhoff impulsive fractional differential equations through variational methods. By leveraging the Nehari manifold and fibering maps, we establish the existence of at least one nontrivial solution to such equations for any pair of parameters $(\lambda, \mu) \in \Theta_*$. Furthermore, using the idea of truncation arguments and Krasnoselskii genus theory, we demonstrate the existence of infinitely many nontrivial solutions for the equation when Kirchhoff functions M_1 and M_2 are degenerate considering any pair of parameters $(\lambda, \mu) \in \Theta_{**}$.

Keywords Kirchhoff fractional differential equations, $p&q$ -Laplacian, impulsive problems, variational methods.

MSC(2010) 34A08, 34K45, 35A15, 58E05.

1. Introduction

This paper focuses on investigating the existence and multiplicity of nontrivial solutions to the $p&q$ -Laplacian impulsive fractional differential equations involving Kirchhoff functions:

$$\left\{ \begin{array}{l} [M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_tD_T^\alpha \Phi_p({}_0D_t^\alpha u(t)) + [M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_tD_T^\beta \Phi_q({}_0D_t^\beta u(t)) \\ = F_u(t, u(t), v(t)), \quad t \neq t_j, \quad \text{a.e. } t \in [0, T], \\ [M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_tD_T^\alpha \Phi_p({}_0D_t^\alpha v(t)) + [M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_tD_T^\beta \Phi_q({}_0D_t^\beta v(t)) \\ = F_v(t, u(t), v(t)), \quad t \neq t'_i, \quad \text{a.e. } t \in [0, T], \\ \Delta([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1} \Phi_p({}_0D_t^\alpha u))(t_j) \\ + \Delta([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1} \Phi_q({}_0D_t^\beta u))(t_j) + \lambda \zeta(t_j) I_j(u(t_j)) = 0, \quad j \in \Delta_1, \\ \Delta([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1} \Phi_p({}_0D_t^\alpha v))(t'_i) \\ + \Delta([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1} \Phi_q({}_0D_t^\beta v))(t'_i) + \mu \varrho(t'_i) S_i(v(t'_i)) = 0, \quad i \in \Delta_2, \\ u(0) = u(T) = 0, \quad v(0) = v(T) = 0, \end{array} \right. \quad (1.1)$$

[†]the corresponding author.

Email address: wangyistum@163.com (Y. Wang), tianlx@ujs.edu.cn (L. Tian)

¹School of Mathematical Sciences, Nanjing Normal University, Nanjing 210023, China

²School of Mathematical Sciences, Jiangsu University, Zhenjiang 212013, China

where $M_1, M_2 \in C^1(R_0^+, R_0^+)$, $1 < p, q < \infty$, $\alpha \in (\frac{1}{p}, 1]$, $\beta \in (\frac{1}{q}, 1]$, ${}_0D_t^\gamma$ and ${}_tD_T^\gamma$ denote left and right standard Riemann-Liouville fractional derivatives, respectively, $\gamma \in \{\alpha, \beta\}$, $\Phi_v(z) = |z|^{v-2}z$, $v > 1$, $v \in \{p, q\}$, λ, μ are real parameters, $\zeta(t), \varrho(t) \in C([0, T], R)$ with $\zeta_\pm(t) = \pm \max\{\pm\zeta(t), 0\} \neq 0$, $\varrho_\pm(t) = \pm \max\{\pm\varrho(t), 0\} \neq 0$, $F : [0, T] \times R^2 \rightarrow R$ is continuous with respect to t , for all $(u, v) \in R^2$, continuously differentiable with respect to u and v for almost every $t \in [0, T]$, F_u and F_v denote the partial derivatives of F with respect to u and v , norms $\|\cdot\|_\alpha, \|\cdot\|_\beta$ are specified later, $0 = t_0 < t_1 < \dots < t_m < t_{m+1} = T$, $0 = t'_0 < t'_1 < \dots < t'_n < t'_{n+1} = T$, $\Delta_1 = \{1, 2, \dots, m\}$, $\Delta_2 = \{1, 2, \dots, n\}$, $I_j, S_i \in C^1(R, R)$ for all $j \in \Delta_1, i \in \Delta_2$ and

$$\begin{aligned} & \Delta([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha u))(t_j) \\ & + \Delta([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta u))(t_j) \\ = & [M_1(\|(u(t_j^+), v(t_j^+))\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha u(t_j^+)) \\ & + [M_2(\|(u(t_j^+), v(t_j^+))\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta u(t_j^+)) \\ & - [M_1(\|(u(t_j^-), v(t_j^-))\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha u(t_j^-)) \\ & - [M_2(\|(u(t_j^-), v(t_j^-))\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta u(t_j^-)), \end{aligned}$$

and

$$\begin{aligned} & \Delta([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha v))(t'_i) \\ & + \Delta([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta v))(t'_i) \\ = & [M_1(\|(u(t_j^+), v(t_j^+))\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha v(t_i^{++})) \\ & + [M_2(\|(u(t_j^+), v(t_j^+))\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta v(t_i^{++})) \\ & - [M_1(\|(u(t_j^-), v(t_j^-))\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha v(t_i^{--})) \\ & - [M_2(\|(u(t_j^-), v(t_j^-))\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta v(t_i^{--})), \end{aligned}$$

where

$$\begin{aligned} & [M_1(\|(u(t_j^\pm), v(t_j^\pm))\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha u(t_j^\pm)) \\ & + [M_2(\|(u(t_j^\pm), v(t_j^\pm))\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta u(t_j^\pm)) \\ = & \lim_{t \rightarrow t_j^\pm} [M_1(\|(u(t), v(t))\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha u(t)) \\ & + [M_2(\|(u(t), v(t))\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta u(t)), \end{aligned}$$

and

$$\begin{aligned} & [M_1(\|(u(t_i'^\pm), v(t_i'^\pm))\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha v(t_i'^\pm)) \\ & + [M_2(\|(u(t_i'^\pm), v(t_i'^\pm))\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta v(t_i'^\pm)) \\ = & \lim_{t \rightarrow t_i'^\pm} [M_1(\|(u(t), v(t))\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1}\Phi_p({}_0D_t^\alpha v(t)) \\ & + [M_2(\|(u(t), v(t))\|_\beta^q)]^{q-1} {}_tD_T^{\beta-1}\Phi_q({}_0D_t^\beta v(t)). \end{aligned}$$

Fractional differential equations (FDEs) are an extension of ordinary differential equations and integration to arbitrarily noninteger orders that have risen in prominence as a result of their extensive applications in many disciplines of research and

engineering. In actuality, several equations incorporating fractional order derivatives are used to simulate a wide range of phenomena, including fluid flow, diffusive transport comparable to diffusion, rheology, probability, and electrical networks; see previous studies (see [2, 6, 12]). It should be emphasized that the existence and multiplicity of solutions to nonlinear boundary value problems of FDEs have been extensively investigated using critical point theory and variational methods (see [10, 11, 15, 18, 21, 30]). In recent years, many scholars have devoted themselves to the study of classical p & q -Laplacian elliptic type equations over bounded and unbounded domains using variational methods and critical point theory (see [7, 20, 24]). At the same time, we are aware of very few contributions concerning FDEs with p & q -Laplacian and impulsive terms (see [16, 17, 31]). For example, Li et al. [17] were the first to study the following impulsive fractional boundary value problem with p & q -Laplacian operators:

$$\begin{cases} {}_t D_T^\alpha \Phi_p({}_0 D_t^\alpha u(t)) + |u(t)|^{p-2} u(t) + {}_t D_T^\beta \Phi_q({}_0 D_t^\beta u(t)) + |u(t)|^{q-2} u(t) \\ = f(t, u(t), {}_0 D_t^\alpha u(t), {}_0 D_t^\beta u(t)), \quad t \neq t_j, \quad \text{a.e. } t \in [0, T], \\ \Delta({}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha u) + {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta u))(t_j) = I_j(u(t_j)), \quad j = 1, 2, \dots, m, \\ u(0) = u(T) = 0, \quad \text{a.e. } t \in [0, T], \end{cases} \quad (1.2)$$

where ${}_0 D_t^\alpha$, ${}_0 D_t^\beta$ and ${}_t D_T^\alpha$, ${}_t D_T^\beta$ denote the left and right Riemann-Liouville fractional derivatives, respectively. Based on the Mountain pass theorem and the iterative technique, Li et al. obtained the existence of at least one nontrivial solution to problem (1.2).

On the other hand, great attention has recently been focused on the study of Kirchhoff-type differential equations which stand out due to the presence of a Kirchhoff function $M \in C(R_0^+, R_0^+)$. In recent years, there have been fruitful achievements in using variational methods to study the existence and multiplicity of solutions for Kirchhoff-type differential equations (see [1, 26, 27, 29]). In addition, some new results have been obtained in the study of Kirchhoff-type FDEs by variational methods and critical point theory when the Kirchhoff function M is given by $M(s) = a + bs$ for all $s \in R_0^+$, where $a, b > 0$. For example, Kratou [14] studied the existence and multiplicity of solutions to the Kirchhoff fractional equation with singular nonlinearity:

$$\begin{cases} \left(a + b \int_0^T |{}_0 D_t^\alpha u(t)|^p dt \right)^{p-1} {}_t D_T^\alpha \Phi_p({}_0 D_t^\alpha u(t)) = \frac{\lambda g(t)}{u^\gamma(t)} + f(t, u(t)), \quad t \in (0, T), \\ u(0) = u(T) = 0, \end{cases} \quad (1.3)$$

where $a \geq 1$, $b, \lambda > 0$, $p > 1$, $\alpha \in (\frac{1}{p}, 1]$, $0 < \gamma < 1 < p < r$, $g \in C([0, T])$. Using the idea of the Nehari manifold technique, Kratou proved the existence of at least two positive solutions to the problem (1.3). When $\frac{\lambda g(t)}{u^\gamma(t)} = 0$, Chen and Liu [4] used the Nehari manifold method to obtain that there exists at least one nontrivial ground state solution to problem (1.3); Chen et al. [5] obtained that problem (1.3) has at least one solution and infinitely many nontrivial weak solutions by using the mountain pass theorem and the genus properties. To the best of my knowledge, there are few related results on the existence and multiplicity of nontrivial solutions for a class of coupled p & q -Laplacian Kirchhoff-type impulsive fractional problems.

Mainly inspired by the above works, we investigate the existence and multiplicity

of solutions for problem (1.1) by using variational methods and critical point theory. Problem (1.1) is more general in the form of the equation. For examples, when $M_1 = M_2 = 1$, problem (1.1) can be reduced to the $p&q$ -Laplacian impulsive fractional differential equations; when $M_1 = M_2 = 1$, $p = q = 2$, and $\alpha = \beta = 1$, problem (1.1) is transformed into the following second-order impulsive differential equations

$$\begin{cases} -2u''(t) = F_u(t, u(t), v(t)), & t \neq t_j, \text{ a.e. } t \in [0, T], \\ -2v''(t) = F_v(t, u(t), v(t)), & t \neq t'_i, \text{ a.e. } t \in [0, T], \\ 2\Delta(u'(t_j)) + \lambda\zeta(t_j)I_j(u(t_j)) = 0, & j \in \Delta_1, \\ 2\Delta(v'(t'_i)) + \mu\rho(t'_i)S_i(v(t'_i)) = 0, & i \in \Delta_2, \\ u(0) = u(T) = 0, & v(0) = v(T) = 0, \end{cases}$$

For that reason, problem (1.1) is an extension of the integer-order impulsive differential equation.

Another novel aspect of this paper is that we provide an existence result utilising the technique of Nehari manifold and fibering maps and a multiplicity result utilising the Krasnoselskii genus for problem (1.1). The main challenge in employing Nehari manifold analysis is that Kirchhoff functions M_1 and M_2 are not specified as $M_1(s) = M_2(s) = a + bs^{\vartheta-1}$ for all $s \in R_0^+$, where $a, b > 0$, $\vartheta \geq 1$, leading to an analysis completely different from Xie and Chen [29], Kratou [14], and Fiscella and Mishra [8]. For the purpose to obtain the critical point of the functional $J_{\lambda, \mu}$ associated with problem (1.1), we are devoted to analyze the behavior of functional $J_{\lambda, \mu}$ over its Nehari manifold $\mathcal{N}_{\lambda, \mu}$. Then, we introduce the decomposition $\mathcal{N}_{\lambda, \mu} = \mathcal{N}_{\lambda, \mu}^+ \cup \mathcal{N}_{\lambda, \mu}^0 \cup \mathcal{N}_{\lambda, \mu}^-$. It is promptly seen that critical points of $J_{\lambda, \mu}|_{\mathcal{N}_{\lambda, \mu}^+ \cup \mathcal{N}_{\lambda, \mu}^-}$ are critical points of $J_{\lambda, \mu}$. Accordingly, it is crucial to make sure that $\mathcal{N}_{\lambda, \mu}^0 = \emptyset$. Considering the generality of the Kirchhoff functions M_1 and M_2 , which lead to $\mathcal{N}_{\lambda, \mu}^0 \neq \emptyset$, condition (M1) effectively avoids this. Moreover, condition (F1) has vital significance in that the homogeneity and Euler identity of F enable us to successfully establish the Nehari manifold $\mathcal{N}_{\lambda, \mu}$. Condition (F2) is also indispensable to ensure the compactness of functional $J_{\lambda, \mu}$. Motivated by Xiang et al. [28], we provide an multiplicity result for problem (1.1) by the Krasnoselskii genus. Xiang et al. [28] investigated the Kirchhoff function $M(s) = a + b\theta s^{\theta-1}$ for all $s \geq 0$, where $a \geq 0$, $b \geq 0$, $a + b > 0$ and $\theta > 1$. Nevertheless, M_1 and M_2 are more general, making the analysis more challenging. Moreover, taking into account the nonlocal nature of Kirchhoff functions M_1 and M_2 and the impact of sign-changing weight functions ζ and ρ , we establish some new estimates under the perturbing influence of the pair of parameters (λ, μ) to gain truncated functional $\bar{J}_{\lambda, \mu}$. Finally, it is evident from conditions (M1') and (M2') that Kirchhoff functions M_1 and M_2 are degenerate, which forces us to discuss the (PS)-sequence compactness of truncated functional $\bar{J}_{\lambda, \mu}$ in three cases, as displayed in (ii) of Lemma 4.2 below.

In order to precisely establish the existence result for problem (1.1) in Section 3, we introduce the following assumptions conditions:

- (M1) There exist constants $1 < p\vartheta_1, q\vartheta_2 < p, q \leq \kappa \leq p^2, q^2$, $\eta_1, \eta_2 > 1$, $0 < \xi_1 < 1 < \delta_1$, $(\sigma - p)\xi_1\eta_1 > (p^2 - p)\delta_1$, $0 < \xi_2 < 1 < \delta_2$, $(\sigma - q)\xi_2\eta_2 > (q^2 - q)\delta_2$, $\xi_1\sigma > \vartheta_1\delta_1p$, $(2p + p^2)\delta_1\theta < \xi_1$, $\xi_2\sigma > \vartheta_2\delta_2q$ and $(2q + q^2)\delta_2\theta < \xi_2$ such that

$$\begin{aligned} \xi_1 x^{\frac{\kappa}{p}} &< \vartheta_1 \bar{M}_1(x) \leq [M_1(x)]^{p-1} x \leq \eta_1 [M_1(x)]^{p-2} M_1'(x) x^2 < \delta_1 x^{\frac{\kappa}{p}}, \\ \xi_2 y^{\frac{\kappa}{q}} &< \vartheta_2 \bar{M}_2(y) \leq [M_2(y)]^{q-1} y \leq \eta_2 [M_2(y)]^{q-2} M_2'(y) y^2 < \delta_2 y^{\frac{\kappa}{q}}, \end{aligned}$$

for all $x, y > 0$, where $\overline{M}_1(x) = \int_0^x [M_1(s)]^{p-1} ds$, $\overline{M}_2(y) = \int_0^y [M_2(s)]^{q-1} ds$;

(H1) There exist positive constants a_j, b_j with $\bar{a} = \max\{a_j\}$, $\bar{b} = \max\{b_i\}$ for all $j \in \Delta_1, i \in \Delta_2, 0 < \theta < 1 < p, q$ such that

$$|I_j(z)z| \leq a_j|z|^\theta, \text{ and } |S_i(z)z| \leq b_i|z|^\theta, \forall z \in R;$$

(H2) For all $j \in \Delta_1, i \in \Delta_2, I_j, K_i : R \rightarrow R$ such that

$$I_j(su(t_j)) = s^{\theta-1}I_j(u(t_j)), \text{ and } S_i(sv(t'_i)) = s^{\theta-1}S_i(v(t'_i)),$$

for all $s > 0$ and $(u, v) \in R^2$;

(F1) $F(t, u, v)$ is σ -homogeneous and satisfies Euler identity with $\sigma\theta > 1, \sigma > p^2, q^2$, that is

$$F(t, su, sv) = s^\sigma F(t, u, v), \text{ and } uF_u(t, u, v) + vF_v(t, u, v) = \sigma F(t, u, v),$$

for all $(u, v) \in R^2$;

(F2) There exist constants $C_* > 0$ such that

$$|F(t, u, v)| \leq C_*(u, v)^\sigma, \text{ for all } (u, v) \in R^2,$$

where $|(u, v)| = (u^2 + v^2)^{\frac{1}{2}}$.

In order to obtain the multiplicity result by the Krasnoselskii genus for problem (1.1) in Section 4, we give the following assumptions conditions:

(M1') There exists $\bar{\vartheta}_1, \bar{\vartheta}_2 \geq 1$ such that

$$[M_1(x)]^{p-1}x \leq \bar{\vartheta}_1\overline{M}_1(x), \text{ and } [M_2(y)]^{q-1}y \leq \bar{\vartheta}_2\overline{M}_2(y), \text{ for all } x, y \in R_0^+;$$

(M2') $M_1(0) = M_2(0) = 0$, and for any $\tau, \bar{\tau} > 0$, there exist $m_1 = m_1(\tau) > 0$ and $m_2 = m_2(\bar{\tau}) > 0$ such that

$$M_1(x) \geq m_1, \text{ for all } x \geq \tau, \text{ and } M_2(y) \geq m_2, \text{ for all } y \geq \bar{\tau};$$

(F1') There exist $\bar{\eta}$ with $\bar{\vartheta}_* = \max\{p\bar{\vartheta}_1, q\bar{\vartheta}_2\} < \bar{\eta}$ such that for any $\varepsilon > 0$, there exists $C_\varepsilon > 0$ for which

$$|\nabla F(t, u, v)| \leq \bar{\vartheta}_*\varepsilon|(u, v)|^{\bar{\vartheta}_*-1} + \bar{\eta}C_\varepsilon|(u, v)|^{\bar{\eta}-1}, \text{ for all } (u, v) \in R^2,$$

holds, where $|(u, v)| = (u^2 + v^2)^{\frac{1}{2}}$ and $\nabla F = (F_u, F_v)$;

(F2') There exists $0 < \bar{\zeta} < \theta < 1$ such that

$$0 \leq \bar{\zeta}F(t, u, v) \leq \nabla F(t, u, v) \cdot (u, v), \text{ for all } (u, v) \in R^2;$$

(F3') $F(t, -u, -v) = F(t, u, v)$, for all $(t, u, v) \in [0, T] \times R^2$, $I_j(-z) = -I_j(z)$ and $S_i(-z) = -S_i(z)$ for all $z \in R, j \in \Delta_1, i \in \Delta_2$.

Remark 1.1. Kirchhoff problems are said to be non-degenerate if $\inf_{s \in R_0^+} M(s) > 0$ and degenerate if $M(0) = 0$ and $\inf_{s \in R^+} M(s) > 0$. some recent results concerning the degenerate case on Kirchhoff-type problems are referred to in [9, 26, 28].

Our main results are as follows:

Theorem 1.1. *Assume that (M1), (H1), (H2) and (F1), (F2) hold. Then, for any $(\lambda, \mu) \in \Theta_*$, problem (1.1) admits at least a nontrivial solution.*

Theorem 1.2. *Assume that (H1), (M1'), (M2') and (F1')-(F3') hold. Then, for any $(\lambda, \mu) \in \Theta_{**}$, problem (1.1) admits infinitely many nontrivial solutions.*

The paper is structured as follows: In Section 2, we recall some definitions of fractional calculus and discuss the variational formulation of problem (1.1). In Section 3, we introduce the Nehari manifold structure and fibering maps analysis related to problem (1.1) to obtain the existence of the solution. In Section 4, we apply Krasnoselskii genus theory to investigate the infinitely many solutions for problem (1.1).

2. Preliminaries

In this section, we present some preliminary findings that will be utilized in the subsequent sections.

For $0 < \gamma \leq 1$ and $1 < v < \infty$, the functional space that incorporates this boundary condition will be denoted by $E_{\gamma,v}$ as the closure of $C_0^\infty([0, T], R)$, where

$$E_{\gamma,v} = \left\{ u \in L^v([0, T], R) \mid {}_0D_t^\gamma u(t) \in L^v([0, T], R), u(0) = u(T) = 0 \right\},$$

endowed with the norm

$$\|z\|_{\gamma,v} = \left(\int_0^T |{}_0D_t^\gamma z(t)|^v dt \right)^{\frac{1}{v}}, \quad \forall z \in E_{\gamma,v}.$$

Lemma 2.1 ([13]). *Let $0 < \gamma \leq 1$ and $1 < v < \infty$. For any $z \in E_{\gamma,v}$, we have*

$$\|z\|_{L^v} \leq \frac{T^\gamma}{\Gamma(\gamma+1)} \|{}_0D_t^\gamma z\|_{L^v} = \mathcal{A}_{\gamma,v} \|z\|_{\gamma,v}, \quad (2.1)$$

furthermore, when $\gamma > \frac{1}{v}$, $\frac{1}{v} + \frac{1}{v'} = 1$, we have

$$\|z\|_\infty \leq \frac{T^{\gamma-\frac{1}{v}}}{\Gamma(\gamma)((\gamma-1)v'+1)^{\frac{1}{v'}}} \|{}_0D_t^\gamma z\|_{L^v} = \bar{\mathcal{A}}_{\gamma,v} \|z\|_{\gamma,v}, \quad (2.2)$$

where $\mathcal{A}_{\gamma,v} = \frac{T^\gamma}{\Gamma(\gamma+1)}$, $\bar{\mathcal{A}}_{\gamma,v} = \frac{T^{\gamma-\frac{1}{v}}}{\Gamma(\gamma)((\gamma-1)v'+1)^{\frac{1}{v'}}$.

Let $E_\alpha = E_{\alpha,p} \times E_{\alpha,p}$ and $E_\beta = E_{\beta,q} \times E_{\beta,q}$, which are reflexive Banach spaces endowed with the norms

$$\|(u, v)\|_\alpha = (\|u\|_{\alpha,p}^p + \|v\|_{\alpha,p}^p)^{\frac{1}{p}}, \quad \text{and} \quad \|(u, v)\|_\beta = (\|u\|_{\beta,q}^q + \|v\|_{\beta,q}^q)^{\frac{1}{q}}.$$

Then, set $E_{\alpha,\beta} = E_\alpha \cap E_\beta$ endowed with the norm

$$\|(u, v)\|_{\alpha,\beta} = \|(u, v)\|_\alpha + \|(u, v)\|_\beta, \quad \forall (u, v) \in E_{\alpha,\beta}. \quad (2.3)$$

For all $\bar{v} \geq p, q$ and $(u, v) \in E_{\alpha,\beta}$, by (2.2), we have

$$\|(u, v)\|_{L^{\bar{v}}}^{\bar{v}} \leq 2^{\bar{v}-1} (\|u\|_{L^{\bar{v}}([0,T])}^{\bar{v}} + \|v\|_{L^{\bar{v}}([0,T])}^{\bar{v}})$$

$$\begin{aligned}
&\leq 2^{\bar{v}-1} T \bar{\mathcal{A}}_{\alpha,p}^{\bar{v}} (\|u\|_{\alpha,p}^{\bar{v}} + \|v\|_{\alpha,p}^{\bar{v}}) \\
&\leq \bar{\mathcal{M}}(\bar{v}) \|(u, v)\|_{\alpha,\beta}^{\bar{v}},
\end{aligned} \tag{2.4}$$

where $\bar{\mathcal{M}}(\bar{v}) = 2^{\bar{v}-1} T \bar{\mathcal{A}}_{\alpha,p}^{\bar{v}}$, the inequality $2^{1-\bar{i}}(a+b)^{\bar{i}} \leq a^{\bar{i}} + b^{\bar{i}} \leq (a+b)^{\bar{i}}$ for all $\bar{i} \geq 1$ and $a, b \geq 0$ is applied.

Lemma 2.2 ([13]). *Let $\frac{1}{p} < \alpha \leq 1$, $\frac{1}{q} < \beta \leq 1$, $1 < p, q < \infty$. The fractional derivative spaces E_α and E_β are reflexive and separable Banach spaces.*

Lemma 2.3 ([13]). *Let $\frac{1}{p} < \alpha \leq 1$, $\frac{1}{q} < \beta \leq 1$, $1 < p, q < \infty$. Assume that the sequence $\{z_n\}$ converges weakly to z in E_α or E_β , i.e., $z_n \rightharpoonup z$. Then $\{z_n\}$ converges strongly to z in $(C([0, T], R))^2$, i.e., $\|z_n - z\|_\infty \rightarrow 0$ as $n \rightarrow \infty$.*

Lemma 2.4. *Let $\{(u_n, v_n)\}_n$ and (u, v) be in $E_{\alpha,\beta}$ such that $(u_n, v_n) \rightharpoonup (u, v)$ weakly in $E_{\alpha,\beta}$ and $(u_n, v_n) \rightarrow (u, v)$ a.e. in R . Then*

$$\begin{aligned}
&\|(u_n, v_n)\|_\alpha^p - \|(u_n - u, v_n - v)\|_\alpha^p = \|(u, v)\|_\alpha^p + o_n(1), \text{ as } n \rightarrow \infty, \\
&\|(u_n, v_n)\|_\beta^q - \|(u_n - u, v_n - v)\|_\beta^q = \|(u, v)\|_\beta^q + o_n(1), \text{ as } n \rightarrow \infty.
\end{aligned}$$

Proof. The basis of our argument is derived from Lemma 3.2 in [19]. However, we have chosen to omit the proof for brevity. \square

Definition 2.1. We say that the couple $(u, v) \in E_{\alpha,\beta}$ is a weak solution to problem (1.1) if for all $(x, y) \in E_{\alpha,\beta}$, it satisfies

$$\begin{aligned}
&[M_1(\|(u, v)\|_\alpha^p)]^{p-1} \int_0^T \Phi_p({}_0D_t^\alpha u(t)) {}_0D_t^\alpha x(t) + \Phi_p({}_0D_t^\alpha v(t)) {}_0D_t^\alpha y(t) dt \\
&+ [M_2(\|(u, v)\|_\beta^q)]^{q-1} \int_0^T \Phi_q({}_0D_t^\beta u(t)) {}_0D_t^\beta x(t) + \Phi_q({}_0D_t^\beta v(t)) {}_0D_t^\beta y(t) dt \\
&- \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) x(t_j) - \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) y(t'_i) \\
&= \int_0^T F_u(t, u(t), v(t)) x(t) + F_v(t, u(t), v(t)) y(t) dt.
\end{aligned} \tag{2.5}$$

Problem (1.1) possesses a variational structure, and its solutions can be regarded as critical points of the functional $J_{\lambda,\mu}$, where

$$\begin{aligned}
J_{\lambda,\mu}(u, v) &= \frac{1}{p} \bar{M}_1(\|(u, v)\|_\alpha^p) + \frac{1}{q} \bar{M}_2(\|(u, v)\|_\beta^q) - \sum_{j=1}^m \int_0^{u(t_j)} \lambda \zeta(t_j) I_j(z) dz \\
&\quad - \sum_{i=1}^n \int_0^{v(t'_i)} \mu \varrho(t'_i) S_i(s) ds - \int_0^T F(t, u(t), v(t)) dt,
\end{aligned} \tag{2.6}$$

for any $(u, v) \in E_{\alpha,\beta}$. Then, $J_{\lambda,\mu} : E_{\alpha,\beta} \rightarrow R$ is of class $C^1(E_{\alpha,\beta}, R)$ and

$$\begin{aligned}
&\langle J'_{\lambda,\mu}(u, v), (x, y) \rangle \\
&= [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \int_0^T \Phi_p({}_0D_t^\alpha u(t)) {}_0D_t^\alpha x(t) + \Phi_p({}_0D_t^\alpha v(t)) {}_0D_t^\alpha y(t) dt \\
&\quad + [M_2(\|(u, v)\|_\beta^q)]^{q-1} \int_0^T \Phi_q({}_0D_t^\beta u(t)) {}_0D_t^\beta x(t) + \Phi_q({}_0D_t^\beta v(t)) {}_0D_t^\beta y(t) dt
\end{aligned}$$

$$\begin{aligned}
& - \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) x(t_j) - \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) y(t'_i) \\
& - \int_0^T F_u(t, u(t), v(t)) x(t) + F_v(t, u(t), v(t)) y(t) dt, \tag{2.7}
\end{aligned}$$

for any $(u, v), (x, y) \in E_{\alpha, \beta}$.

Lemma 2.5. *If the function $(u, v) \in E_{\alpha, \beta}$ is a weak solution of problem (1.1), then (u, v) is a classical solution of problem (1.1).*

Proof. If $(u, v) \in E_{\alpha, \beta}$ is a weak solution of problem (1.1), then (2.5) holds for any $(x, y) \in E_{\alpha, \beta}$. Choose a function $(x, y) \in C_0^\infty(t_j, t_{j+1}) \times C_0^\infty(t'_i, t'_{i+1})$ for any $j \in \Delta_1$ and $i \in \Delta_2$. Then, (2.5) becomes

$$\begin{aligned}
& [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \left(\int_{t_j}^{t_{j+1}} \Phi_p({}_0D_t^\alpha u)_0 D_t^\alpha x dt + \int_{t'_i}^{t'_{i+1}} \Phi_p({}_0D_t^\alpha v)_0 D_t^\alpha y dt \right) \\
& + [M_2(\|(u, v)\|_\beta^q)]^{q-1} \left(\int_{t_j}^{t_{j+1}} \Phi_q({}_0D_t^\beta u)_0 D_t^\beta x dt + \int_{t'_i}^{t'_{i+1}} \Phi_q({}_0D_t^\beta v)_0 D_t^\beta y dt \right) \\
& = \int_{t_j}^{t_{j+1}} F_u(t, u, v) x dt + \int_{t'_i}^{t'_{i+1}} F_v(t, u, v) y dt. \tag{2.8}
\end{aligned}$$

By using Definitions 2.1, 2.2 and Proposition 2.4 of [30], we have

$$\begin{aligned}
& [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \left(\int_{t_j}^{t_{j+1}} \Phi_p({}_0D_t^\alpha u)_0 D_t^\alpha x dt + \int_{t'_i}^{t'_{i+1}} \Phi_p({}_0D_t^\alpha v)_0 D_t^\alpha y dt \right) \\
& + [M_2(\|(u, v)\|_\beta^q)]^{q-1} \left(\int_{t_j}^{t_{j+1}} \Phi_q({}_0D_t^\beta u)_0 D_t^\beta x dt + \int_{t'_i}^{t'_{i+1}} \Phi_q({}_0D_t^\beta v)_0 D_t^\beta y dt \right) \\
& = [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \left(\int_{t_j}^{t_{j+1}} {}_tD_T^\alpha(\Phi_p({}_0D_t^\alpha u)) x dt + \int_{t'_i}^{t'_{i+1}} {}_tD_T^\alpha(\Phi_p({}_0D_t^\alpha v)) y dt \right) \\
& + [M_2(\|(u, v)\|_\beta^q)]^{q-1} \left(\int_{t_j}^{t_{j+1}} {}_tD_T^\beta(\Phi_q({}_0D_t^\beta u)) x dt + \int_{t'_i}^{t'_{i+1}} {}_tD_T^\beta(\Phi_q({}_0D_t^\beta v)) y dt \right). \tag{2.9}
\end{aligned}$$

Hence, we can obtain from (2.8) and (2.9) that

$$\begin{aligned}
& [M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_tD_T^\alpha \Phi_p({}_0D_t^\alpha u(t)) + [M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_tD_T^\beta \Phi_q({}_0D_t^\beta u(t)) \\
& = F_u(t, u(t), v(t)), \quad \forall t \in (t_j, t_{j+1}), \tag{2.10}
\end{aligned}$$

and

$$\begin{aligned}
& [M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_tD_T^\alpha \Phi_p({}_0D_t^\alpha v(t)) + [M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_tD_T^\beta \Phi_q({}_0D_t^\beta v(t)) \\
& = F_v(t, u(t), v(t)), \quad \forall t \in (t'_i, t'_{i+1}), \tag{2.11}
\end{aligned}$$

which shows that (u, v) satisfies the equation of problem (1.1) a.e. on $[0, T] \setminus \{t_1, \dots, t_m\} \times [0, T] \setminus \{t'_1, \dots, t'_n\}$. Moreover, according to Proposition 2.6 of [30] and $M_1, M_2 \in C^1(R_0^+, R_0^+)$, we have

$$[M_1(\|(u(t_j^\pm), v(t_j^\pm))\|_\alpha^p)]^{p-1} {}_tD_T^{\alpha-1} \Phi_p({}_0D_t^\alpha u(t_j^\pm))$$

$$\begin{aligned}
& + [M_2(\|(u(t_j^\pm), v(t_j^\pm))\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta u(t_j^\pm)) \\
= & \lim_{t \rightarrow t_j^\pm} [M_1(\|(u(t), v(t))\|_\alpha^p)]^{p-1} {}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha u(t)) \\
& + [M_2(\|(u(t), v(t))\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta u(t)),
\end{aligned}$$

and

$$\begin{aligned}
& [M_1(\|(u(t_i^\pm), v(t_i^\pm))\|_\alpha^p)]^{p-1} {}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha v(t_i^\pm)) \\
& + [M_2(\|(u(t_i^\pm), v(t_i^\pm))\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta v(t_i^\pm)) \\
= & \lim_{t \rightarrow t_i^\pm} [M_1(\|(u(t), v(t))\|_\alpha^p)]^{p-1} {}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha v(t)) \\
& + [M_2(\|(u(t), v(t))\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta v(t))
\end{aligned}$$

exist. Multiplying (2.10) by x and multiplying (2.11) by y . Then, integrating between 0 and T , we have

$$\begin{aligned}
& [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \int_0^T {}_t D_T^\alpha (\Phi_p({}_0 D_t^\alpha u))x + {}_t D_T^\alpha (\Phi_p({}_0 D_t^\alpha v))y dt \\
& + [M_2(\|(u, v)\|_\beta^q)]^{q-1} \int_0^T {}_t D_T^\beta (\Phi_q({}_0 D_t^\beta u))x + {}_t D_T^\beta (\Phi_q({}_0 D_t^\beta v))y dt \\
= & \int_0^T F_u(t, u, v)x + F_v(t, u, v)y dt.
\end{aligned} \tag{2.12}$$

On the basis of Proposition 2.5 in [30], we have

$$\begin{aligned}
& [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \int_0^T {}_t D_T^\alpha (\Phi_p({}_0 D_t^\alpha u))x + {}_t D_T^\alpha (\Phi_p({}_0 D_t^\alpha v))y dt \\
& + [M_2(\|(u, v)\|_\beta^q)]^{q-1} \int_0^T {}_t D_T^\beta (\Phi_q({}_0 D_t^\beta u))x + {}_t D_T^\beta (\Phi_q({}_0 D_t^\beta v))y dt \\
= & \sum_{j=1}^m \left[\Delta \left([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha u) \right) (t_j) \right. \\
& \left. + \Delta \left([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta u) \right) (t_j) \right] x(t_j) \\
& + \sum_{i=1}^n \left[\Delta \left([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha v) \right) (t'_i) \right. \\
& \left. + \Delta \left([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta v) \right) (t'_i) \right] y(t'_i) \\
& + [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \int_0^T \Phi_p({}_0 D_t^\alpha u)_0 D_t^\alpha x + \Phi_p({}_0 D_t^\alpha v)_0 D_t^\alpha y dt \\
& + [M_2(\|(u, v)\|_\beta^q)]^{q-1} \int_0^T \Phi_q({}_0 D_t^\beta u)_0 D_t^\beta x + \Phi_q({}_0 D_t^\beta v)_0 D_t^\beta y dt
\end{aligned} \tag{2.13}$$

Hence, combining with (2.5), (2.12) and (2.13), we have

$$\sum_{j=1}^m \left[\Delta \left([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha u) \right) (t_j) \right]$$

$$\begin{aligned}
& + \Delta \left([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta u) \right) (t_j) + \lambda \zeta(t_j) I_j(u(t_j)) \Big] x(t_j) \\
& + \sum_{i=1}^n \left[\Delta \left([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha v) \right) (t'_i) \right. \\
& \left. + \Delta \left([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta v) \right) (t'_i) + \mu \varrho(t'_i) S_i(v(t'_i)) \right] y(t'_i) \\
& = 0.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
& \Delta([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha u))(t_j) \\
& + \Delta([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta u))(t_j) + \lambda \zeta(t_j) I_j(u(t_j)) = 0, \quad j \in \Delta_1,
\end{aligned}$$

and

$$\begin{aligned}
& \Delta([M_1(\|(u, v)\|_\alpha^p)]^{p-1} {}_t D_T^{\alpha-1} \Phi_p({}_0 D_t^\alpha v))(t'_i) \\
& + \Delta([M_2(\|(u, v)\|_\beta^q)]^{q-1} {}_t D_T^{\beta-1} \Phi_q({}_0 D_t^\beta v))(t'_i) + \mu \varrho(t'_i) S_i(v(t'_i)) = 0, \quad i \in \Delta_2,
\end{aligned}$$

which means that (u, v) satisfies the impulsive conditions of problem (1.1). Meanwhile, since $(u, v) \in E_{\alpha, \beta}$, it follows that (u, v) also satisfies the boundary conditions of problem (1.1). We complete the proof. \square

Definition 2.2. Let $c \in R$ and $J_{\lambda, \mu} \in C^1(E_{\alpha, \beta}, R)$. The functional $J_{\lambda, \mu}$ satisfies the $(PS)_c$ condition if any sequence $\{(u_n, v_n)\} \subset E_{\alpha, \beta}$ such that $J_{\lambda, \mu}(u_n, v_n) \rightarrow c$ and $J'_{\lambda, \mu}(u_n, v_n) \rightarrow 0$ as $n \rightarrow \infty$ in $E_{\alpha, \beta}^*$, where $E_{\alpha, \beta}^*$ is the dual space of $E_{\alpha, \beta}$, has a convergent subsequence in $E_{\alpha, \beta}$.

3. Existence of nontrivial solutions

In this section, we introduce the Nehari manifold for problem (1.1) to analyze the behavior of $J_{\lambda, \mu}$. It is assumed, without further mention, that the structural assumptions required in Theorem 1.1 are satisfied.

Now, we define the Nehari manifold associated with functional problem $J_{\lambda, \mu}$ as

$$\mathcal{N}_{\lambda, \mu} = \left\{ (u, v) \in E_{\alpha, \beta} \setminus \{(0, 0)\} : \langle J'_{\lambda, \mu}(u, v), (u, v) \rangle = 0 \right\}.$$

It is easy to see that $(u, v) \in \mathcal{N}_{\lambda, \mu}$ if and only if

$$\begin{aligned}
& [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \|(u, v)\|_\alpha^p + [M_2(\|(u, v)\|_\beta^q)]^{q-1} \|(u, v)\|_\beta^q \\
& - \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) - \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \\
& - \sigma \int_0^T F(t, u(t), v(t)) dt = 0. \tag{3.1}
\end{aligned}$$

Specifically, we derive several topological properties of the $\mathcal{N}_{\lambda, \mu}$, which are expressed as

$$\mathcal{N}_{\lambda, \mu} = \left\{ (u, v) \in E_{\alpha, \beta} \setminus \{(0, 0)\} : \mathcal{J}'_{u, v}(1) = 0 \right\}$$

$$= \left\{ (su, sv) \in E_{\alpha, \beta} \setminus \{(0, 0)\} : \mathcal{J}'_{u,v}(s) = 0 \right\},$$

where $\mathcal{J}_{u,v} : [0, +\infty) \rightarrow R$ is the fibering map given by $\mathcal{J}_{u,v}(s) = J_{\lambda, \mu}(su, sv)$ for any $s \geq 0$ and $(u, v) \in E_{\alpha, \beta}$, i.e.

$$\begin{aligned} \mathcal{J}_{u,v}(s) &= \frac{1}{p} \overline{M}_1(s^p \|(u, v)\|_{\alpha}^p) + \frac{1}{q} \overline{M}_2(s^q \|(u, v)\|_{\beta}^q) \\ &\quad - \sum_{j=1}^m \int_0^{su(t_j)} \lambda \zeta(t_j) I_j(z) dz - \sum_{i=1}^n \int_0^{sv(t'_i)} \mu \varrho(t'_i) S_i(z) dz \\ &\quad - s^{\sigma} \int_0^T F(t, u(t), v(t)) dt. \end{aligned} \quad (3.2)$$

Therefore, for any $(u, v) \in E_{\alpha, \beta}$, we have

$$\begin{aligned} \mathcal{J}'_{u,v}(s) &= [M_1(s^p \|(u, v)\|_{\alpha}^p)]^{p-1} s^{p-1} \|(u, v)\|_{\alpha}^p + [M_2(s^q \|(u, v)\|_{\beta}^q)]^{q-1} s^{q-1} \|(u, v)\|_{\beta}^q \\ &\quad - s^{\theta-1} \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) - s^{\theta-1} \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \\ &\quad - \sigma s^{\sigma-1} \int_0^T F(t, u(t), v(t)) dt, \end{aligned} \quad (3.3)$$

and

$$\begin{aligned} \mathcal{J}''_{u,v}(s) &= s^{p-2} (p-1) [M_1(s^p \|(u, v)\|_{\alpha}^p)]^{p-1} \|(u, v)\|_{\alpha}^p \\ &\quad + s^{q-2} (q-1) [M_2(s^q \|(u, v)\|_{\beta}^q)]^{q-1} \|(u, v)\|_{\beta}^q \\ &\quad + s^{2p-2} p(p-1) [M_1(s^p \|(u, v)\|_{\alpha}^p)]^{p-2} M'_1(s^p \|(u, v)\|_{\alpha}^p) \|(u, v)\|_{\alpha}^{2p} \\ &\quad + s^{2q-2} q(q-1) [M_2(s^q \|(u, v)\|_{\beta}^q)]^{q-2} M'_2(s^q \|(u, v)\|_{\beta}^q) \|(u, v)\|_{\beta}^{2q} \\ &\quad - (\theta-1) s^{\theta-2} \left(\sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \right) \\ &\quad - \sigma(\sigma-1) s^{\sigma-2} \int_0^T F(t, u(t), v(t)) dt. \end{aligned} \quad (3.4)$$

Then, for any $(u, v) \in \mathcal{N}_{\lambda, \mu}$, by (3.3), we have

$$\begin{aligned} \mathcal{J}'_{u,v}(1) &= [M_1(\|(u, v)\|_{\alpha}^p)]^{p-1} \|(u, v)\|_{\alpha}^p + [M_2(\|(u, v)\|_{\beta}^q)]^{q-1} \|(u, v)\|_{\beta}^q \\ &\quad - \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) - \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \\ &\quad - \sigma \int_0^T F(t, u(t), v(t)) dt, \end{aligned}$$

and by (3.1) and (3.4), we have

$$\begin{aligned} \mathcal{J}''_{u,v}(1) &= (p-1) [M_1(\|(u, v)\|_{\alpha}^p)]^{p-1} \|(u, v)\|_{\alpha}^p + (q-1) [M_2(\|(u, v)\|_{\beta}^q)]^{q-1} \|(u, v)\|_{\beta}^q \\ &\quad + p(p-1) [M_1(\|(u, v)\|_{\alpha}^p)]^{p-2} M'_1(\|(u, v)\|_{\alpha}^p) \|(u, v)\|_{\alpha}^{2p} \\ &\quad + q(q-1) [M_2(\|(u, v)\|_{\beta}^q)]^{q-2} M'_2(\|(u, v)\|_{\beta}^q) \|(u, v)\|_{\beta}^{2q} \end{aligned}$$

$$\begin{aligned}
& -(\theta - 1) \left(\sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \right) \\
& - \sigma(\sigma - 1) \int_0^T F(t, u(t), v(t)) dt \\
& = (p - \theta) [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \|(u, v)\|_\alpha^p + (q - \theta) [M_2(\|(u, v)\|_\beta^q)]^{q-1} \|(u, v)\|_\beta^q \\
& \quad + p(p - 1) [M_1(\|(u, v)\|_\alpha^p)]^{p-2} M_1'(\|(u, v)\|_\alpha^p) \|(u, v)\|_\alpha^{2p} \\
& \quad + q(q - 1) [M_2(\|(u, v)\|_\beta^q)]^{q-2} M_2'(\|(u, v)\|_\beta^q) \|(u, v)\|_\beta^{2q} \\
& - \sigma(\sigma - \theta) \int_0^T F(t, u(t), v(t)) dt \\
& = (p - \sigma) [M_1(\|(u, v)\|_\alpha^p)]^{p-1} \|(u, v)\|_\alpha^p + (q - \sigma) [M_2(\|(u, v)\|_\beta^q)]^{q-1} \|(u, v)\|_\beta^q \\
& \quad + p(p - 1) [M_1(\|(u, v)\|_\alpha^p)]^{p-2} M_1'(\|(u, v)\|_\alpha^p) \|(u, v)\|_\alpha^{2p} \\
& \quad + q(q - 1) [M_2(\|(u, v)\|_\beta^q)]^{q-2} M_2'(\|(u, v)\|_\beta^q) \|(u, v)\|_\beta^{2q} \\
& - (\theta - \sigma) \left(\sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \right). \tag{3.5}
\end{aligned}$$

Following the methodology developed by [25], $\mathcal{N}_{\lambda, \mu}$ is partitioned into the following three constituent elements:

$$\mathcal{N}_{\lambda, \mu}^\pm = \left\{ (u, v) \in \mathcal{N}_{\lambda, \mu} \mid \mathcal{J}_{u, v}''(1) \geq 0 \right\}, \text{ and } \mathcal{N}_{\lambda, \mu}^0 = \left\{ (u, v) \in \mathcal{N}_{\lambda, \mu} \mid \mathcal{J}_{u, v}''(1) = 0 \right\}.$$

Lemma 3.1. *For any $(\lambda, \mu) \in \Theta^0$, we have $\mathcal{N}_{\lambda, \mu}^0 = \emptyset$, where*

$$\Theta^0 = \left\{ (\lambda, \mu) \in R^2 \setminus \{(0, 0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \leq Q_0 \right\},$$

and

$$Q_0 = \left(\frac{\min\{\mathcal{B}_\alpha^*, \mathcal{B}_\beta^*\}}{2^\kappa(\sigma - \theta)\mathcal{A}_{\alpha, p}^\theta} \right)^{\frac{p}{p-\theta}} \left(\frac{\min\{\mathcal{B}_\alpha, \mathcal{B}_\beta\} 2^{1-\kappa}}{\sigma(\sigma - \theta)C_*\overline{\mathcal{M}}(\sigma)} \right)^{\frac{\kappa-\theta}{\sigma-\kappa} \cdot \frac{p}{p-\theta}}.$$

Proof. We debate through contradiction. Suppose there exists a pair $(\lambda, \mu) \in \Theta^0$ such that $\mathcal{N}_{\lambda, \mu}^0 \neq \emptyset$; in this scenario, we can distinguish between two cases.

CaseI: $(u, v) \in \mathcal{N}_{\lambda, \mu}^0$ and $\sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) = 0$. Evidently, considering (3.5) in conjunction with assumption (M1), it can be inferred that

$$\begin{aligned}
0 & < \left((p - \sigma)\xi_1 + p(p - 1) \frac{\delta_1}{\eta_1} \right) \|(u, v)\|_\alpha^\kappa + \left((q - \sigma)\xi_2 + q(q - 1) \frac{\delta_2}{\eta_2} \right) \|(u, v)\|_\beta^\kappa \\
& = -\mathcal{B}_\alpha^* \|(u, v)\|_\alpha^\kappa - \mathcal{B}_\beta^* \|(u, v)\|_\beta^\kappa \\
& < 0,
\end{aligned}$$

which is a contradiction, where $\mathcal{B}_\alpha^* = (\sigma - p)\xi_1 + p(1 - p) \frac{\delta_1}{\eta_1} > 0$, $\mathcal{B}_\beta^* = (\sigma - q)\xi_2 + q(1 - q) \frac{\delta_2}{\eta_2} > 0$, due to $\sigma > p^2, q^2$, $(\sigma - p)\xi_1 \eta_1 > (p^2 - p)\delta_1$ and $(\sigma - q)\xi_2 \eta_2 > (q^2 - q)\delta_2$.

CaseII: $(u, v) \in \mathcal{N}_{\lambda, \mu}^0$ and $\sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \neq 0$. Considering (M1) and (3.5), it can be concluded that

$$\sigma(\sigma - \theta) \int_0^T F(t, u(t), v(t)) dt > \left((p - \theta)\xi_1 + p(p - 1) \frac{\xi_1}{\eta_1} \right) \|(u, v)\|_\alpha^\kappa$$

$$\begin{aligned}
& + \left((q - \theta)\xi_2 + q(q - 1)\frac{\xi_2}{\eta_2} \right) \|(u, v)\|_{\beta}^{\kappa} \\
& \geq \min\{\mathcal{B}_{\alpha}, \mathcal{B}_{\beta}\} 2^{1-\kappa} \|(u, v)\|_{\alpha, \beta}^{\kappa}. \tag{3.6}
\end{aligned}$$

where $\mathcal{B}_{\alpha} = (p - \theta)\xi_1 + p(p - 1)\frac{\xi_1}{\eta_1}$, $\mathcal{B}_{\beta} = (q - \theta)\xi_2 + q(q - 1)\frac{\xi_2}{\eta_2}$. Moreover, by (F2) and (2.4), we have

$$\begin{aligned}
\sigma(\sigma - \theta) \int_0^T F(t, u(t), v(t)) dt & \leq \sigma(\sigma - \theta) C_* \|(u, v)\|_{\sigma}^{\sigma} \\
& \leq \sigma(\sigma - \theta) C_* \overline{\mathcal{M}}(\sigma) \|(u, v)\|_{\alpha, \beta}^{\sigma}. \tag{3.7}
\end{aligned}$$

Consequently, we can deduce from (3.6) and (3.7) that

$$\|(u, v)\|_{\alpha, \beta} > \left(\frac{\min\{\mathcal{B}_{\alpha}, \mathcal{B}_{\beta}\} 2^{1-\kappa}}{\sigma(\sigma - \theta) C_* \overline{\mathcal{M}}(\sigma)} \right)^{\frac{1}{\sigma - \kappa}}. \tag{3.8}$$

On the other hand, by (M1) and (3.5), we have

$$\begin{aligned}
& (\sigma - \theta) \left(\sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \right) \\
& > \min\{\mathcal{B}_{\alpha}^*, \mathcal{B}_{\beta}^*\} 2^{1-\kappa} \|(u, v)\|_{\alpha, \beta}^{\kappa}. \tag{3.9}
\end{aligned}$$

In addition, according to (H1) and Hölder inequality, we have

$$\begin{aligned}
& \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \\
& \leq \left(m\bar{a}|\lambda| \|\zeta\|_{\infty} + n\bar{b}|\mu| \|\varrho\|_{\infty} \right) \mathcal{A}_{\alpha, p}^{\theta} (\|u\|_{\alpha, p}^{\theta} + \|v\|_{\alpha, p}^{\theta}) \\
& \leq \mathcal{A}_{\alpha, p}^{\theta} \left((m\bar{a}|\lambda| \|\zeta\|_{\infty} + n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \left((\|u\|_{\alpha, p}^{\theta} + \|v\|_{\alpha, p}^{\theta})^{\frac{p}{\theta}} \right)^{\frac{\theta}{p}} \\
& \leq 2\mathcal{A}_{\alpha, p}^{\theta} \left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \|(u, v)\|_{\alpha, \beta}^{\theta}. \tag{3.10}
\end{aligned}$$

Apparently, from (3.9) and (3.10), we have

$$\|(u, v)\|_{\alpha, \beta} < \left(\frac{2^{\kappa}(\sigma - \theta) \mathcal{A}_{\alpha, p}^{\theta} \left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}}}{\min\{\mathcal{B}_{\alpha}^*, \mathcal{B}_{\beta}^*\}} \right)^{\frac{1}{\kappa - \theta}}. \tag{3.11}$$

From (3.8) and (3.11), we get

$$\begin{aligned}
& (m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}} \\
& > \left(\frac{\min\{\mathcal{B}_{\alpha}^*, \mathcal{B}_{\beta}^*\}}{2^{\kappa}(\sigma - \theta) \mathcal{A}_{\alpha, p}^{\theta}} \right)^{\frac{p}{p-\theta}} \left(\frac{\min\{\mathcal{B}_{\alpha}, \mathcal{B}_{\beta}\} 2^{1-\kappa}}{\sigma(\sigma - \theta) C_* \overline{\mathcal{M}}(\sigma)} \right)^{\frac{\kappa - \theta}{\sigma - \kappa} \cdot \frac{p}{p - \theta}},
\end{aligned}$$

which contradicts $(\lambda, \mu) \in \Theta^0$. This completes the proof. \square

Lemma 3.2. *For any $(\lambda, \mu) \in \Theta^0$, the functional $J_{\lambda, \mu}(u, v)$ is coercive and bounded below on $\mathcal{N}_{\lambda, \mu}$.*

Proof. For any $(u, v) \in \mathcal{N}_{\lambda, \mu}$, we can deduce from (2.6), (3.1), (M1) and (3.10) that

$$\begin{aligned} J_{\lambda, \mu}(u, v) &= J_{\lambda, \mu}(u, v) - \frac{1}{\sigma} \langle J'_{\lambda, \mu}(u, v), (u, v) \rangle \\ &> \left(\frac{1}{p} \frac{\xi_1}{\vartheta_1} - \frac{1}{\sigma} \delta_1 \right) \| (u, v) \|_{\alpha}^{\kappa} + \left(\frac{1}{q} \frac{\xi_2}{\vartheta_2} - \frac{1}{\sigma} \delta_2 \right) \| (u, v) \|_{\beta}^{\kappa} \\ &\quad - \left(\frac{1}{\theta} + \frac{1}{\sigma} \right) 2\mathcal{A}_{\alpha, p}^{\theta} \left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \| (u, v) \|_{\alpha, \beta}^{\theta} \\ &\geq \min\{\mathcal{C}_{\alpha}, \mathcal{C}_{\beta}\} 2^{1-\kappa} \| (u, v) \|_{\alpha, \beta}^{\kappa} \\ &\quad - \left(\frac{1}{\theta} + \frac{1}{\sigma} \right) 2\mathcal{A}_{\alpha, p}^{\theta} \left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \| (u, v) \|_{\alpha, \beta}^{\theta}, \end{aligned}$$

where $\mathcal{C}_{\alpha} = \frac{1}{p} \frac{\xi_1}{\vartheta_1} - \frac{1}{\sigma} \delta_1 > 0$, $\mathcal{C}_{\beta} = \frac{1}{q} \frac{\xi_2}{\vartheta_2} - \frac{1}{\sigma} \delta_2 > 0$, considering $\xi_1 \sigma > \vartheta_1 \delta_1 p$, $\xi_2 \sigma > \vartheta_2 \delta_2 q$ and $\theta < \kappa$. This completes the proof. \square

In the following result, we illustrate that $\mathcal{N}_{\lambda, \mu}^{+}$ and $\mathcal{N}_{\lambda, \mu}^{-}$ are non-empty. In order to deal with sign-changing functions ζ and ϱ , we define

$$\begin{aligned} \mathfrak{A}^{\pm} &= \left\{ (u, v) \in E_{\alpha, \beta} \mid \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \geq 0 \right\}, \\ \mathfrak{B}^{\pm} &= \left\{ (u, v) \in E_{\alpha, \beta} \mid \int_0^T F(t, u(t), v(t)) dt \geq 0 \right\}. \end{aligned}$$

Lemma 3.3. (i) For any $(u, v) \in \mathfrak{A}^{+} \cap \mathfrak{B}^{+}$ and $(\lambda, \mu) \in \Lambda_1$, there exist positive constants $s_* = s_*(u, v)$, $s_{**} = s_{**}(u, v)$, $s^+ = s^+(u, v)$ and $s^- = s^-(u, v)$ with $0 < s^+ < s_* < s^- < s_{**}$ such that $(s^+u, s^+v) \in \mathcal{N}_{\lambda, \mu}^{+}$, $(s^-u, s^-v) \in \mathcal{N}_{\lambda, \mu}^{-}$ and

$$J_{\lambda, \mu}(s^+u, s^+v) = \min_{0 \leq s \leq s^-} J_{\lambda, \mu}(su, sv), \quad J_{\lambda, \mu}(s^-u, s^-v) = \max_{s_* \leq s \leq s_{**}} J_{\lambda, \mu}(su, sv).$$

(ii) For any $(u, v) \in \mathfrak{A}^{+} \cap \mathfrak{B}^{-}$, there exists a unique $s^+ = s^+(u, v) > 0$ such that $(s^+u, s^+v) \in \mathcal{N}_{\lambda, \mu}^{+}$ and

$$J_{\lambda, \mu}(s^+u, s^+v) = \min_{s \geq 0} J_{\lambda, \mu}(su, sv).$$

Proof. For given $(u, v) \in E_{\alpha, \beta} \setminus \{(0, 0)\}$, let us consider the function $\mathfrak{L}_{u, v}(s) : R^+ \rightarrow R$ as

$$\begin{aligned} \mathfrak{L}_{u, v}(s) &= s^{p-\theta} [M_1(s^p \|(u, v)\|_{\alpha}^p)]^{p-1} \|(u, v)\|_{\alpha}^p + s^{q-\theta} [M_2(s^q \|(u, v)\|_{\beta}^q)]^{q-1} \|(u, v)\|_{\beta}^q \\ &\quad - \sigma s^{\sigma-\theta} \int_0^T F(t, u(t), v(t)) dt. \end{aligned} \quad (3.12)$$

Then, for any $s > 0$, if $(su, sv) \in \mathcal{N}_{\lambda, \mu}$, we can infer that

$$\mathfrak{L}_{u, v}(s) = \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i).$$

By (3.12), we have

$$\mathfrak{L}'_{u, v}(s) = (p - \theta) s^{p-\theta-1} [M_1(s^p \|(u, v)\|_{\alpha}^p)]^{p-1} \|(u, v)\|_{\alpha}^p$$

$$\begin{aligned}
& + p(p-1)s^{2p-\theta-1}[M_1(s^p\|(u,v)\|_\alpha^p)]^{p-2}M_1'(s^p\|(u,v)\|_\alpha^p)\|(u,v)\|_\alpha^{2p} \\
& + (q-\theta)s^{q-\theta-1}[M_2(s^q\|(u,v)\|_\beta^q)]^{q-1}\|(u,v)\|_\beta^q \\
& + q(q-1)s^{2q-\theta-1}[M_2(s^q\|(u,v)\|_\beta^q)]^{q-2}M_2'(s^q\|(u,v)\|_\beta^q)\|(u,v)\|_\beta^{2q} \\
& - \sigma(\sigma-\theta)s^{\sigma-\theta-1}\int_0^T F(t,u(t),v(t))dt, \tag{3.13}
\end{aligned}$$

and so if $(su, sv) \in \mathcal{N}_{\lambda,\mu}$, then

$$\mathcal{J}''_{su,sv}(1) = s^{\theta+1}\mathfrak{L}'_{u,v}(s). \tag{3.14}$$

Hence, $(su, sv) \in \mathcal{N}_{\lambda,\mu}^+$ (or $\mathcal{N}_{\lambda,\mu}^-$) if and only if $\mathfrak{L}'_{u,v}(s) > 0$ (or < 0).

(i) Let $(u, v) \in \mathfrak{A}^+ \cap \mathfrak{B}^+$. We note that $\mathfrak{L}_{u,v}(0) = 0$, $\mathfrak{L}_{u,v}(s) \rightarrow -\infty$ as $s \rightarrow +\infty$. By (3.7), (M1) and (3.13), we have

$$\begin{aligned}
\mathfrak{L}'_{u,v}(s) & > \min\{\mathcal{B}_\alpha, \mathcal{B}_\beta\}2^{1-\kappa}\|(u,v)\|_{\alpha,\beta}^\kappa s^{\kappa-\theta-1} \\
& - (\sigma(\sigma-\theta)C_*\overline{\mathcal{M}}(\sigma)\|(u,v)\|_{\alpha,\beta}^\sigma)s^{\sigma-\theta-1}, \tag{3.15}
\end{aligned}$$

which implies $\lim_{s \rightarrow 0^+} \mathfrak{L}'_{u,v}(s) > 0$ and $\lim_{s \rightarrow \infty} \mathfrak{L}'_{u,v}(s) = -\infty$. Then, define the function $\mathcal{P}(s) : R^+ \rightarrow R$ as

$$\begin{aligned}
\mathcal{P}(s) & = \min\{\mathcal{B}_\alpha, \mathcal{B}_\beta\}2^{1-\kappa}\|(u,v)\|_{\alpha,\beta}^\kappa s^{\kappa-\theta-1} \\
& - (\sigma(\sigma-\theta)C_*\overline{\mathcal{M}}(\sigma)\|(u,v)\|_{\alpha,\beta}^\sigma)s^{\sigma-\theta-1},
\end{aligned}$$

Due to $\kappa < \sigma$, there exists a unique s_{\max} , where

$$s_{\max} = \left(\frac{\min\{\mathcal{B}_\alpha, \mathcal{B}_\beta\}2^{1-\kappa}(\kappa-\theta-1)}{\sigma(\sigma-\theta)C_*\overline{\mathcal{M}}(\sigma)\|(u,v)\|_{\alpha,\beta}^{\sigma-\kappa}(\sigma-\theta-1)} \right)^{\frac{1}{\sigma-\kappa}},$$

such that $\mathcal{P}(s)$ is increasing on $(0, s_{\max})$, decreasing on $(s_{\max}, +\infty)$. Hence, there exists a unique $s_0 > s_{\max}$ such that $\mathcal{P}(s_0) = 0$. Then, we can deduce that $\mathfrak{L}'_{u,v}(s)$ exists a minimal root $s_* \geq s_0$ such that $\mathfrak{L}'_{u,v}(s_*) = 0$, $\mathfrak{L}''_{u,v}(s_*) < 0$. Therefore, there exists $s_{**} > s_*$ such that $\mathfrak{L}_{u,v}$ is increasing on $(0, s_*)$, decreasing on (s_*, s_{**}) . Taking into account the sign of $\mathfrak{L}_{u,v}(s_{**})$, we discuss it in the following two cases.

CaseI: $\mathfrak{L}_{u,v}(s_{**}) \leq 0$. By (3.7), (3.12) and (M1), we have

$$\mathfrak{L}_{u,v}(s) \geq \min\{\xi_1, \xi_2\}2^{1-\kappa}s^{\kappa-\theta}\|(u,v)\|_{\alpha,\beta}^\kappa - \sigma s^{\sigma-\theta}C_*\overline{\mathcal{M}}(\sigma)\|(u,v)\|_{\alpha,\beta}^\sigma.$$

Let the function $\mathcal{P}_1(s) : R^+ \rightarrow R$ be defined by

$$\mathcal{P}_1(s) = \min\{\xi_1, \xi_2\}2^{1-\kappa}\|(u,v)\|_{\alpha,\beta}^\kappa s^{\kappa-\theta} - \sigma C_*\overline{\mathcal{M}}(\sigma)\|(u,v)\|_{\alpha,\beta}^\sigma s^{\sigma-\theta}.$$

Then, there exists unique

$$s_1 = \left(\frac{\min\{\xi_1, \xi_2\}2^{1-\kappa}(\kappa-\theta)}{\sigma C_*\overline{\mathcal{M}}(\sigma)\|(u,v)\|_{\alpha,\beta}^{\sigma-\kappa}(\sigma-\theta)} \right)^{\frac{1}{\sigma-\kappa}},$$

such that

$$\max_{s>0} \mathcal{P}_1(s) = \mathcal{P}_1(s_1)$$

$$= \min\{\xi_1, \xi_2\} 2^{1-\kappa} \|(u, v)\|_{\alpha, \beta}^\theta \frac{\sigma - \kappa}{\sigma - \theta} \left(\frac{\min\{\xi_1, \xi_2\} 2^{1-\kappa} (\kappa - \theta)}{\sigma C_* \overline{\mathcal{M}}(\sigma) (\sigma - \theta)} \right)^{\frac{\kappa - \theta}{\sigma - \kappa}}.$$

Hence, we have $\mathfrak{L}_{u, v}(s_*) > \max_{s > 0} \mathcal{P}_1(s) = \mathcal{P}_1(s_1) > 0$. Moreover, for any $(\lambda, \mu) \in \Theta_1^0$, where

$$\Theta_1^0 = \left\{ (\lambda, \mu) \in R^2 \setminus \{(0, 0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \leq Q_1 \right\},$$

and

$$Q_1 = \left(\frac{\min\{\xi_1, \xi_2\} 2^{-\kappa} (\sigma - \kappa)}{\mathcal{A}_{\alpha, \beta}^\theta (\sigma - \theta)} \right)^{\frac{p}{p-\theta}} \left(\frac{\min\{\xi_1, \xi_2\} 2^{1-\kappa} (\kappa - \theta)}{\sigma C_* \overline{\mathcal{M}}(\sigma) (\sigma - \theta)} \right)^{\frac{\kappa - \theta}{\sigma - \kappa} \cdot \frac{p}{p-\theta}}.$$

Then, for any $(\lambda, \mu) \in \Theta_1^0$, we have

$$\begin{aligned} 0 &< \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \\ &\leq 2\mathcal{A}_{\alpha, \beta}^\theta \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \|(u, v)\|_{\alpha, \beta}^\theta \\ &\leq \mathcal{P}_1(s_1). \end{aligned}$$

Up to now, there exist unique $s^+ < s_*$ and $s_* < s^- < s_{**}$ such that

$$\mathfrak{L}_{u, v}(s^+) = \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) = \mathfrak{L}_{u, v}(s^-).$$

That is, (s^+u, s^+v) , $(s^-u, s^-v) \in \mathcal{N}_{\lambda, \mu}$. Moreover, by (3.14), we have $(s^+u, s^+v) \in \mathcal{N}_{\lambda, \mu}^+$, $(s^-u, s^-v) \in \mathcal{N}_{\lambda, \mu}^-$. Further, considering

$$\mathcal{J}'_{u, v}(s) = s^{\theta-1} \left(\mathfrak{L}_{u, v}(s) - \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) - \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \right),$$

we have $\mathcal{J}'_{u, v}(s) < 0$ for all $s \in [0, s^+)$, $\mathcal{J}'_{u, v}(s) > 0$ for all $s \in (s^+, s^-)$ and $\mathcal{J}'_{u, v}(s) < 0$ for all $s \in [s^-, s_{**})$. Hence, we have

$$J_{\lambda, \mu}(s^+u, s^+v) = \min_{0 \leq s \leq s^-} J_{\lambda, \mu}(su, sv), \text{ and } J_{\lambda, \mu}(s^-u, s^-v) = \max_{s_* \leq s \leq s_{**}} J_{\lambda, \mu}(su, sv).$$

CaseII: $0 < \mathfrak{L}_{u, v}(s_{**}) < \mathfrak{L}_{u, v}(s_*)$. Then, there exists sufficiently small $\varepsilon \in (0, 2\mathcal{A}_{\alpha, \beta}^\theta \|(u, v)\|_{\alpha, \beta}^\theta Q_1^{\frac{p-\theta}{p}})$ such that for any $(\lambda, \mu) \in \Theta^1$, where

$$\begin{aligned} \Theta^1 &= \left\{ (\lambda, \mu) \in R^2 \setminus \{(0, 0)\} : \left(Q_1^{\frac{p-\theta}{p}} - \frac{\varepsilon}{2\mathcal{A}_{\alpha, \beta}^\theta \|(u, v)\|_{\alpha, \beta}^\theta} \right)^{\frac{p}{p-\theta}} \right. \\ &\quad \left. \leq (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \leq Q_1 \right\}, \end{aligned}$$

we have

$$\mathfrak{L}_{u, v}(s_{**}) \leq \mathcal{P}_1(s_1) - \varepsilon$$

$$\begin{aligned}
&\leq \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i) \\
&\leq 2\mathcal{A}_{\alpha,p}^\theta \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \|(u,v)\|_{\alpha,\beta}^\theta \\
&\leq \mathcal{P}_1(s_1).
\end{aligned}$$

As analyzed in case1, there exist unique $s^+ < s_*$ and $s_* < s^- < s_{**}$ such that $(s^+u, s^+v) \in \mathcal{N}_{\lambda,\mu}^+$ and $(s^-u, s^-v) \in \mathcal{N}_{\lambda,\mu}^-$. moreover, we have

$$J_{\lambda,\mu}(s^+u, s^+v) = \min_{0 \leq s \leq s^-} J_{\lambda,\mu}(su, sv), \text{ and } J_{\lambda,\mu}(s^-u, s^-v) = \max_{s_* \leq s \leq s_{**}} J_{\lambda,\mu}(su, sv).$$

(ii) Let $(u, v) \in \mathfrak{A}^+ \cap \mathfrak{B}^-$. Apparently, $\mathfrak{L}_{u,v}(0) = 0$, $\mathfrak{L}_{u,v}(s) \rightarrow +\infty$ as $s \rightarrow +\infty$. According to (3.13), we can deduce that $\mathfrak{L}'_{u,v}(s) > 0$ for all $s > 0$, that is $\mathfrak{L}_{u,v}$ is increasing on $(0, +\infty)$. Since, $(u, v) \in \mathfrak{A}^+$, there exists unique $s^+ > 0$ such that

$$\mathfrak{L}_{u,v}(s^+) = \sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j)) u(t_j) + \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i)) v(t'_i).$$

That is, $(s^+u, s^+v) \in \mathcal{N}_{\lambda,\mu}$. In addition, (3.14) implies that $(s^+u, s^+v) \in \mathcal{N}_{\lambda,\mu}^+$ and $J_{\lambda,\mu}(s^+u, s^+v) = \min_{s \geq 0} J_{\lambda,\mu}(su, sv)$. \square

Lemma 3.4. For any $(\lambda, \mu) \in \Theta^2$, then $c_{\lambda,\mu}^+ = \inf_{(u,v) \in \mathcal{N}_{\lambda,\mu}^+} J_{\lambda,\mu}(u, v) < 0$, where

$$\Theta^2 = \left\{ (\lambda, \mu) \in R^2 \setminus \{(0, 0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \leq Q_2 \right\},$$

and

$$Q_2 = \left(\frac{\max\{Q_1, Q_2\} 2^{1-\kappa}}{4\mathcal{A}_{\alpha,p}^\theta} \right)^{\frac{p}{p-\theta}} \|(u, v)\|_{\alpha,\beta}^{\frac{p(\kappa-\theta)}{p-\theta}}.$$

Proof. Let $(u, v) \in \mathcal{N}_{\lambda,\mu}^+$. By (3.5), we have

$$\begin{aligned}
&\sigma(\sigma - \theta) \int_0^T F(t, u(t), v(t)) dt \\
&< \left((p - \theta)\delta_1 + p(p - 1) \frac{\delta_1}{\eta_1} \right) \|(u, v)\|_\alpha^\kappa + \left((q - \theta)\delta_2 + p(p - 1) \frac{\delta_2}{\eta_2} \right) \|(u, v)\|_\beta^\kappa.
\end{aligned}$$

Then, from (2.6), (3.1), (M1), we have

$$\begin{aligned}
c_{\lambda,\mu}^+ &\leq J_{\lambda,\mu}(u, v) \\
&< \left(\frac{1}{p} \frac{\delta_1}{\vartheta_1} + \frac{1}{\theta\sigma} \left((p - \theta)\delta_1 + p(p - 1) \frac{\delta_1}{\eta_1} \right) - \frac{1}{\theta} \xi_1 \right) \|(u, v)\|_\alpha^\kappa \\
&\quad + \left(\frac{1}{q} \frac{\delta_2}{\vartheta_2} + \frac{1}{\theta\sigma} \left((q - \theta)\delta_2 + q(q - 1) \frac{\delta_2}{\eta_2} \right) - \frac{1}{\theta} \xi_2 \right) \|(u, v)\|_\beta^\kappa \\
&\quad + \frac{4}{\theta} \mathcal{A}_{\alpha,p}^\theta \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \|(u, v)\|_{\alpha,\beta}^\theta. \quad (3.16)
\end{aligned}$$

Taking into account that $p\vartheta_1, q\vartheta_2 > 1$, $\theta\sigma > 1$, $(2p + p^2)\delta_1\theta < \xi_1$, and $(2q + q^2)\delta_2\theta < \xi_2$, it follows from (3.16) that

$$c_{\lambda,\mu}^+ \leq J_{\lambda,\mu}(u, v) < -\frac{1}{\theta} \max\{Q_1, Q_2\} 2^{1-\kappa} \|(u, v)\|_{\alpha,\beta}^\kappa$$

$$\begin{aligned}
& + \frac{4}{\theta} \mathcal{A}_{\alpha,p}^\theta \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \|(u,v)\|_{\alpha,\beta}^\theta \\
& < 0,
\end{aligned}$$

where $\mathcal{Q}_1 = \xi_1 - (2p+p^2)\delta_1\theta$ and $\mathcal{Q}_2 = \xi_2 - (2q+q^2)\delta_2\theta$. This completes the proof. \square

Lemma 3.5. *Let $(u,v) \in \mathcal{N}_{\lambda,\mu}^+$ and $(\lambda,\mu) \in \Theta^0$, there is a number $\varepsilon > 0$ and a continuous function $\varphi : \Theta_\varepsilon(0,0) \subseteq E_{\alpha,\beta} \rightarrow R^+$ such that*

$$\varphi(0,0) = 1, \text{ and } \varphi(z,\bar{z})((u,v) - (z,\bar{z})) \in \mathcal{N}_{\lambda,\mu}^+,$$

where $\Theta_\varepsilon(0,0) = \{(z,\bar{z}) \in E_{\alpha,\beta} : \|(z,\bar{z})\|_{\alpha,\beta} < \varepsilon\}$.

Proof. To begin, we define a function $\Upsilon : R^+ \times E_{\alpha,\beta} \rightarrow R$ as follows:

$$\begin{aligned}
\Upsilon(s, (z,\bar{z})) &= s^p [M_1(s^p \|(u,v) - (z,\bar{z})\|_\alpha^p)]^{p-1} \|(u,v) - (z,\bar{z})\|_\alpha^p \\
& + s^q [M_2(s^q \|(u,v) - (z,\bar{z})\|_\beta^q)]^{q-1} \|(u,v) - (z,\bar{z})\|_\beta^q \\
& - s^\theta \left(\sum_{j=1}^m \lambda \zeta(t_j) I_j(u(t_j) - z(t_j))(u(t_j) - z(t_j)) \right. \\
& \left. - \sum_{i=1}^n \mu \varrho(t'_i) S_i(v(t'_i) - \bar{z}(t'_i))(v(t'_i) - \bar{z}(t'_i)) \right) \\
& - \sigma s^\sigma \int_0^T F(t, u - z, v - \bar{z}) dt.
\end{aligned}$$

Thanks to $(u,v) \in \mathcal{N}_{\lambda,\mu}^+$, we have $\Upsilon(1, (0,0)) = 0$ and $\frac{\partial \Upsilon}{\partial s}(1, (0,0)) > 0$. Applying the implicit function theorem at $(1, (0,0))$, we obtain that there exists $\varepsilon_0 > 0$ such that for any $(z,\bar{z}) \in E_{\alpha,\beta}$ with $\|(z,\bar{z})\|_{\alpha,\beta} < \varepsilon_0$, the equation $\Upsilon(s, (z,\bar{z})) = 0$ has a unique continuous solution $s = \varphi(z,\bar{z}) > 0$ and $\varphi(0,0) = 1$. Furthermore, we have

$$\langle \varphi'(0,0), (z,\bar{z}) \rangle = \frac{W_*}{\mathcal{J}_{u,v}''(1)}, \quad (3.17)$$

where

$$\begin{aligned}
W_* &= \left((p-1) [M_1(\|(u,v)\|_\alpha^p)]^{p-2} M_1'(\|(u,v)\|_\alpha^p) \|(u,v)\|_\alpha^p + [M_1(\|(u,v)\|_\alpha^p)]^{p-1} \right) \\
& \times p \int_0^T (\Phi_p({}_0D_t^\alpha u)_0 D_t^\alpha z + \Phi_p({}_0D_t^\alpha v)_0 D_t^\alpha \bar{z}) dt \\
& + \left((q-1) [M_2(\|(u,v)\|_\beta^q)]^{q-2} M_2'(\|(u,v)\|_\beta^q) \|(u,v)\|_\beta^q + [M_2(\|(u,v)\|_\beta^q)]^{q-1} \right) \\
& \times q \int_0^T (\Phi_q({}_0D_t^\beta u)_0 D_t^\beta z + \Phi_q({}_0D_t^\beta v)_0 D_t^\beta \bar{z}) dt \\
& - \sum_{j=1}^m \lambda \zeta(t_j) (I_j'(u(t_j)) z(t_j) u(t_j) + I_j(u(t_j)) z(t_j)) \\
& - \sum_{i=1}^n \mu \varrho(t'_i) (S_i'(v(t'_i)) \bar{z}(t'_i) v(t'_i) + S_i(v(t'_i)) \bar{z}(t'_i))
\end{aligned}$$

$$- \sigma \int_0^T F_u(t, u, v)z + F_v(t, u, v)\bar{z}dt,$$

and $\Upsilon(\varphi(z, \bar{z}), (z, \bar{z})) = 0$ for all $(z, \bar{z}) \in B_{\varepsilon_0}(0, 0)$, which means that $\varphi(z, \bar{z})((u, v) - (z, \bar{z})) \in \mathcal{N}_\lambda$ for all $(z, \bar{z}) \in \Theta_{\varepsilon_0}(0, 0)$. Moreover, we can choose $0 < \varepsilon < \varepsilon_0$ such that for any $(z, \bar{z}) \in E_{\alpha, \beta}$ with $\|(z, \bar{z})\|_{\alpha, \beta} \leq \varepsilon$ we have $\frac{\partial \Upsilon}{\partial s}(\varphi(z, \bar{z}), (z, \bar{z})) > 0$, which implies that $\varphi(z, \bar{z})((u, v) - (z, \bar{z})) \in \mathcal{N}_\lambda^+$ for all $(z, \bar{z}) \in \Theta_\varepsilon(0, 0)$. \square

Lemma 3.6. *If $(\lambda, \mu) \in \Theta^0$, then there exists a minimizing sequence $\{(u_n, v_n)\} \subset \mathcal{N}_{\lambda, \mu}$ such that*

$$J_{\lambda, \mu}(u_n, v_n) = c_{\lambda, \mu} + o_n(1), \quad \text{and} \quad J'_{\lambda, \mu}(u_n, v_n) = o_n(1), \quad (3.18)$$

where $c_{\lambda, \mu} = \inf_{(u, v) \in \mathcal{N}_{\lambda, \mu}} J_{\lambda, \mu}(u, v)$.

Proof. By Lemma 3.2 and Ekeland's variational principle, there exists a minimizing sequence $\{(u_n, v_n)\} \subset \mathcal{N}_{\lambda, \mu}$ for $J_{\lambda, \mu}$ such that

$$c_{\lambda, \mu} < J_{\lambda, \mu}(u_n, v_n) < c_{\lambda, \mu} + \frac{1}{n}, \quad (3.19)$$

and

$$J_{\lambda, \mu}(u_n, v_n) < J_{\lambda, \mu}(u, v) + \frac{1}{n} \|(u, v) - (u_n, v_n)\|_{\alpha, \beta}, \quad \forall (u, v) \in \mathcal{N}_{\lambda, \mu}. \quad (3.20)$$

apparently, by (3.19), we get (3.18)₁. Now, we show that $\|J'_{\lambda, \mu}(u_n, v_n)\|_{\alpha, \beta} \rightarrow 0$ as $n \rightarrow \infty$. By Lemma 3.5, we obtain the function $\varphi_n : \Theta_{\varepsilon_n}(0, 0) \rightarrow R$ such that $\varphi_n(z, \bar{z})((u_n, v_n) - (z, \bar{z})) \in \mathcal{N}_{\lambda, \mu}$ for all $(z, \bar{z}) \in \Theta_{\varepsilon_n}(0, 0)$. For fixed n , choose $0 < \varepsilon < \varepsilon_n$ and define $(z_\varepsilon, \bar{z}_\varepsilon) = \varepsilon \frac{(u, v)}{\|(u, v)\|_{\alpha, \beta}}$ with $(u, v) \in E_{\alpha, \beta} \setminus (0, 0)$. Set $(z_{1, \varepsilon}, z_{2, \varepsilon}) = \varphi_n(z_\varepsilon, \bar{z}_\varepsilon)((u_n, v_n) - (z_\varepsilon, \bar{z}_\varepsilon))$, then it is clear that $(z_{1, \varepsilon}, z_{2, \varepsilon}) \in \mathcal{N}_{\lambda, \mu}$. According to (3.20), we have

$$J_{\lambda, \mu}(z_{1, \varepsilon}, z_{2, \varepsilon}) - J_{\lambda, \mu}(u_n, v_n) \geq -\frac{1}{n} \|(z_{1, \varepsilon}, z_{2, \varepsilon}) - (u_n, v_n)\|_{\alpha, \beta}.$$

Applying the mean value theorem, we have

$$\begin{aligned} & \langle J'_{\lambda, \mu}(u_n, v_n), (z_{1, \varepsilon}, z_{2, \varepsilon}) - (u_n, v_n) \rangle + o_n(\|(z_{1, \varepsilon}, z_{2, \varepsilon}) - (u_n, v_n)\|_{\alpha, \beta}) \\ & \geq -\frac{1}{n} \|(z_{1, \varepsilon}, z_{2, \varepsilon}) - (u_n, v_n)\|_{\alpha, \beta}. \end{aligned}$$

Then,

$$\begin{aligned} & -\langle J'_{\lambda, \mu}(u_n, v_n), (z_\varepsilon, \bar{z}_\varepsilon) \rangle + (\varphi_n(z_\varepsilon, \bar{z}_\varepsilon) - 1) \langle J'_{\lambda, \mu}(u_n, v_n), (u_n, v_n) - (z_\varepsilon, \bar{z}_\varepsilon) \rangle \\ & \geq -\frac{1}{n} \|(z_{1, \varepsilon}, z_{2, \varepsilon}) - (u_n, v_n)\|_{\alpha, \beta} + o_n(\|(z_{1, \varepsilon}, z_{2, \varepsilon}) - (u_n, v_n)\|_{\alpha, \beta}). \end{aligned}$$

Considering $\langle J'_{\lambda, \mu}(z_{1, \varepsilon}, z_{2, \varepsilon}), (u_n, v_n) - (z_\varepsilon, \bar{z}_\varepsilon) \rangle = 0$, we get

$$\begin{aligned} & (\varphi_n(z_\varepsilon, \bar{z}_\varepsilon) - 1) \langle J'_{\lambda, \mu}(u_n, v_n) - J'_{\lambda, \mu}(z_{1, \varepsilon}, z_{2, \varepsilon}), (u_n, v_n) - (z_\varepsilon, \bar{z}_\varepsilon) \rangle \\ & - \varepsilon \langle J'_{\lambda, \mu}(u_n, v_n), \frac{(u, v)}{\|(u, v)\|_{\alpha, \beta}} \rangle \end{aligned}$$

$$\geq -\frac{1}{n} \| (z_{1,\epsilon}, z_{2,\epsilon}) - (u_n, v_n) \|_{\alpha,\beta} + o_n(\| (z_{1,\epsilon}, z_{2,\epsilon}) - (u_n, v_n) \|_{\alpha,\beta}).$$

Hence, we have

$$\begin{aligned} & \langle J'_{\lambda,\mu}(u_n, v_n), \frac{(u, v)}{\|(u, v)\|_{\alpha,\beta}} \rangle \\ & \leq \frac{1}{\epsilon n} \| (z_{1,\epsilon}, z_{2,\epsilon}) - (u_n, v_n) \|_{\alpha,\beta} + \frac{1}{\epsilon} o_n(\| (z_{1,\epsilon}, z_{2,\epsilon}) - (u_n, v_n) \|_{\alpha,\beta}) \\ & \quad + \frac{1}{\epsilon} (\varphi_n(z_\epsilon, \bar{z}_\epsilon) - 1) \langle J'_{\lambda,\mu}(u_n, v_n) - J'_{\lambda,\mu}(z_{1,\epsilon}, z_{2,\epsilon}), (u_n, v_n) - (z_\epsilon, \bar{z}_\epsilon) \rangle. \end{aligned} \quad (3.21)$$

Since

$$\| (z_{1,\epsilon}, z_{2,\epsilon}) - (u_n, v_n) \|_{\alpha,\beta} \leq \epsilon |\varphi_n(z_\epsilon, \bar{z}_\epsilon)| + |\varphi_n(z_\epsilon, \bar{z}_\epsilon) - 1| \| (u_n, v_n) \|_{\alpha,\beta},$$

and

$$\lim_{\epsilon \rightarrow 0} \frac{\varphi_n(z_\epsilon, \bar{z}_\epsilon) - 1}{\epsilon} \leq \|\varphi'_n(0, 0)\|_{\alpha,\beta},$$

passing to the limit $\epsilon \rightarrow 0^+$ in (3.21), there exists $K_0 > 0$ such that

$$\langle J'_{\lambda,\mu}(u_n, v_n), \frac{(u, v)}{\|(u, v)\|_{\alpha,\beta}} \rangle \leq \frac{K_0}{n} (1 + \|\varphi'_n(0, 0)\|_{\alpha,\beta}). \quad (3.22)$$

Below, we demonstrate that $\|\varphi'_n(0, 0)\|_{\alpha,\beta}$ is bounded. Arguing by contradiction, we assume that $\langle \varphi'_n(0, 0), (z, \bar{z}) \rangle \rightarrow \infty$ as $n \rightarrow \infty$. By (3.19) and Lemma 3.2, we have $\sup_n \|(u_n, v_n)\|_{\alpha,\beta} < \infty$. Then, from (3.17), the boundedness of $\{(u_n, v_n)\}$ and Hölder's inequality, there exists positive constant L_0 such that

$$\langle \varphi'_n(0, 0), (z, \bar{z}) \rangle = \frac{L_0(\|z\|_{\alpha,p} + \|\bar{z}\|_{\alpha,p} + \|z\|_{\beta,q} + \|\bar{z}\|_{\beta,q})}{\mathcal{J}''_{u_n, v_n}(1)},$$

which implies that there exists a subsequence $\{(u_n, v_n)\}$ such that $\mathcal{J}''_{u_n, v_n}(1) = o_n(1)$ as $n \rightarrow \infty$. Following the proof of Lemma 3.1, we have $(\lambda, \mu) \notin \Theta^0$, which gives a contradiction. \square

Lemma 3.7. *Let $\{(u_n, v_n)\} \subset \mathcal{N}_{\lambda,\mu}$. $J_{\lambda,\mu}$ satisfies the (PS)-condition at level $c_{\lambda,\mu}$.*

Proof. It follows from Lemma 3.6 that there exists $\{(u_n, v_n)\}$ be a (PS) $_{c_{\lambda,\mu}}$ sequence for $J_{\lambda,\mu}$, i.e.

$$J_{\lambda,\mu}(u_n, v_n) \rightarrow c_{\lambda,\mu}, \text{ and } J'_{\lambda,\mu}(u_n, v_n) \rightarrow 0 \text{ in } E_{\alpha,\beta}^*, \text{ as } n \rightarrow \infty.$$

Lemma 3.2 shows that $\{(u_n, v_n)\}$ is bounded in $E_{\alpha,\beta}$. Hence, there exists $(u, v) \in E_{\alpha,\beta}$ and a subsequence of $\{(u_n, v_n)\}_n$, still denoted by $\{(u_n, v_n)\}_n$ such that $(u_n, v_n) \rightharpoonup (u, v)$ in $E_{\alpha,\beta}$. Additionally, by Lemma 2.3, we have $(u_n, v_n) \rightarrow (u, v)$ uniformly in $(C([0, T], R))^2$, i.e. $\|(u_n, v_n) - (u, v)\|_\infty \rightarrow 0$ as $n \rightarrow \infty$.

Now, let $(\mathbf{a}, \mathbf{b}) \in E_{\gamma,v}$ be fixed, $0 < \gamma \leq 1$, $1 < v < \infty$, and denote by $\mathcal{S}_{\gamma,v}(\mathbf{a}, \mathbf{b})$ the linear functional on $E_{\gamma,v}$

$$\langle \mathcal{S}_{\gamma,v}(\mathbf{a}, \mathbf{b}), (\mathbf{c}_1, \mathbf{c}_2) \rangle = \int_0^T \Phi_v({}_0D_t^\gamma \mathbf{a}(t)) {}_0D_t^\gamma \mathbf{c}_1(t) + \Phi_v({}_0D_t^\gamma \mathbf{b}(t)) {}_0D_t^\gamma \mathbf{c}_2(t) dt,$$

for any $(\mathbf{c}_1, \mathbf{c}_2) \in E_{\gamma, v}$. By the Hölder inequality, we have

$$|\langle \mathcal{S}_{\gamma, v}(\mathbf{a}, \mathbf{b}), (\mathbf{c}_1, \mathbf{c}_2) \rangle| \leq \|\mathbf{a}\|_{\gamma, v}^{v-1} \|\mathbf{c}_1\|_{\gamma, v} + \|\mathbf{b}\|_{\gamma, v}^{v-1} \|\mathbf{c}_2\|_{\gamma, v}.$$

Thus, for any $(\mathbf{c}_1, \mathbf{c}_2) \in E_{\gamma, v}$, the linear functional $\mathcal{S}_{\gamma, v}(\mathbf{a}, \mathbf{b})$ is also continuous. Hence, combined with $(u_n, v_n) \rightharpoonup (u, v)$ in $E_{\alpha, \beta}$, we have

$$\langle \mathcal{S}_{\gamma, v}(u, v), (u_n - u, v_n - v) \rangle \rightarrow 0, \text{ as } n \rightarrow \infty. \quad (3.23)$$

Then, we can deduce that

$$\begin{aligned} o_n(1) &= \langle J'_{\lambda, \mu}(u_n, v_n) - J_\lambda(u, v), (u_n - u, v_n - v) \rangle \\ &= [M_1(\|(u_n, v_n)\|_\alpha^p)]^{p-1} \langle \mathcal{S}_{\alpha, p}(u_n, v_n) - \mathcal{S}_{\alpha, p}(u, v), (u_n - u, v_n - v) \rangle \\ &\quad + [M_2(\|(u_n, v_n)\|_\beta^q)]^{q-1} \langle \mathcal{S}_{\beta, q}(u_n, v_n) - \mathcal{S}_{\beta, q}(u, v), (u_n - u, v_n - v) \rangle \\ &\quad + ([M_1(\|(u_n, u_n)\|_\alpha^p)]^{p-1} - [M_1(\|(u, v)\|_\alpha^p)]^{p-1}) \langle \mathcal{S}_{\alpha, p}(u, v), (u_n - u, v_n - v) \rangle \\ &\quad + ([M_2(\|(u_n, u_n)\|_\beta^q)]^{q-1} - [M_2(\|(u, v)\|_\beta^q)]^{q-1}) \langle \mathcal{S}_{\beta, q}(u, v), (u_n - u, v_n - v) \rangle \\ &\quad - \sum_{j=1}^m \lambda \zeta(t_j) (I_j(u_n(t_j)) - I_j(u(t_j)))(u_n(t_j) - u(t_j)) \\ &\quad - \sum_{i=1}^n \mu \varrho(t'_i) (S_i(v_n(t'_i)) - S_i(v(t'_i)))(v_n(t'_i) - v(t'_i)) \\ &\quad - \int_0^T (F_{u_n}(t, u_n, v_n) - F_u(t, u, v))(u_n - u) dt \\ &\quad - \int_0^T (F_{v_n}(t, u_n, v_n) - F_v(t, u, v))(v_n - v) dt. \end{aligned} \quad (3.24)$$

Moreover,

$$\begin{aligned} &\left| \sum_{j=1}^m \lambda \zeta(t_j) (I_j(u_n(t_j)) - I_j(u(t_j)))(u_n(t_j) - u(t_j)) \right| \\ &\leq m \bar{a} |\lambda| \|\zeta\|_\infty (\sup_{n \in N} |u_n|^{\theta-1} + \|u\|_\infty^{\theta-1}) \|u_n - u\|_\infty \rightarrow 0, \text{ as } n \rightarrow \infty, \end{aligned} \quad (3.25)$$

$$\begin{aligned} &\left| \sum_{i=1}^n \mu \varrho(t'_i) (K_i(v_n(t'_i)) - K_i(v(t'_i)))(v_n(t'_i) - v(t'_i)) \right| \\ &\leq n \bar{b} |\mu| \|\varrho\|_\infty (\sup_{n \in N} |v_n|^{\theta-1} + \|v\|_\infty^{\theta-1}) \|v_n - v\|_\infty \rightarrow 0, \text{ as } n \rightarrow \infty, \end{aligned} \quad (3.26)$$

and

$$\begin{aligned} &\left| \int_0^T (F_{u_n}(t, u_n, v_n) - F_u(t, u, v))(u_n - u) \right. \\ &\quad \left. + (F_{v_n}(t, u_n, v_n) - F_v(t, u, v))(v_n - v) dt \right| \\ &\leq T \|F_{u_n}(t, u_n, v_n) - F_u(t, u, v)\| \|u_n - u\|_\infty \\ &\quad + T \|F_{v_n}(t, u_n, v_n) - F_v(t, u, v)\| \|v_n - v\|_\infty \rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned} \quad (3.27)$$

From (3.23)-(3.27), we have

$$\langle \mathcal{S}_{\alpha,p}(u_n, v_n) - \mathcal{S}_{\alpha,p}(u, v), (u_n - u, v_n - v) \rangle \rightarrow 0, \text{ as } n \rightarrow \infty, \quad (3.28)$$

and

$$\langle \mathcal{S}_{\beta,q}(u_n, v_n) - \mathcal{S}_{\beta,q}(u, v), (u_n - u, v_n - v) \rangle \rightarrow 0, \text{ as } n \rightarrow \infty. \quad (3.29)$$

To illustrate our findings, we require the following well-established Simon inequalities.

$$|d_1 - d_2|^h \leq \begin{cases} \overline{\mathcal{D}}(|d_1|^{h-2}d_1 - |d_2|^{h-2}d_2)(d_1 - d_2), h \geq 2, \\ \overline{\mathcal{D}}((|d_1|^{h-2}d_1 - |d_2|^{h-2}d_2)(d_1 - d_2))^{\frac{h}{2}}(|d_1|^h + |d_2|^h)^{\frac{2-h}{2}}, 1 < h < 2, \end{cases}$$

for all $d_1, d_2 \in \mathbb{R}$, where $\overline{\mathcal{D}}$ is positive constant.

When $1 < p < 2$, by (3.28) and the Hölder inequality, we have

$$\begin{aligned} \|(u_n - u, v_n - v)\|_{\alpha}^p &= \|u_n - u\|_{\alpha,p}^p + \|v_n - v\|_{\alpha,p}^p \\ &\leq \overline{\mathcal{D}} \left(\int_0^T (\Phi_p({}_0D_t^\alpha u_n) - \Phi_p({}_0D_t^\alpha u))({}_0D_t^\alpha u_n - {}_0D_t^\alpha u) dt \right)^{\frac{p}{2}} \\ &\quad \times \left(\int_0^T (|{}_0D_t^\alpha u_n|^p + |{}_0D_t^\alpha u|^p) dt \right)^{\frac{2-p}{2}} \\ &\quad + \overline{\mathcal{D}} \left(\int_0^T (\Phi_p({}_0D_t^\alpha v_n) - \Phi_p({}_0D_t^\alpha v))({}_0D_t^\alpha v_n - {}_0D_t^\alpha v) dt \right)^{\frac{p}{2}} \\ &\quad \times \left(\int_0^T (|{}_0D_t^\alpha v_n|^p + |{}_0D_t^\alpha v|^p) dt \right)^{\frac{2-p}{2}} \\ &\rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned} \quad (3.30)$$

When $p \geq 2$, by (3.28), we have

$$\begin{aligned} \|(u_n - u, v_n - v)\|_{\alpha}^p &= \|u_n - u\|_{\alpha,p}^p + \|v_n - v\|_{\alpha,p}^p \\ &\leq \overline{\mathcal{D}} \langle \mathcal{S}_{\alpha,p}(u_n, v_n) - \mathcal{S}_{\alpha,p}(u, v), (u_n - u, v_n - v) \rangle \\ &\rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned} \quad (3.31)$$

Hence, we have $\|(u_n - u, v_n - v)\|_{\alpha}^p \rightarrow 0$ as $n \rightarrow \infty$. Similarly, we can obtain $\|(u_n - u, v_n - v)\|_{\beta}^q \rightarrow 0$, as $n \rightarrow \infty$. Then, we have $(u_n, v_n) \rightarrow (u, v)$ as $n \rightarrow \infty$ in $E_{\alpha,\beta}$. This ends the proof. \square

Proof of Theorem 1.1. Let $(\lambda, \mu) \in \Theta_* = \Theta^0 \cap \Theta^1 \cap \Theta^2$. Thanks to the Lemma 3.6, we can consider a $(PS)_{c_{\lambda,\mu}^+}$ sequence $\{(u_n, v_n)\} \subset \mathcal{N}_{\lambda,\mu}^+ \subset \mathcal{N}_{\lambda,\mu}$ such that

$$J_{\lambda,\mu}(u_n, v_n) \rightarrow c_{\lambda,\mu}^+, \text{ and } J'_{\lambda,\mu}(u_n, v_n) \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Then, by Lemma 3.4 and 3.7, there exists $(u_{\lambda,\mu}, v_{\lambda,\mu}) \in \mathcal{N}_{\lambda,\mu}$ such that

$$J'_{\lambda,\mu}(u_{\lambda,\mu}, v_{\lambda,\mu}) = 0, \text{ and } J_{\lambda,\mu}(u_{\lambda,\mu}, v_{\lambda,\mu}) = c_{\lambda,\mu}^+ < 0.$$

Further, we show that $(u_{\lambda,\mu}, v_{\lambda,\mu}) \in \mathcal{N}_{\lambda,\mu}^+$. We debate through contradiction. Assume that $(u_{\lambda,\mu}, v_{\lambda,\mu}) \in \mathcal{N}_{\lambda,\mu}^-$. Considering (3.5), we have $(u_{\lambda,\mu}, v_{\lambda,\mu}) \in \mathfrak{B}^+$. Moreover, $(u_{\lambda,\mu}, v_{\lambda,\mu}) \in \mathcal{N}_{\lambda,\mu}$ and $J_{\lambda,\mu}(u_{\lambda,\mu}, v_{\lambda,\mu}) = c_{\lambda,\mu}^+ < 0$ imply $(u_{\lambda,\mu}, v_{\lambda,\mu}) \in \mathfrak{A}^+$.

Then, by Lemma 3.3, there exist $s^-(u_{\lambda,\mu}, v_{\lambda,\mu}) > s^+(u_{\lambda,\mu}, v_{\lambda,\mu}) > 0$ such that $(s^-u_{\lambda,\mu}, s^-v_{\lambda,\mu}) \in \mathcal{N}_{\lambda,\mu}^-$ and $(s^+u_{\lambda,\mu}, s^+v_{\lambda,\mu}) \in \mathcal{N}_{\lambda,\mu}^+$, which implies $s^- = 1$ and $s^+ < 1$. Hence, there exists $s_0 \in (s^+, s^-)$ such that

$$\begin{aligned} J_{\lambda,\mu}(s^+u_{\lambda,\mu}, s^+v_{\lambda,\mu}) &= \min_{0 \leq s \leq s^-} J_{\lambda,\mu}(su_{\lambda,\mu}, sv_{\lambda,\mu}) < J_{\lambda,\mu}(s_0u_{\lambda,\mu}, s_0v_{\lambda,\mu}) \\ &< J_{\lambda,\mu}(s^-u_{\lambda,\mu}, s^-v_{\lambda,\mu}) = J_{\lambda,\mu}(u_{\lambda,\mu}, v_{\lambda,\mu}) = c_{\lambda,\mu}^+. \end{aligned}$$

That's a contradiction. Thus, $(u_{\lambda,\mu}, v_{\lambda,\mu}) \in \mathcal{N}_{\lambda,\mu}^+$. Upon next, we show that $(u_{\lambda,\mu}, v_{\lambda,\mu})$ is a local minimizer of $J_{\lambda,\mu}$ in $E_{\alpha,\beta}$, which is a critical point of $J_{\lambda,\mu}$ in $E_{\alpha,\beta}$ (refer to reference [23]). Since $(u_{\lambda,\mu}, v_{\lambda,\mu}) \in \mathcal{N}_{\lambda,\mu}^+$, by Lemma 3.3, we have $s^+(u_{\lambda,\mu}, v_{\lambda,\mu}) = 1 < s_*(u_{\lambda,\mu}, v_{\lambda,\mu})$. Thus, by continuity of $(u, v) \mapsto s_*(u, v)$, for fixed $\epsilon_0 > 0$, there exists $\rho_0 = \rho_0(\epsilon_0) > 0$ such that $1 + \epsilon_0 < s_*((u_{\lambda,\mu}, v_{\lambda,\mu}) - (u, v))$ for all $\|(u, v)\|_{\alpha,\beta} < \rho_0$. Also, from Lemma 3.5, it is easy to see that for a given $\rho_1 > 0$, there exists a C^1 map $\varphi : \Theta_{\rho_1}(0, 0) \subseteq E_{\alpha,\beta} \rightarrow R^+$ such that $\varphi(u, v)((u_{\lambda,\mu}, v_{\lambda,\mu}) - (u, v)) \in \mathcal{N}_{\lambda,\mu}^+$ and $\varphi(0, 0) = 1$. Consequently, for any $0 < \rho_* = \min\{\rho_0, \rho_1\}$ and the uniqueness of zeros of fibering map, we have $s^+((u_{\lambda,\mu}, v_{\lambda,\mu}) - (u, v)) = \varphi(u, v) < 1 + \epsilon_0 < s_*((u_{\lambda,\mu}, v_{\lambda,\mu}) - (u, v))$ for all $\|(u, v)\|_{\alpha,\beta} < \rho_*$. Since $s_*((u_{\lambda,\mu}, v_{\lambda,\mu}) - (u, v)) > 1$, we obtain

$$\begin{aligned} J_{\lambda,\mu}(u_{\lambda,\mu}, v_{\lambda,\mu}) &\leq J_{\lambda,\mu}(s^+((u_{\lambda,\mu}, v_{\lambda,\mu}) - (u, v))((u_{\lambda,\mu}, v_{\lambda,\mu}) - (u, v))) \\ &\leq J_{\lambda,\mu}((u_{\lambda,\mu}, v_{\lambda,\mu}) - (u, v)), \end{aligned}$$

for all $\|(u, v)\|_{\alpha,\beta} < \rho_*$. This shows that $(u_{\lambda,\mu}, v_{\lambda,\mu})$ is a local minimizer of $J_{\lambda,\mu}$ in $E_{\alpha,\beta}$. Moreover, since $(0, 0) \notin \mathcal{N}_{\lambda,\mu}^+$, we have $(u_{\lambda,\mu}, v_{\lambda,\mu}) \neq (0, 0)$. Hence, $(u_{\lambda,\mu}, v_{\lambda,\mu})$ is a nontrivial solution of problem (1.1). \square

4. Existence of infinity many nontrivial solutions

In this section, we explore the multiplicity of solutions for problem (1.1) by applying the Krasnoselskii genus theory. First of all, we refer to [22] to recall some basic notions about Krasnoselskii genus.

Let X be a Banach space, and Ξ denote the family of sets $A \subset X \setminus \{0\}$, where A is closed in X and symmetric with respect to 0, that is if $a \in A$ implies $-a \in A$.

Definition 4.1. Let $A \in \Xi$. The Krasnoselskii genus $\bar{\gamma}(A)$ of A is defined as the least positive integer n such that there is an odd mapping $\bar{h} \in C(A, R^n \setminus \{0\})$ for all $a \in A$. If n does not exist, $\bar{\gamma}(A) = \infty$. Moreover, $\bar{\gamma}(\emptyset) = 0$.

Note that condition (M1') implies that

$$\bar{M}_1(x) \geq \bar{M}_1(1)x^{\bar{q}_1}, \quad \forall x \in [0, 1], \quad \text{and} \quad \bar{M}_2(y) \geq \bar{M}_2(1)y^{\bar{q}_2}, \quad \forall y \in [0, 1]. \quad (4.1)$$

Moreover, for any $\epsilon > 0$, condition (F1') gives the existence of $C_\epsilon > 0$ such that for a.e. $t \in [0, T]$

$$|F(t, u, v)| \leq \epsilon |(u, v)|^{\bar{q}_*} + C_\epsilon |(u, v)|^{\bar{q}}, \quad \text{for all } (u, v) \in R^2. \quad (4.2)$$

Considering the nonlocal characteristics of M_1 and M_2 , let us assume that $0 < \|(u, v)\|_{\alpha,\beta} \leq 1$. Consequently, based on (2.4), (2.6), (3.10), (4.1) and (4.2), we can deduce that

$$J_{\lambda,\mu}(u, v) \geq \frac{1}{p} \bar{M}_1(1) \|(u, v)\|_{\alpha}^{p\bar{q}_1} + \frac{1}{q} \bar{M}_2(1) \|(u, v)\|_{\beta}^{q\bar{q}_2}$$

$$\begin{aligned}
& - \frac{2}{\theta} \mathcal{A}_{\alpha,p}^\theta \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \|(u,v)\|_{\alpha,\beta}^\theta \\
& - \varepsilon \overline{\mathcal{M}}(\bar{\vartheta}_*) \|(u,v)\|_{\alpha,\beta}^{\bar{\vartheta}_*} - C_\varepsilon \overline{\mathcal{M}}(\bar{\eta}) \|(u,v)\|_{\alpha,\beta}^{\bar{\eta}} \\
& \geq \overline{M}_{1,2} \|(u,v)\|_{\alpha,\beta}^{\bar{\vartheta}_*} - \mathcal{M}_* \|(u,v)\|_{\alpha,\beta}^{\bar{\eta}} \\
& - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \|(u,v)\|_{\alpha,\beta}^\theta. \quad (4.3)
\end{aligned}$$

Then, we can choose $\varepsilon > 0$ sufficiently small such that $\overline{M}_{1,2} > 0$, where

$$\overline{M}_{1,2} = \min\left\{\frac{1}{p}\overline{M}_1(1), \frac{1}{q}\overline{M}_2(1)\right\}2^{1-\bar{\vartheta}_*} - \varepsilon \overline{\mathcal{M}}(\bar{\vartheta}_*), \quad \mathcal{A}_* = \frac{2}{\theta} \mathcal{A}_{\alpha,p}^\theta, \quad \mathcal{M}_* = C_\varepsilon \overline{\mathcal{M}}(\bar{\eta}).$$

Define a function $\mathcal{T}_{\lambda,\mu} : (0, +\infty) \rightarrow R$ as

$$\begin{aligned}
\mathcal{T}_{\lambda,\mu}(X) &= \overline{M}_{1,2} X^{\bar{\vartheta}_*} - \mathcal{M}_* X^{\bar{\eta}} \\
& - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} X^\theta,
\end{aligned}$$

then $J_{\lambda,\mu}(u,v) \geq \mathcal{T}_{\lambda,\mu}(\|(u,v)\|_{\alpha,\beta})$ for all $(u,v) \in E_{\alpha,\beta}$ with $0 < \|(u,v)\|_{\alpha,\beta} \leq 1$. Noticeably, $\mathcal{T}'_{\lambda,\mu}(X) = X^{\theta-1} \overline{\mathcal{T}}_{\lambda,\mu}(X)$, where

$$\begin{aligned}
\overline{\mathcal{T}}_{\lambda,\mu}(X) &= \overline{M}_{1,2} \bar{\vartheta}_* X^{\bar{\vartheta}_* - \theta} - \mathcal{M}_* \bar{\eta} X^{\bar{\eta} - \theta} \\
& - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \theta, \quad \forall X > 0.
\end{aligned}$$

Then, there exists a unique

$$X_0 = \left(\frac{\overline{M}_{1,2} \bar{\vartheta}_* (\bar{\vartheta}_* - \theta)}{\mathcal{M}_* \bar{\eta} (\bar{\eta} - \theta)} \right)^{\frac{1}{\bar{\eta} - \bar{\vartheta}_*}} > 0,$$

such that $\overline{\mathcal{T}}'_{\lambda,\mu}(X_0) = 0$. Namely, $\overline{\mathcal{T}}_{\lambda,\mu}(X)$ is increasing on $(0, X_0)$, decreasing on $(X_0, +\infty)$. Let $(\lambda, \mu) \in \Theta_0$, where

$$\Theta_0 = \left\{ (\lambda, \mu) \in R^2 \setminus \{(0,0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} < \Omega_0 \right\},$$

and

$$\Omega_0 = \left(\frac{\overline{M}_{1,2} \bar{\vartheta}_* \frac{\bar{\eta} - \bar{\vartheta}_*}{\bar{\eta} - \theta}}{\mathcal{A}_* \theta} \right)^{\frac{p}{p-\theta}} \left(\frac{\overline{M}_{1,2} \bar{\vartheta}_* (\bar{\vartheta}_* - \theta)}{\mathcal{M}_* \bar{\eta} (\bar{\eta} - \theta)} \right)^{\frac{\bar{\vartheta}_* - \theta}{\bar{\eta} - \bar{\vartheta}_*} \cdot \frac{p}{p-\theta}},$$

then we have

$$\begin{aligned}
\max \overline{\mathcal{T}}_{\lambda,\mu}(X) &= \overline{\mathcal{T}}_{\lambda,\mu}(X_0) \\
&= \overline{M}_{1,2} \bar{\vartheta}_* \frac{\bar{\eta} - \bar{\vartheta}_*}{\bar{\eta} - \theta} \left(\frac{\overline{M}_{1,2} \bar{\vartheta}_* (\bar{\vartheta}_* - \theta)}{\mathcal{M}_* \bar{\eta} (\bar{\eta} - \theta)} \right)^{\frac{\bar{\vartheta}_* - \theta}{\bar{\eta} - \bar{\vartheta}_*}} \\
& - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \theta > 0.
\end{aligned}$$

Hence, there exist $X_1 \in (0, X_0)$ and $X_2 \in (X_0, +\infty)$ such that $\mathcal{T}'_{\lambda,\mu}(X_1) = \mathcal{T}'_{\lambda,\mu}(X_2) = 0$. Taking into account that $\lim_{X \rightarrow 0^+} \overline{\mathcal{T}}_{\lambda,\mu}(X) < 0$ and $\lim_{X \rightarrow \infty} \overline{\mathcal{T}}_{\lambda,\mu}(X) = -\infty$,

we have $\mathcal{T}_{\lambda,\mu}(X)$ is decreasing on $(0, X_1)$, increasing on (X_1, X_2) , decreasing on $(X_2, +\infty)$. Moreover, let $(\lambda, \mu) \in \Theta_1$, where

$$\Theta_1 = \left\{ (\lambda, \mu) \in \mathbb{R}^2 \setminus \{(0, 0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} < \Omega_1 \right\},$$

and

$$\Omega_1 = \left(\frac{\bar{M}_{1,2} \frac{\bar{\eta} - \bar{\vartheta}_*}{\bar{\eta} - \theta}}{\mathcal{A}_*} \right)^{\frac{p}{p-\theta}} \left(\frac{\bar{M}_{1,2}(\bar{\vartheta}_* - \theta)}{\mathcal{M}_*(\bar{\eta} - \theta)} \right)^{\frac{\bar{\vartheta}_* - \theta}{\bar{\eta} - \bar{\vartheta}_*} \cdot \frac{p}{p-\theta}}.$$

Then, if we define a function $\mathcal{T}_{\lambda,\mu}^1 : (0, +\infty) \rightarrow \mathbb{R}$ as

$$\mathcal{T}_{\lambda,\mu}^1(X) = \bar{M}_{1,2} X^{\bar{\vartheta}_* - \theta} - \mathcal{M}_* X^{\bar{\eta} - \theta},$$

we can deduce that there exist unique $X_* = \left(\frac{\bar{M}_{1,2}(\bar{\vartheta}_* - \theta)}{\mathcal{M}_*(\bar{\eta} - \theta)} \right)^{\frac{1}{\bar{\eta} - \bar{\vartheta}_*}}$ such that

$$\max \mathcal{T}_{\lambda,\mu}^1(X) = \mathcal{T}_{\lambda,\mu}^1(X_*) = \bar{M}_{1,2} \frac{\bar{\eta} - \bar{\vartheta}_*}{\bar{\eta} - \theta} \left(\frac{\bar{M}_{1,2}(\bar{\vartheta}_* - \theta)}{\mathcal{M}_*(\bar{\eta} - \theta)} \right)^{\frac{\bar{\vartheta}_* - \theta}{\bar{\eta} - \bar{\vartheta}_*}} > 0.$$

Hence, for any $(\lambda, \mu) \in \Theta_1$, we have

$$\begin{aligned} \mathcal{T}_{\lambda,\mu}(X_2) &= \max_{X>0} \mathcal{T}_{\lambda,\mu}(X) \\ &\geq X_*^\theta \left(\mathcal{T}_{\lambda,\mu}^1(X_*) - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \right) \\ &> 0. \end{aligned}$$

At this point, we arrive at there exist $0 < K_0(\lambda, \mu) < K_1(\lambda, \mu)$ with $\mathcal{T}_{\lambda,\mu}(K_0(\lambda, \mu)) = \mathcal{T}_{\lambda,\mu}(K_1(\lambda, \mu)) = 0$. From the structure of $\mathcal{T}_{\lambda,\mu}(X)$, we have $\mathcal{T}_{\lambda,\mu}(X) \leq 0$ if $X \in (0, K_0(\lambda, \mu)]$, $\mathcal{T}_{\lambda,\mu}(X) > 0$ if $X \in (K_0(\lambda, \mu), K_1(\lambda, \mu))$, and $\mathcal{T}_{\lambda,\mu}(X) \leq 0$ if $X \in [K_1(\lambda, \mu), +\infty)$.

Lemma 4.1. $\lim_{(\lambda,\mu) \rightarrow (0,0)} K_0(\lambda, \mu) = 0$.

Proof. From $\mathcal{T}_{\lambda,\mu}(K_0(\lambda, \mu)) = 0$ and $\mathcal{T}'_{\lambda,\mu}(K_0(\lambda, \mu)) > 0$, we have

$$\begin{aligned} \bar{M}_{1,2} K_0(\lambda, \mu)^{\bar{\vartheta}_*} - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} K_0(\lambda, \mu)^\theta \\ - \mathcal{M}_* K_0(\lambda, \mu)^{\bar{\eta}} = 0, \end{aligned} \quad (4.4)$$

and

$$\begin{aligned} \bar{M}_{1,2} \bar{\vartheta}_* K_0(\lambda, \mu)^{\bar{\vartheta}_* - 1} - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \theta K_0(\lambda, \mu)^{\theta-1} \\ - \mathcal{M}_* \bar{\eta} K_0(\lambda, \mu)^{\bar{\eta}-1} > 0. \end{aligned} \quad (4.5)$$

Subsequently, based on (4.4) and (4.5), we have

$$K_0(\lambda, \mu) < \left(\frac{\bar{M}_{1,2}(\bar{\vartheta}_* - \theta)}{\mathcal{M}_*(\bar{\eta} - \theta)} \right)^{\frac{1}{\bar{\eta} - \bar{\vartheta}_*}}.$$

The uniform boundedness of $K_0(\lambda, \mu)$ with respect to (λ, μ) can be inferred. Consequently, we choose a sequence $\{(\lambda_n, \mu_n)\}_{n=1}^\infty$ with $(\lambda_n, \mu_n) \rightarrow (0, 0)$ as $n \rightarrow \infty$.

This choice ensures that $K_0(\lambda_n, \mu_n) \rightarrow K_{0*} \geq 0$ as $n \rightarrow \infty$. Then, (4.4) and (4.5) imply that

$$\overline{M}_{1,2} K_{0*}^{\overline{\vartheta}_*} - \mathcal{M}_* K_{0*}^{\overline{\eta}} = 0, \quad (4.6)$$

and

$$\overline{M}_{1,2} \overline{\vartheta}_* K_{0*}^{\overline{\vartheta}_* - 1} - \mathcal{M}_* \overline{\eta} K_{0*}^{\overline{\eta} - 1} \geq 0. \quad (4.7)$$

Combining (4.6) and (4.7), we have

$$\mathcal{M}_* (\overline{\eta} - \overline{\vartheta}_*) K_{0*}^{\overline{\eta}} \leq 0,$$

which implies that $K_{0*} = 0$. Considering the arbitrary nature of $\{(\lambda_n, \mu_n)\}$, we can deduce that $\lim_{(\lambda, \mu) \rightarrow (0, 0)} K_0(\lambda, \mu) = 0$. \square

Take $\aleph : R^+ \rightarrow [0, 1]$, nonincreasing and C^∞ with $\aleph(z) = 1$ if $z \in [0, K_0(\lambda, \mu)]$, $\aleph(z) = 0$ if $z \in [\min\{K_1(\lambda, \mu), 1\}, \infty)$. From Lemma 4.1, there exists $\Omega_2 > 0$ sufficiently small when $(\lambda, \mu) \in \Theta_2$ where

$$\Theta_2 = \left\{ (\lambda, \mu) \in R^2 \setminus \{(0, 0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} < \Omega_2 \right\},$$

such that $K_0(\lambda, \mu) < \min\{K_1(\lambda, \mu), 1\}$. Subsequently, for any $(\lambda, \mu) \in \overline{\Theta}$, where $\overline{\Theta} = \Theta_0 \cap \Theta_1 \cap \Theta_2$, we hereby introduce the truncated functional

$$\begin{aligned} \overline{J}_{\lambda, \mu}(u, v) &= \frac{1}{p} \overline{M}_1 (\| (u, v) \|_\alpha^p) + \frac{1}{q} \overline{M}_2 (\| (u, v) \|_\beta^q) \\ &\quad - \sum_{j=1}^m \int_0^{u(t_j)} \lambda \zeta(t_j) I_j(z) dz - \sum_{i=1}^n \int_0^{v(t'_i)} \mu \varrho(t'_i) S_i(s) ds \\ &\quad - \aleph(\| (u, v) \|_{\alpha, \beta}) \int_0^T F(t, u(t), v(t)) dt. \end{aligned} \quad (4.8)$$

According to (4.3), for any $0 < \| (u, v) \|_{\alpha, \beta} \leq 1$, we can observe that

$$\begin{aligned} \overline{J}_{\lambda, \mu}(u, v) &\geq \overline{M}_{1,2} \| (u, v) \|_{\alpha, \beta}^{\overline{\vartheta}_*} - \mathcal{M}_* \aleph(\| (u, v) \|_{\alpha, \beta}) \| (u, v) \|_{\alpha, \beta}^{\overline{\eta}} \\ &\quad - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \| (u, v) \|_{\alpha, \beta}^\theta. \end{aligned}$$

Define a function $\mathcal{S}_{\lambda, \mu} : (0, +\infty) \rightarrow R$ as

$$\begin{aligned} \mathcal{S}_{\lambda, \mu}(X) &= \overline{M}_{1,2} X^{\overline{\vartheta}_*} - \mathcal{M}_* \aleph(X) X^{\overline{\eta}} \\ &\quad - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} X^\theta, \end{aligned}$$

then $\overline{J}_{\lambda, \mu}(u, v) \geq \mathcal{S}_{\lambda, \mu}(\| (u, v) \|_{\alpha, \beta})$ for all $(u, v) \in E_{\alpha, \beta}$ with $0 < \| (u, v) \|_{\alpha, \beta} \leq 1$. Moreover, we can deduce that $\mathcal{S}_{\lambda, \mu}(X) \geq \mathcal{T}_{\lambda, \mu}(X)$ for all $X \geq 0$, $\mathcal{S}_{\lambda, \mu}(X) = \mathcal{T}_{\lambda, \mu}(X)$ for all $0 \leq X \leq K_0(\lambda, \mu)$.

Lemma 4.2. *For all $(\lambda, \mu) \in \Theta_{**}$, the following hold:*

- (i) *If $\overline{J}_{\lambda, \mu}(u, v) < 0$, then $\| (u, v) \|_{\alpha, \beta} < K_0(\lambda, \mu)$ and $J_{\lambda, \mu}(\overline{u}, \overline{v}) = \overline{J}_{\lambda, \mu}(\overline{u}, \overline{v})$ for all $(\overline{u}, \overline{v})$ in a sufficiently small neighborhood of (u, v) .*

(ii) $\bar{J}_{\lambda,\mu}(u, v)$ satisfies the local $(PS)_c$ condition for all $c < 0$.

Proof. (i) We distinguish two different cases.

Case1: $\min\{K_1(\lambda, \mu), 1\} = 1$. If $K_0(\lambda, \mu) \leq \|(u, v)\|_{\alpha,\beta} < 1$, then $\bar{J}_{\lambda,\mu}(u, v) \geq \mathcal{S}_{\lambda,\mu}(\|(u, v)\|_{\alpha,\beta}) \geq \mathcal{T}_{\lambda,\mu}(\|(u, v)\|_{\alpha,\beta}) \geq 0$. If $\|(u, v)\|_{\alpha,\beta} \geq 1$, by the definition of \aleph we know that $\aleph(\|(u, v)\|_{\alpha,\beta}) = 0$. Due to the impacts of M_1 and M_2 , we have classified Case 1 into four subcases.

Case1₁: $\|(u, v)\|_{\alpha,\beta} \geq 1$ with $\|(u, v)\|_{\alpha} \geq 1$ and $\|(u, v)\|_{\beta} \geq 1$. Then, by (M1'), (M2') with $\tau = \bar{\tau} = 1$ and (3.10), we have

$$\begin{aligned} \bar{J}_{\lambda,\mu}(u, v) &\geq \min\left\{\frac{1}{p\bar{\vartheta}_1}m_1^{p-1}, \frac{1}{q\bar{\vartheta}_2}m_2^{q-1}\right\}2^{1-\varsigma}\|(u, v)\|_{\alpha,\beta}^{\varsigma} \\ &\quad - \mathcal{A}_*\left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}}\right)^{\frac{p-\theta}{p}}\|(u, v)\|_{\alpha,\beta}^{\theta}, \end{aligned}$$

where $\varsigma = \min\{p, q\}$. Define the function $\mathcal{S}_1 : (0, +\infty) \rightarrow R$ as

$$\mathcal{S}_1(X) = \Lambda_{1*}X^{\varsigma} - \mathcal{A}_*\left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}}\right)^{\frac{p-\theta}{p}}X^{\theta}, \quad \forall X \geq 0,$$

where $\Lambda_{1*} = \min\{\frac{1}{p\bar{\vartheta}_1}m_1^{p-1}, \frac{1}{q\bar{\vartheta}_2}m_2^{q-1}\}2^{1-\varsigma}$. Undoubtedly, $\mathcal{S}_1(X)$ has a global minimum point at

$$X_{\min} = \left(\frac{\mathcal{A}_*\left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}}\right)^{\frac{p-\theta}{p}}\theta}{\Lambda_{1*}\varsigma}\right)^{\frac{1}{\varsigma-\theta}},$$

with

$$\begin{aligned} \min_{X>0} \mathcal{S}_1(X) &= \mathcal{S}_1(X_{\min}) \\ &= X_{\min}^{\theta} \mathcal{A}_*\left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}}\right)^{\frac{p-\theta}{p}} \left(\frac{\theta}{\varsigma} - 1\right) < 0. \end{aligned}$$

We point out that $\mathcal{S}_1(X) \geq 0$ if and only if $X \geq X_{\lambda,\mu}^1$, where

$$X_{\lambda,\mu}^1 = \left(\frac{\mathcal{A}_*\left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}}\right)^{\frac{p-\theta}{p}}}{\Lambda_{1*}}\right)^{\frac{1}{\varsigma-\theta}}.$$

Hence, choosing $(\lambda, \mu) \in \Theta_{1*}$, where

$$\begin{aligned} \Theta_{1*} &= \left\{(\lambda, \mu) \in R^2 \setminus \{(0, 0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}} \leq \left(\frac{\Lambda_{1*}}{\mathcal{A}_*}\right)^{\frac{p}{p-\theta}}\right\}, \end{aligned}$$

we have $\bar{J}_{\lambda,\mu}(u, v) \geq \mathcal{S}_1(\|(u, v)\|_{\alpha,\beta}) \geq 0$ for all $\|(u, v)\|_{\alpha,\beta} \geq 1$.

Case1₂: $\|(u, v)\|_{\alpha,\beta} \geq 1$ with $0 < \|(u, v)\|_{\alpha} < 1$ and $0 < \|(u, v)\|_{\beta} < 1$. Then, by (3.10) and (4.1), we have

$$\begin{aligned} \bar{J}_{\lambda,\mu}(u, v) &\geq \min\left\{\frac{1}{p}\bar{M}_1(1), \frac{1}{q}\bar{M}_2(1)\right\}2^{1-\bar{\vartheta}_*}\|(u, v)\|_{\alpha,\beta}^{\bar{\vartheta}_*} \\ &\quad - \mathcal{A}_*\left((m\bar{a}|\lambda| \|\zeta\|_{\infty})^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_{\infty})^{\frac{p}{p-\theta}}\right)^{\frac{p-\theta}{p}}\|(u, v)\|_{\alpha,\beta}^{\theta}. \end{aligned}$$

Define the function $\mathcal{S}_2 : (0, +\infty) \rightarrow R$ as

$$\mathcal{S}_2(X) = \Lambda_{2*} X^{\bar{\vartheta}^*} - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\bar{\theta}}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\bar{\theta}}} \right)^{\frac{p-\bar{\theta}}{p}} X^\theta, \quad \forall X \geq 0,$$

where $\Lambda_{2*} = \min\{\frac{1}{p}\bar{M}_1(1), \frac{1}{q}\bar{M}_2(1)\}2^{1-\bar{\vartheta}^*}$. Likewise, $\mathcal{S}_2(X) \geq 0$ if and only if $X \geq X_{\lambda,\mu}^2$, where

$$X_{\lambda,\mu}^2 = \left(\frac{\mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\bar{\theta}}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\bar{\theta}}} \right)^{\frac{p-\bar{\theta}}{p}}}{\Lambda_{2*}} \right)^{\frac{1}{\bar{\theta}^* - \bar{\theta}}}.$$

Hence, choosing $(\lambda, \mu) \in \Theta_{2*}$, where

$$\begin{aligned} & \Theta_{2*} \\ &= \left\{ (\lambda, \mu) \in R^2 \setminus \{(0, 0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\bar{\theta}}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\bar{\theta}}} \leq \left(\frac{\Lambda_{2*}}{\mathcal{A}_*} \right)^{\frac{p}{p-\bar{\theta}}} \right\}, \end{aligned}$$

we have $\bar{J}_{\lambda,\mu}(u, v) \geq \mathcal{S}_2(\|(u, v)\|_{\alpha,\beta}) \geq 0$ for all $\|(u, v)\|_{\alpha,\beta} \geq 1$.

Case1₃: $\|(u, v)\|_{\alpha,\beta} \geq 1$ with $\|(u, v)\|_\alpha \geq 1$ and $0 < \|(u, v)\|_\beta < 1$. Then, by (M1'), (M2') with $\tau = 1$ and (3.10), we have

$$\begin{aligned} & \bar{J}_{\lambda,\mu}(u, v) \\ & \geq \frac{1}{p\bar{\vartheta}_1} m_1^{p-1} \|(u, v)\|_\alpha^p - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\bar{\theta}}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\bar{\theta}}} \right)^{\frac{p-\bar{\theta}}{p}} \|(u, v)\|_\alpha^\theta. \end{aligned}$$

Define the function $\mathcal{S}_3 : (0, +\infty) \rightarrow R$ as

$$\mathcal{S}_3(X) = \Lambda_{3*} X^p - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\bar{\theta}}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\bar{\theta}}} \right)^{\frac{p-\bar{\theta}}{p}} X^\theta, \quad \forall X \geq 0,$$

where $\Lambda_{3*} = \frac{1}{p\bar{\vartheta}_1} m_1^{p-1}$. Then, $\mathcal{S}_3(X) \geq 0$ if and only if $X \geq X_{\lambda,\mu}^3$, where

$$X_{\lambda,\mu}^3 = \left(\frac{\mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\bar{\theta}}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\bar{\theta}}} \right)^{\frac{p-\bar{\theta}}{p}}}{\Lambda_{3*}} \right)^{\frac{1}{p-\bar{\theta}}}.$$

Hence, choosing $(\lambda, \mu) \in \Theta_{3*}$, where

$$\begin{aligned} & \Theta_{3*} \\ &= \left\{ (\lambda, \mu) \in R^2 \setminus \{(0, 0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\bar{\theta}}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\bar{\theta}}} \leq \left(\frac{\Lambda_{3*}}{\mathcal{A}_*} \right)^{\frac{p}{p-\bar{\theta}}} \right\}, \end{aligned}$$

we have $\bar{J}_{\lambda,\mu}(u, v) \geq \mathcal{S}_3(\|(u, v)\|_{\alpha,\beta}) \geq 0$ for all $\|(u, v)\|_{\alpha,\beta} \geq 1$.

Case1₄: $\|(u, v)\|_{\alpha,\beta} \geq 1$ with $0 < \|(u, v)\|_\alpha < 1$ and $\|(u, v)\|_\beta \geq 1$. Then, by (M1'), (M2') with $\bar{\tau} = 1$ and (3.10), we have

$$\begin{aligned} & \bar{J}_{\lambda,\mu}(u, v) \\ & \geq \frac{1}{q\bar{\vartheta}_2} m_2^{q-1} \|(u, v)\|_\beta^q - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\bar{\theta}}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\bar{\theta}}} \right)^{\frac{p-\bar{\theta}}{p}} \|(u, v)\|_\beta^\theta. \end{aligned}$$

Define the function $\mathcal{S}_4 : (0, +\infty) \rightarrow R$ as

$$\mathcal{S}_4(X) = \Lambda_{4*} X^q - \mathcal{A}_* \left((|\lambda| \|\zeta\|_{\gamma_*})^{\frac{p}{p-\bar{\theta}}} + (|\mu| \|\varrho\|_{\gamma_*})^{\frac{p}{p-\bar{\theta}}} \right)^{\frac{p-\bar{\theta}}{p}} X^\theta, \quad \forall X \geq 0,$$

where $\Lambda_{4*} = \frac{1}{q\bar{v}_2} m_2^{q-1}$. Then, $\mathcal{S}_4(X) \geq 0$ if and only if $X \geq X_{\lambda,\mu}^4$, where

$$X_{\lambda,\mu}^4 = \left(\frac{\mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}}}{\Lambda_{4*}} \right)^{\frac{1}{q-\theta}}.$$

Hence, choosing $(\lambda, \mu) \in \Theta_{4*}$, where

$$\begin{aligned} & \Theta_{4*} \\ &= \left\{ (\lambda, \mu) \in R^2 \setminus \{(0, 0)\} : 0 < (m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \leq \left(\frac{\Lambda_{4*}}{\mathcal{A}_*} \right)^{\frac{p}{p-\theta}} \right\}, \end{aligned}$$

we have $\bar{J}_{\lambda,\mu}(u, v) \geq \mathcal{S}_4(\|(u, v)\|_{\alpha,\beta}) \geq 0$ for all $\|(u, v)\|_{\alpha,\beta} \geq 1$.

Case2: $\min\{K_1(\lambda, \mu), 1\} = K_1(\lambda, \mu)$. If $K_0(\lambda, \mu) \leq \|(u, v)\|_{\alpha,\beta} < K_1(\lambda, \mu)$, then $\bar{J}_{\lambda,\mu}(u, v) \geq \mathcal{S}_{\lambda,\mu}(\|(u, v)\|_{\alpha,\beta}) \geq \mathcal{T}_{\lambda,\mu}(\|(u, v)\|_{\alpha,\beta}) \geq 0$. If $K_1(\lambda, \mu) \leq \|(u, v)\|_{\alpha,\beta} < 1$, considering $\aleph(\|(u, v)\|_{\alpha,\beta}) = 0$, we have

$$\begin{aligned} \bar{J}_{\lambda,\mu}(u, v) &\geq \bar{M}_{1,2} \|(u, v)\|_{\alpha,\beta}^{\bar{v}_*} \\ &\quad - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \|(u, v)\|_{\alpha,\beta}^\theta. \end{aligned}$$

Due to $\mathcal{T}_{\lambda,\mu}(K_1(\lambda, \mu)) = 0$, that is

$$\begin{aligned} & \bar{M}_{1,2} K_1(\lambda, \mu)^{\bar{v}_*} - \mathcal{A}_* \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} K_1(\lambda, \mu)^\theta \\ &= \mathcal{M}_* K_1(\lambda, \mu)^{\bar{\eta}} > 0, \end{aligned}$$

we have $\bar{J}_{\lambda,\mu}(u, v) > 0$ for all $K_1(\lambda, \mu) \leq \|(u, v)\|_{\alpha,\beta} < 1$. If $\|(u, v)\|_{\alpha,\beta} \geq 1$, arguing similarly to Case1₁-Case1₄, we have $\mathcal{J}_\lambda(u, v) \geq 0$.

Hence, from **Case1** and **Case2**, for all $(\lambda, \mu) \in \Theta_{**}$, where $\Theta_{**} = \bar{\Theta} \cap \Theta_{1*} \cap \dots \cap \Theta_{4*}$, we obtain $\bar{J}_{\lambda,\mu}(u, v) < 0$, then $\|(u, v)\|_{\alpha,\beta} < K_0(\lambda, \mu)$. Moreover, we observe that $J_{\lambda,\mu}(\bar{u}, \bar{v}) = \bar{J}_{\lambda,\mu}(\bar{u}, \bar{v})$ for all $\|(\bar{u}, \bar{v}) - (u, v)\|_{\alpha,\beta} < K_1(\lambda, \mu) - \|(u, v)\|_{\alpha,\beta}$.

(ii) Let $\{(u_n, v_n)\}$ be a (PS)-sequence for $\bar{J}_{\lambda,\mu}$ on the level $c < 0$, that is

$$\bar{J}_{\lambda,\mu}(u_n, v_n) \rightarrow c, \text{ and } \bar{J}'_{\lambda,\mu}(u_n, v_n) \rightarrow 0 \text{ in } E_{\alpha,\beta}^*, \text{ as } n \rightarrow \infty.$$

This implies that $\bar{J}_{\lambda,\mu}(u_n, v_n) < 0$ for $n \in N$ large enough. Then, by (i) of Lemma 4.2, we have $\|(u, v)\|_{\alpha,\beta} < K_0(\lambda, \mu)$ and $J_{\lambda,\mu}(u_n, v_n) = \bar{J}_{\lambda,\mu}(u_n, v_n) \rightarrow c < 0$ and $J'_{\lambda,\mu}(u_n, v_n) = \bar{J}'_{\lambda,\mu}(u_n, v_n) \rightarrow 0$ as $n \rightarrow \infty$. By virtue of the coerciveness of $\bar{J}_{\lambda,\mu}$ in $E_{\alpha,\beta}$, it can be inferred that $\{(u_n, v_n)\}$ is bounded in $E_{\alpha,\beta}$. Hence, there exists $(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}) \in E_{\alpha,\beta}$, $e_\alpha \geq 0$, $e_\beta \geq 0$ and a subsequence of $\{(u_n, v_n)\}_n$, still denoted by $\{(u_n, v_n)\}_n$ such that

$$(u_n, v_n) \rightharpoonup (\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}) \text{ in } E_{\alpha,\beta}, \|(u_n, v_n)\|_\alpha \rightarrow e_\alpha, \|(u_n, v_n)\|_\beta \rightarrow e_\beta, \text{ as } n \rightarrow \infty.$$

As a result of (F1'), we have

$$\begin{aligned} & \left| \int_0^T (F_{u_n}(t, u_n, v_n) - F_{\bar{u}_{\lambda,\mu}}(t, \bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}))(u_n - \bar{u}_{\lambda,\mu}) \right. \\ & \quad \left. + (F_{v_n}(t, u_n, v_n) - F_{\bar{v}_{\lambda,\mu}}(t, \bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}))(v_n - \bar{v}_{\lambda,\mu}) dt \right| \end{aligned}$$

$\rightarrow 0$, as $n \rightarrow \infty$.

Following a similar line of argument as presented in (3.23)-(3.26), we can deduce that

$$\begin{aligned}
o_n(1) &= \left\langle J'_{\lambda,\mu}(u_n, v_n) - J_\lambda(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}), (u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu}) \right\rangle \\
&= [M_1(\|(u_n, v_n)\|_\alpha^p)]^{p-1} \\
&\quad \times \left\langle \mathcal{S}_{\alpha,p}(u_n, v_n) - \mathcal{S}_{\alpha,p}(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}), (u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu}) \right\rangle \\
&\quad + [M_2(\|(u_n, v_n)\|_\beta^q)]^{q-1} \\
&\quad \times \left\langle \mathcal{S}_{\beta,q}(u_n, v_n) - \mathcal{S}_{\beta,q}(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}), (u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu}) \right\rangle. \quad (4.9)
\end{aligned}$$

Due to the degeneracy of M_1 and M_2 , we discuss them in the following three cases and conclude that only the first one holds.

Casei: $e_\alpha > 0$ and $e_\beta > 0$. It is apparent that condition (M2') implies that $M_1(e_\alpha^p) > 0$, $M_2(e_\beta^q) > 0$. Then, we have

$$\left\langle \mathcal{S}_{\alpha,p}(u_n, v_n) - \mathcal{S}_{\alpha,p}(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}), (u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu}) \right\rangle \rightarrow 0, \text{ as } n \rightarrow \infty, \quad (4.10)$$

$$\left\langle \mathcal{S}_{\beta,q}(u_n, v_n) - \mathcal{S}_{\beta,q}(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}), (u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu}) \right\rangle \rightarrow 0, \text{ as } n \rightarrow \infty. \quad (4.11)$$

Hence, arguing similarly to (3.30) and (3.31), we have

$$\|(u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu})\|_\alpha^p \rightarrow 0, \text{ and } \|(u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu})\|_\beta^q \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Additionally, by employing Lemma 2.4, we derive that

$$\|(u_n, v_n)\|_\alpha^p = \|(u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu})\|_\alpha^p + \|(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu})\|_\alpha^p + o_n(1), \quad (4.12)$$

$$\|(u_n, v_n)\|_\beta^q = \|(u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu})\|_\beta^q + \|(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu})\|_\beta^q + o_n(1). \quad (4.13)$$

Then, we have

$$\|(u_n, v_n)\|_\alpha \rightarrow \|(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu})\|_\alpha, \text{ and } \|(u_n, v_n)\|_\beta \rightarrow \|(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu})\|_\beta, \text{ as } n \rightarrow \infty.$$

Therefore, we get $(u_n, v_n) \rightarrow (\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu})$ as $n \rightarrow \infty$ in $E_{\alpha,\beta}$.

Caseii: $e_\alpha = 0$ and $e_\beta > 0$. The condition (M2') implies $M_1(e_\alpha^p) = 0$, $M_2(e_\beta^q) > 0$. Then, (4.11) is valid. Hence, we obtain $\|(u_n - \bar{u}_{\lambda,\mu}, v_n - \bar{v}_{\lambda,\mu})\|_\beta^q \rightarrow 0$ as $n \rightarrow \infty$. Moreover, it follows from (4.12) that $(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu}) = (0, 0)$. Then, according to equation (4.13), we can deduce that the norm of $\|(u_n, v_n)\|_\beta \rightarrow e_\beta = \|(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu})\|_\beta = 0 > 0$, which leads to a contradiction.

Caseiii: $e_\alpha > 0$ and $e_\beta = 0$. The condition (M2') implies $M_1(e_\alpha^p) > 0$, $M_2(e_\beta^q) = 0$. Consequently, (4.10) holds. Repeating the analysis procedure as in case Caseii yields $\|(u_n, v_n)\|_\alpha \rightarrow e_\alpha = \|(\bar{u}_{\lambda,\mu}, \bar{v}_{\lambda,\mu})\|_\alpha = 0 > 0$, which is a contradiction. \square

Lemma 4.3. *For each $n \in N$, there exists a real number $\bar{\varepsilon} = \bar{\varepsilon}(n) > 0$ such that $\bar{\gamma}(\bar{J}_{\lambda,\mu}^{-\bar{\varepsilon}}) \geq n$, where $\bar{J}_{\lambda,\mu}^{-\bar{\varepsilon}} = \{(u, v) \in E_{\alpha,\beta} : \bar{J}_{\lambda,\mu}(u, v) \leq -\bar{\varepsilon}\}$.*

Proof. By (F2'), there exists $W_0, W_1 > 0$ such that

$$F(t, u, v) \geq W_0(|u|^\bar{\varepsilon} + |v|^\bar{\varepsilon}) - W_1, \text{ for any } (u, v) \in R^2.$$

Consider a fixed positive integer n , and let $E_{\alpha,\beta}^n$ denote an n -dimensional subspace of $E_{\alpha,\beta}$. The fact that $E_{\alpha,\beta}^n$ is a space of finite dimension implies that all its norms in $E_{\alpha,\beta}^n$ are mutually equivalent. Subsequently, we define

$$C_*^1 = \inf \left\{ \int_0^T (|u|^{\bar{\zeta}} + |v|^{\bar{\zeta}}) dt \mid (u, v) \in E_{\alpha,\beta}^n, \|(u, v)\|_{\alpha,\beta} = 1 \right\} > 0.$$

Then, taking $(u, v) \in E_{\alpha,\beta}$ with $\|(u, v)\|_{\alpha,\beta} = 1$, for $0 < \xi < K_0(\lambda, \mu)$, we can deduce that

$$\begin{aligned} \bar{J}_{\lambda,\mu}(\xi u, \xi v) &\leq \frac{1}{p} \max_{0 < \iota_1 < 1} [M_1(\iota_1)]^{p-1} \xi^p + \frac{1}{q} \max_{0 < \iota_2 < 1} [M_2(\iota_2)]^{q-1} \xi^q \\ &\quad + \frac{\xi^\theta}{\theta} 2\mathcal{A}_{\alpha,p}^\theta \left((m\bar{a}|\lambda| \|\zeta\|_\infty)^{\frac{p}{p-\theta}} + (n\bar{b}|\mu| \|\varrho\|_\infty)^{\frac{p}{p-\theta}} \right)^{\frac{p-\theta}{p}} \\ &\quad - \xi^{\bar{\zeta}} W_0 C_*^1 + W_1 T. \end{aligned}$$

Given $0 < \bar{\zeta} < \theta < 1$, for any $\bar{\varepsilon} > 0$, there exists a sufficiently small $\xi \in (0, K_0(\lambda, \mu))$ such that $\bar{J}_{\lambda,\mu}(\xi u, \xi v) \leq -\bar{\varepsilon}$ holds true for all $(u, v) \in E_{\alpha,\beta}$ with $\|(u, v)\|_{\alpha,\beta} = 1$. Let $K_\xi = \{(u, v) \in E_{\alpha,\beta} \mid \|(u, v)\|_{\alpha,\beta} = \xi\}$, then $K_\xi \cap E_{\alpha,\beta}^n \subset \{(u, v) \in E_{\alpha,\beta} : \bar{J}_{\lambda,\mu}(u, v) \leq -\bar{\varepsilon}\}$. By the mapping property of the genus [22, Proposition 7.5], we have $\bar{\gamma}(\bar{J}_{\lambda,\mu}^{-\bar{\varepsilon}}) \geq \bar{\gamma}(K_\xi \cap E_{\alpha,\beta}^n) = n$. \square

Proof of Theorem 1.2. Let us commence by introducing some notation.

$$\begin{aligned} \Sigma_n &= \left\{ A \subset \Xi \setminus \{(0, 0)\}, \bar{\gamma}(A) \geq n \right\}, \quad c_n = \inf_{A \in \Sigma_n} \sup_{(u,v) \in A} \bar{J}_{\lambda,\mu}(u, v), \\ L_c &= \left\{ (u, v) \in E_{\alpha,\beta} \mid \bar{J}_{\lambda,\mu}(u, v) = c, \bar{J}'_{\lambda,\mu}(u, v) = 0 \right\}. \end{aligned}$$

Apparently, condition (F3') ensures that $\bar{J}_{\lambda,\mu}(u, v)$ is even and Lemma 4.3 reveals that $\bar{J}_{\lambda,\mu}^{-\bar{\varepsilon}} \in \Sigma_n$. Moreover, as $\bar{J}_{\lambda,\mu}(u, v)$ is bounded from below, it follows that $-\infty < c_n \leq -\bar{\varepsilon}(n) < 0$. The $(PS)_{c_n}$ -condition of $\bar{J}_{\lambda,\mu}(u, v)$ can be inferred from (ii) of Lemma 4.2. Thus, c_n is a critical value of $\bar{J}_{\lambda,\mu}(u, v)$ for any $n \in \mathbb{N}$.

The inclusion $\Sigma_{n+1} \subset \Sigma_n$ implies $c_n \leq c_{n+1}$. We proceed by considering two distinct scenarios.

CaseI: If $-\infty < c_1 < c_2 < \dots < c_n < c_{n+1} < \dots$, then $\bar{\gamma}(L_{c_n}) \geq 1$, indicating that $\{c_n\}$ represents a sequence of distinct negative critical values of $\bar{J}_{\lambda,\mu}(u, v)$.

CaseII: If there exists a positive integer n_0 such that $c = c_{n_0} = c_{n_0+1} = \dots = c_{n_0+N_*}$ for some $N_* \geq 1$, then it follows that $\bar{\gamma}(L_c) \geq N_* + 1$. Assuming a contradiction, we hypothesize that $c = c_{n_0} = c_{n_0+1} = \dots = c_{n_0+N_*} < 0$, then $\bar{\gamma}(L_c) \leq N_*$. Considering that $\bar{J}_{\lambda,\mu}$ satisfies $(PS)_c$ condition according to Lemma 4.2, we obtain that L_c is a compact set and $\bar{\gamma}(L_c) \leq N_* < \infty$. According to the continuity property of the genus, there exists $\delta^* > 0$ such that $\bar{\gamma}(U_{\delta^*}) = \bar{\gamma}(L_c) \leq N_* < \infty$. Since $\bar{J}_{\lambda,\mu}$ is even, by the Deformation Theorem [22], there exists an odd homeomorphism $\mathcal{T} : E_{\alpha,\beta} \rightarrow E_{\alpha,\beta}$ such that $\mathcal{T}(\bar{J}_{\lambda,\mu}^{c+\bar{\varepsilon}_0} \setminus U_{\delta^*}) \subset \bar{J}_{\lambda,\mu}^{c-\bar{\varepsilon}_0}$ for some $0 < \bar{\varepsilon}_0 < -c$. In addition, owing to $c = c_{n_0+N_*} = \inf_{A \in \Sigma_{n_0+N_*}} \sup_{(u,v) \in A} \bar{J}_{\lambda,\mu}(u, v)$, there exists an $A \in \Sigma_{n_0+N_*}$ such that $\sup_{(u,v) \in A} \bar{J}_{\lambda,\mu}(u, v) = c < c + \bar{\varepsilon}_0$, i.e. $A \subset \bar{J}_{\lambda,\mu}^{c+\bar{\varepsilon}_0}$. Then, $\mathcal{T}(A \setminus U_{\delta^*}) \subset \mathcal{T}(\bar{J}_{\lambda,\mu}^{c+\bar{\varepsilon}_0} \setminus U_{\delta^*}) \subset \bar{J}_{\lambda,\mu}^{c-\bar{\varepsilon}_0}$, i.e.

$$\sup_{(u,v) \in \mathcal{T}(A \setminus U_{\delta^*})} \bar{J}_{\lambda,\mu}(u, v) \leq c - \bar{\varepsilon}_0. \quad (4.14)$$

Hence, we have $\overline{\gamma}(A \setminus \overline{U_{\delta^*}}) \geq \overline{\gamma}(A) - \overline{\gamma}(U_{\delta^*}) \geq n_0 + N_* - N_* = n_0$, i.e. $\mathcal{T}(\overline{A \setminus \overline{U_{\delta^*}}}) \in \Sigma_{n_0}$ and $\sup_{(u,v) \in \mathcal{T}(A \setminus \overline{U_{\delta^*}})} \overline{J}_{\lambda,\mu}(u,v) \geq c_{n_0} = c$, which contradicts (4.14). Consequently, it follows that $\overline{\gamma}(L_c) \geq N_* + 1 > 2$, which shows that L_c contains infinitely many point. Furthermore, $\overline{J}_{\lambda,\mu}(u,v) = c < 0 = \overline{J}_{\lambda,\mu}(0,0)$, then $(u,v) \neq (0,0)$. And also, it is established that $\overline{J}_{\lambda,\mu}(u,v) = J_{\lambda,\mu}(u,v)$ if $\overline{J}_{\lambda,\mu}(u,v) < 0$, then there are infinitely many nontrivial critical points of $J_{\lambda,\mu}(u,v)$. Therefore, problem (1.1) has infinitely many nontrivial solutions. \square

5. Examples

In this section we give two examples to illustrate the application of our results.

Example 5.1. We consider the following impulsive fractional differential equation:

$$\left\{ \begin{array}{l} 15\|(u,v)\|_{\alpha}^2 {}_t D_T^{\alpha}({}_0 D_t^{\alpha} u(t)) + 15.5\|(u,v)\|_{\beta}^2 {}_t D_T^{\beta}({}_0 D_t^{\beta} u(t)) \\ = (t+1)|(u,v)|^{500} u, \quad t \neq t_1, \quad \text{a.e. } t \in [0, T], \\ 15\|(u,v)\|_{\alpha}^2 {}_t D_T^{\alpha}({}_0 D_t^{\alpha} v(t)) + 15.5\|(u,v)\|_{\beta}^2 {}_t D_T^{\beta}({}_0 D_t^{\beta} v(t)) \\ = (t+1)|(u,v)|^{500} v, \quad t \neq t'_1, \quad \text{a.e. } t \in [0, T], \\ \Delta(15\|(u,v)\|_{\alpha}^2 {}_t D_T^{\alpha-1}({}_0 D_t^{\alpha} u))(t_1) \\ + \Delta(15.5\|(u,v)\|_{\beta}^2 {}_t D_T^{\beta-1}({}_0 D_t^{\beta} u))(t_1) + \lambda \zeta(t_1) \frac{|u(t_1)|^{0.002}}{u(t_1)} = 0, \\ \Delta(15\|(u,v)\|_{\alpha}^2 {}_t D_T^{\alpha-1}({}_0 D_t^{\alpha} v))(t'_1) \\ + \Delta(15.5\|(u,v)\|_{\beta}^2 {}_t D_T^{\beta-1}({}_0 D_t^{\beta} v))(t'_1) + \mu \varrho(t'_1) \frac{|v(t'_1)|^{0.002}}{v(t'_1)} = 0, \\ u(0) = u(T) = 0, \quad v(0) = v(T) = 0, \end{array} \right. \quad (5.1)$$

For this case, $M_1(x) = 15x$, $M_2(y) = 15.5y$, $p = q = 2$, $I_1(z) = S_1(z) = \frac{|z|^{0.002}}{z}$, $z \in \mathbb{R}$, $F(t, u, v) = (t+1) \frac{|(u,v)|^{502}}{502}$.

We verify that all the conditions of Theorem 1.1 are satisfied. Obviously, M_1 and M_2 satisfy (M1) with

$$\begin{aligned} 0.5x^2 &< 0.8 \times 7.5x^2 \leq 15x^2 \leq 1.2 \times 15x^2 < 20x^2, \\ 0.4y^2 &< 0.9 \times 7.75y^2 \leq 15.5y^2 \leq 1.4 \times 15.5y^2 < 22y^2, \end{aligned}$$

where $\xi_1 = 0.5$, $\vartheta_1 = 0.8$, $\eta_1 = 1.2$, $\delta_1 = 20$, $\sigma = 502$, $\theta = 0.002$, $\xi_2 = 0.4$, $\vartheta_2 = 0.9$, $\eta_2 = 1.4$, $\delta_2 = 22$, $\kappa = 4$; I_1 and S_1 satisfy (H1) and (H2) with

$$\begin{aligned} |I_1(z)z| &\leq 1.2|z|^{0.002}, \quad \text{and } |S_1(z)z| \leq 2.4|z|^{0.002}, \\ I_1(su) &= s^{0.002-1} I_1(u), \quad \text{and } S_1(sv) = s^{0.002-1} S_1(v), \end{aligned}$$

where $a_1 = 1.2$, $b_1 = 2.4$; F satisfies (F1) and (F2) with

$$\begin{aligned} F(t, su, sv) &= s^{502} F(t, u, v), \quad \text{and } uF_u(t, u, v) + vF_v(t, u, v) = 502F(t, u, v), \\ |F(t, u, v)| &\leq \frac{T+1}{502} |(u,v)|^{502}, \end{aligned}$$

where $C_* = \frac{T+1}{502}$. Thus, all the conditions of Theorem 1.1 are satisfied. Then, Theorem 1.1 implies that there exists $\Theta_* > 0$ such that for any $(\lambda, \mu) \in \Theta_*$, problem (5.1) admits at least a nontrivial solution.

Example 5.2. We consider the following impulsive fractional differential equation:

$$\left\{ \begin{array}{l} [a\|(u, v)\|_{\alpha}^3]^2 {}_t D_T^{\alpha} \Phi_3({}_0 D_t^{\alpha} u(t)) + [b\|(u, v)\|_{\beta}^3]^2 {}_t D_T^{\beta} \Phi_3({}_0 D_t^{\beta} u(t)) \\ = h(t)|(u, v)|^{12} u, \quad t \neq t_1, \quad \text{a.e. } t \in [0, T], \\ [a\|(u, v)\|_{\alpha}^3]^2 {}_t D_T^{\alpha} \Phi_3({}_0 D_t^{\alpha} v(t)) + [b\|(u, v)\|_{\beta}^3]^2 {}_t D_T^{\beta} \Phi_3({}_0 D_t^{\beta} v(t)) \\ = h(t)|(u, v)|^{12} v, \quad t \neq t'_1, \quad \text{a.e. } t \in [0, T], \\ \Delta([a\|(u, v)\|_{\alpha}^3]^2 {}_t D_T^{\alpha-1} \Phi_3({}_0 D_t^{\alpha} u))(t_1) \\ + \Delta([b\|(u, v)\|_{\beta}^3]^2 {}_t D_T^{\beta-1} \Phi_3({}_0 D_t^{\beta} u))(t_1) + \lambda \zeta(t_1) \bar{a} \frac{|u(t_1)|^{0.9}}{u(t_1)} = 0, \\ \Delta([a\|(u, v)\|_{\alpha}^3]^2 {}_t D_T^{\alpha-1} \Phi_3({}_0 D_t^{\alpha} v))(t'_1) \\ + \Delta([b\|(u, v)\|_{\beta}^3]^2 {}_t D_T^{\beta-1} \Phi_3({}_0 D_t^{\beta} v))(t'_1) + \mu \varrho(t'_1) \bar{b} \frac{|v(t'_1)|^{0.9}}{v(t'_1)} = 0, \\ u(0) = u(T) = 0, \quad v(0) = v(T) = 0, \end{array} \right. \quad (5.2)$$

where $a > 0$, $b > 0$, $\bar{a} > 0$, $\bar{b} > 0$, $0 \leq h \in C([0, T], \mathbb{R}^+)$. For this case, $p = q = 3$, $M_1(x) = ax$, $M_2(y) = by$, $F(t, u, v) = \frac{1}{13} h(t)|(u, v)|^{13}$, $I_1(s) = \bar{a} \frac{|s|^{0.9}}{s}$, $S_1(s) = \bar{b} \frac{|s|^{0.9}}{s}$.

We verify that all the conditions of Theorem 1.2 are satisfied.

(M1')

$$[M_1(x)]^2 x \leq 3\bar{M}_1(x), \quad \text{and} \quad [M_2(y)]^2 y \leq 4\bar{M}_2(y),$$

where $\bar{\vartheta}_1 = 3$, $\bar{\vartheta}_2 = 4$.

(M2') $M_1(0) = M_2(0) = 0$, and for any $\tau, \bar{\tau} > 0$, there exist $m_1 = m_1(\tau) > 0$ and $m_2 = m_2(\bar{\tau}) > 0$ such that

$$M_1(x) \geq m_1, \quad \text{for all } x \geq \tau, \quad \text{and} \quad M_2(y) \geq m_2, \quad \text{for all } y \geq \bar{\tau}.$$

(F1') There exist $\bar{\eta}$ with $\bar{\vartheta}_* = \max\{p\bar{\vartheta}_1, q\bar{\vartheta}_2\} < \bar{\eta}$ such that for any $\varepsilon > 0$, there exists $C_{\varepsilon} > 0$ for which

$$|\nabla F(t, u, v)| \leq \bar{\vartheta}_* \varepsilon |(u, v)|^{\bar{\vartheta}_*-1} + \bar{\eta} C_{\varepsilon} |(u, v)|^{\bar{\eta}-1}, \quad \text{for all } (u, v) \in \mathbb{R}^2$$

holds, where $|(u, v)| = (u^2 + v^2)^{\frac{1}{2}}$ and $\nabla F = (F_u, F_v)$.

(F2') There exists $0 < \bar{\zeta} < \theta < 1$ such that

$$0 \leq \bar{\zeta} F(t, u, v) \leq \nabla F(t, u, v) \cdot (u, v), \quad \text{for all } (u, v) \in \mathbb{R}^2.$$

Obviously, M_1 and M_2 satisfy (M1'), (M2') with

$$[M_1(x)]^2 x \leq 3\bar{M}_1(x), \quad \text{and} \quad [M_2(y)]^2 y \leq 4\bar{M}_2(y),$$

where $\bar{\vartheta}_1 = 3$, $\bar{\vartheta}_2 = 4$; F satisfies (F1')-(F3') with

$$\begin{aligned} |\nabla F(t, u, v)| &\leq 12\varepsilon |(u, v)|^{11} + 13C_{\varepsilon} |(u, v)|^{12}, \\ 0 &\leq 0.8F(t, u, v) \leq \nabla F(t, u, v) \cdot (u, v), \end{aligned}$$

where $\bar{\vartheta}_* = 12$, $\bar{\eta} = 13$, $\bar{\zeta} = 0.8$, $\theta = 0.9$; I_1 and S_1 satisfy (H1), (F3') with

$$|I_1(z)z| \leq 2\bar{a}|z|^{0.9}, \quad \text{and} \quad |S_1(z)z| \leq 2\bar{b}|z|^{0.9},$$

where $a_1 = 2\bar{a}$, $b_1 = 2\bar{b}$. Then, by Theorem 1.2, there exists $\Theta_* > 0$ such that for any $(\lambda, \mu) \in \Theta_{**}$, problem (5.2) admits infinitely many nontrivial solutions.

References

- [1] G. A. Afrouzi, S. Heidarkhani and S. Moradi, *Multiple solutions for a Kirchhoff-type second-order impulsive differential equation on the half-line*, Quaest. Math., 2022, 45(1), 109–141.
- [2] R. P. Agarwal, M. Benchohra and S. Hamani, *A survey on existence results for boundary value problems of nonlinear fractional differential equations and inclusions*, Acta. Appl. Math., 2010, 109, 973–1033.
- [3] M. F. Chaves, G. Ercole and O. H. Miyagaki, *Existence of a nontrivial solution for the (p, q) -Laplacian in R^N without the Ambrosetti-Rabinowitz condition*, Nonlinear Anal.-Theory Methods Appl., 2015, 114, 133–141.
- [4] T. Y. Chen and W. B. Liu, *Ground state solutions of Kirchhoff-type fractional Dirichlet problem with p -Laplacian*, Adv. Differ. Equ., 2018, 2018, 1–9.
- [5] T. Y. Chen, W. B. Liu and H. Jin, *Nontrivial solutions of the Kirchhoff-type fractional p -Laplacian Dirichlet problem*, J. Funct. Spaces, 2020, 2020, 1–8.
- [6] K. Diethelm, *The Analysis of Fractional Differential Equation*, Springer, New York, 2010.
- [7] C. Farkas, A. Fiscella and P. Winkert, *On a class of critical double phase problems*, J. Math. Anal. Appl., 2022, 515(2), 1–16.
- [8] A. Fiscella and P. K. Mishra, *The Nehari manifold for fractional Kirchhoff problems involving singular and critical terms*, Nonlinear Anal., 2019, 186, 6–32.
- [9] A. Fiscella and P. Pucci, *Degenerate Kirchhoff (p, q) -fractional systems with critical nonlinearities*, Fract. Calc. Appl. Anal., 2020, 23(3), 723–752.
- [10] J. R. Graef, S. Heidarkhani, L. J. Kong and S. Moradi, *Existence results for impulsive fractional differential equations with p -Laplacian via variational methods*, Bound. Value Probl., 2022, 147(1), 95–112.
- [11] S. Heidarkhani and A. Salari, *Nontrivial solutions for impulsive fractional differential systems through variational methods*, Math. Methods. Appl. Sci., 2020, 43(10), 6529–6541.
- [12] R. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific, Singapore, 2000.
- [13] F. Jiao and Y. Zhou, *Existence results for fractional boundary value problem via critical point theory*, Int. J. Bifurcation Chaos, 2012, 22(4), 1–17.
- [14] M. Kratou, *Ground state solutions of p -Laplacian singular Kirchhoff problem involving a Riemann-Liouville fractional derivative*, Filomat, 2019, 33(7), 2073–2088.
- [15] D. P. Li, F. Q. Chen and Y. K. An, *Positive solutions for a p -Laplacian type system of impulsive fractional boundary value problem*, J. Appl. Anal. Comput., 2020, 10(2), 740–759.
- [16] D. P. Li, F. Q. Chen and Y. K. An, *The existence of solutions for an impulsive fractional coupled system of (p, q) -Laplacian type without the Ambrosetti-Rabinowitz condition*, Math. Meth. Appl. Sci., 2019, 42: 1449–1464.

- [17] D. P. Li, F. Q. Chen and Y. K. An, *Variational formulation for nonlinear impulsive fractional differential equations with (p, q) -Laplacian operator*, Math. Meth. Appl. Sci., 2022, 45, 515–531.
- [18] N. Nyamoradi and S. Tersian, *Existence of solutions for nonlinear fractional order p -Laplacian differential equations via critical point theory*, Fract. Calc. Appl. Anal., 2019, 22(4), 945–967.
- [19] K. Perera, M. Squassina and Y. Yang, *Bifurcation and multiplicity results for critical fractional p -Laplacian problems*, Math. Nachr., 2016, 289(2-3), 332–342.
- [20] P. Pucci and L. Temperini, *Existence for fractional (p, q) systems with critical and Hardy terms in R^N* , Nonlinear Anal., 2021, 211, 1–33.
- [21] Y. Qiao, F. Q. Chen and Y. K. An, *Nontrivial solutions of a class of fractional differential equations with p -Laplacian via variational methods*, Bound. Value. Probl., 2020, 2020: 1–15.
- [22] R. H. Rabinowitz, *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, American Mathematical Society Providence, Rhode Island, 1986.
- [23] K. Saoudi, P. Agarwal, P. Kumam, A. Ghanmi and P. Thounthong, *The Nehari manifold for a boundary value problem involving Riemann-Liouville fractional derivative*, Adv. Differ. Equ., 2018, 2018, 1–18.
- [24] R. Steglański, *Infinitely many solutions for double phase problem with unbounded potential in R^N* , Nonlinear Anal., 2022, 214, 1–20.
- [25] G. Tarantello, *On nonhomogeneous elliptic involving critical Sobolev exponent*, Ann. Inst. Henri Poincaré-Analyse non linéaire, 1992, 9(3), 281–304.
- [26] L. Wang, K. Xie and B. L. Zhang, *Existence and multiplicity of solutions for critical Kirchhoff-type p -Laplacian problems*, J. Math. Anal. Appl., 2018, 458(1), 361–378.
- [27] M. Q. Xiang, D. Hu and D. Yang, *Least energy solutions for fractional Kirchhoff problems with logarithmic nonlinearity*, Nonlinear Anal., 2020, 198, 1–20.
- [28] M. Q. Xiang, B. L. Zhang and D. Repovš, *Existence and multiplicity of solutions for fractional Schrödinger-Kirchhoff equations with Trudinger-Moser nonlinearity*, Nonlinear Anal., 2019, 186, 74–98.
- [29] W. H. Xie and H. B. Chen, *Multiple positive solutions for the critical Kirchhoff type problems involving sign-changing weight functions*, J. Math. Anal. Appl., 2019, 479(1), 135–161.
- [30] Y. L. Zhao and L. Tang, *Multiplicity results for impulsive fractional differential equations with p -Laplacian via variational methods*, Bound. Value. Probl., 2017, 2017, 1–15.
- [31] J. W. Zhou, Y. Q. Liu, Y. N. Wang and J. F. Suo, *Solvability of nonlinear impulsive generalized fractional differential equations with (p, q) -Laplacian operator via critical point theory*, Fractal Fract., 2022, 6(12): 1–24.