

1 **ON THE STUDY TO A TYPE OF**
2 **SINGULARLY PERTURBED BOUNDARY**
3 **VALUE PROBLEM WITH TWO DOUBLE**
4 **ROOTS OF THE DEGENERATE EQUATION***

5 Ruixiang Li¹, Mingkang Ni^{1,2,†}

6 **Abstract** This paper addresses a singularly perturbed boundary value prob-
7 lem where the degenerate equation has three distinct roots: two double roots
8 and one simple root. It is shown that for a sufficiently small parameter, the
9 solution of the problem switches between the two double roots in a neighbor-
10 hood of the transition point. As a result, the inner layer can be divided into
11 multiple regions. An asymptotic expansion is constructed, and the existence
12 of smooth solutions is established. Additionally, an estimate for the remainder
13 term is provided.

14 **Keywords** Singular perturbations, Asymptotic theory, Multiple roots.

15 **MSC(2010)** 35A21, 35B25, 35C20.

16 **1. Introduction**

17 This paper discusses the important topic of singularly perturbed problems, specif-
18 ically contrast structures. The Tikhonov school in the former Soviet Union first
19 introduced the concept of contrast structures in the late 1990s. A contrast struc-
20 ture occurs in singular perturbation problems where the degenerate equation has
21 distinct roots, known as critical manifolds in geometric singular perturbation theory.
22 As the solution switches between these isolated roots, a complex solution structure
23 forms.

24 Currently, most research focuses on steptype contrast structures. The main
25 challenge in this area is that the position and timing of the switching are unknown.
26 Moreover, the switching happens over a very short time scale. Since the 1970s,
27 Nefedov and Ni [1–4] have conducted numerous studies on singularly perturbed
28 problems involving contrast structures. However, all of their studies assume that
29 the critical manifolds are normal hyperbolic manifolds of saddle type. This raises
30 an important issue: whether contrast structures exist when the critical manifold

[†]The corresponding author. Email: xiaovikdo@163.com(M. Ni).

¹School of Mathematical Sciences, East China Normal University, Shanghai, 200062, PR China.

²Shanghai Key Laboratory of Pure Mathematics and Mathematical Practice, Shanghai, 200062, PR China.

*The authors were supported by National Natural Science Foundation of China(NO.12371168) and in part by Science and Technology Commission of Shanghai Municipality(NO.18dz2271000).

Email: 843271281@qq.com(R. Li), xiaovikdo@163.com(M. Ni).

31 is non-normal hyperbolic, or where the degenerate equation has repeated roots.
 32 Butuzov [8] was one of the first to investigate such types of problems

$$\begin{cases} \varepsilon^2 \frac{d^2 u}{dx^2} = f(u, x, \varepsilon), & 0 < x < 1, \\ \frac{du}{dx}(0, \varepsilon) = 0, \quad \frac{du}{dx}(1, \varepsilon) = 0, \end{cases} \quad (1.1)$$

where

$$f(u, x, \varepsilon) = (u - \varphi_1(x))^2(u - \varphi_2(x))(u - \varphi_3(x)) - \varepsilon f_1(u, x, \varepsilon).$$

In [8], authors established a new method for studying singularly perturbed problems with repeated roots, which we call the non-standard boundary layer method. This approach successfully addressed the limitation of Vasil'eva's method [5], which cannot be applied to non-hyperbolic manifolds. As a result, problems involving multiple roots have become a major focus in the study of singularly perturbed problems. Butuzov [7, 9–11] not only conducted extensive research on double roots, but also expanded his work to include triple roots and elliptic problems [17–19]. Yang [14–16] studied piecewise-smooth systems based on equation (1.1), where

$$f(u, x, \varepsilon) = \begin{cases} f^{(-)}(u, x, \varepsilon), & 0 \leq x < x_0, \\ f^{(+)}(u, x, \varepsilon), & x_0 \leq x \leq 1. \end{cases}$$

33 Both $f^{(-)}(u, x, \varepsilon)$ and $f^{(+)}(u, x, \varepsilon)$ contain repeated roots. Yang [12, 13] had also
 34 extended the research to reaction-diffusion equation.

In this paper, we consider the problem (1.1), where

$$f(u, x, \varepsilon) = -(u - \varphi_1(x))^2(u - \varphi_2(x))(u - \varphi_3(x))^2 - \varepsilon f_1(u, x, \varepsilon).$$

35 In this paper, following Butuzov [8], we study a singularly perturbed boundary
 36 value problem where the degenerate equation has three distinct roots: two double
 37 roots and one simple root. The key difference between [8] and our study is that
 38 we focus on two double roots, whereas Butuzov studied a single root and a double
 39 root. The difficulty arises because the degenerate roots can “jump” from one non-
 40 hyperbolic manifold to another, making the existence of a smooth solution more
 41 uncertain.

42 This problem can be treated as two separate sub-problems, referred to as the
 43 left and right problems. We construct the formal asymptotic solution using the
 44 boundary layer method and solve it term by term. Finally, the solutions to the left
 45 and right problems are smoothly matched using the seaming method, leading to a
 46 smooth solution for the original problem.

47 We show that when the degenerate equation has repeated roots, i.e., when the
 48 critical manifolds are non-normal hyperbolic, there exists a contrast structure be-
 49 tween the roots. Unlike the case where the critical manifold is normal hyperbolic,
 50 not only is the formal asymptotic solution expanded in terms of fractional powers
 51 of the small parameter, but a non-standard method is also used to solve the in-
 52 ternal layer functions. These internal layer functions exhibit complex variations,
 53 transitioning from exponential decay to power-law decay.

2. Problem statement

We consider singularly perturbed problems with Neumann boundary conditions

$$\begin{cases} \varepsilon^2 \frac{d^2 u}{dx^2} = f(u, x, \varepsilon), & 0 < x < 1, \\ \frac{du}{dx}(0, \varepsilon) = 0, & \frac{du}{dx}(1, \varepsilon) = 0, \end{cases} \quad (2.1)$$

$$(2.2)$$

where $\varepsilon > 0$ is a small parameter. Here $f(u, x, \varepsilon)$ has the form

$$f(u, x, \varepsilon) = -(u - \varphi_1(x))^2(u - \varphi_2(x))(u - \varphi_3(x))^2 - \varepsilon f_1(u, x, \varepsilon).$$

We will call a multivariable function smooth if it is infinitely differentiable with respect to all arguments.

Assume that the following conditions are hold.

A1. The functions $\varphi_i(x), i = 1, 2, 3$, are smooth and satisfy

$$\varphi_1(x) < \varphi_2(x) < \varphi_3(x)$$

for $0 \leq x \leq 1$.

Condition A1 guarantees that the roots of the degenerate equation are distinct.

A2. The function $f_1(u, x, \varepsilon)$ is smooth, not identically equal to zero for $0 \leq x \leq 1$, and satisfies

$$\begin{aligned} \bar{f}_1^{(-)}(x) &:= f_1(\varphi_1(x), x, 0) > 0, \\ \bar{f}_1^{(+)}(x) &:= f_1(\varphi_3(x), x, 0) < 0. \end{aligned}$$

A3. The equation

$$I(\bar{x}_0) := \int_{\varphi_1(x_0)}^{\varphi_3(x_0)} f(u, x_0, 0) du = 0,$$

has the root $x_0 \in (0, 1)$, and $I'(x_0) \neq 0$. The root x_0 is called the transition point.

The boundary layer method is commonly used to solve such singularly perturbed problems. The general process involves expanding the solution as a power series form with respect to small parameters ε , and then finding each coefficient iteratively. This process is known as constructing asymptotic solutions.

3. Construction of asymptotic solutions

To determine the steptype asymptotic solution of the problem (2.1)-(2.2), we treat the original problem as two separate problems, namely, the left and right problems. The solutions of the left and right problems are then smoothly matched using the seaming method, yielding a smooth solution of the original problem. The asymptotic for function $u(x, \varepsilon)$ has the form

$$u(x, \varepsilon) = \begin{cases} u^{(-)}(x, \varepsilon), & 0 \leq x < x_*, \\ u^{(+)}(x, \varepsilon), & x_* \leq x \leq 1. \end{cases} \quad (3.1)$$

72 The left and right problems can be written as follows.

73 Left problem is defined for $0 \leq x \leq x_*$ as

$$\begin{cases} \varepsilon^2 \frac{d^2 u^{(-)}}{dx^2} = f(u^{(-)}, x, \varepsilon), \\ \frac{du^{(-)}}{dx}(0, \varepsilon) = 0, \quad u^{(-)}(x_*, \varepsilon) = \varphi_2(x_*). \end{cases} \quad (3.2)$$

74 Right problem is defined for $x_* \leq x \leq 1$ as

$$\begin{cases} \varepsilon^2 \frac{d^2 u^{(+)}}{dx^2} = f(u^{(+)}, x, \varepsilon), \\ u^{(+)}(x_*, \varepsilon) = \varphi_2(x_*), \quad \frac{du^{(-)}}{dx}(1, \varepsilon) = 0, \end{cases} \quad (3.3)$$

75 where

$$u^{(-)}(x, \varepsilon) = \bar{u}^{(-)}(x, \varepsilon) + Q^{(-)}(\tau, \varepsilon) + \Pi^{(-)}(\xi, \varepsilon), \quad (3.4)$$

$$u^{(+)}(x, \varepsilon) = \bar{u}^{(+)}(x, \varepsilon) + Q^{(+)}(\tau, \varepsilon) + \Pi^{(+)}(\tilde{\xi}, \varepsilon), \quad (3.5)$$

77 Here functions $\bar{u}^{(\pm)}(x, \varepsilon)$ are the regular parts of asymptotic, $Q^{(\pm)}(\tau, \varepsilon)$ are the
78 asymptotic inner layer, and $\Pi^{(-)}(\xi, \varepsilon), \Pi^{(+)}(\tilde{\xi}, \varepsilon)$ are the boundary layer functions.
79 We represent these functions in power series expansion as follows:

$$\bar{u}^{(\pm)}(x, \varepsilon) = \bar{u}_0^{(\pm)}(x) + \varepsilon^{\frac{1}{2}} \bar{u}_1^{(\pm)}(x) + \cdots + \varepsilon^{\frac{i}{2}} \bar{u}_i^{(\pm)}(x) + \cdots, \quad (3.6)$$

$$Q^{(\pm)}(\tau, \varepsilon) = Q_0^{(\pm)}(\tau) + \varepsilon^{\frac{1}{4}} Q_1^{(\pm)}(\tau) + \cdots + \varepsilon^{\frac{i}{4}} Q_i^{(\pm)}(\tau) + \cdots, \quad \tau = \frac{x - x_*}{\varepsilon}, \quad (3.7)$$

$$\Pi^{(-)}(\xi, \varepsilon) = \varepsilon^{\frac{3}{4}} (\Pi_0^{(-)}(\xi) + \varepsilon^{\frac{1}{4}} \Pi_1^{(-)}(\xi) + \cdots + \varepsilon^{\frac{i}{4}} \Pi_i^{(-)}(\xi) + \cdots), \quad \xi = \frac{x}{\varepsilon^{3/4}}, \quad (3.8)$$

$$\Pi^{(+)}(\tilde{\xi}, \varepsilon) = \varepsilon^{\frac{3}{4}} (\Pi_0^{(+)}(\tilde{\xi}) + \varepsilon^{\frac{1}{4}} \Pi_1^{(+)}(\tilde{\xi}) + \cdots + \varepsilon^{\frac{i}{4}} \Pi_i^{(+)}(\tilde{\xi}) + \cdots), \quad \tilde{\xi} = \frac{1 - x}{\varepsilon^{3/4}}. \quad (3.9)$$

83 Functions $Q^{(\pm)}(\tau, \varepsilon)$, $\Pi^{(-)}(\xi, \varepsilon)$ and $\Pi^{(+)}(\tilde{\xi}, \varepsilon)$ are expanded in terms of $\varepsilon^{1/4}$,
84 because the degenerate equation contains repeated roots. Then, the refined algo-
85 rithms for accurate searching coefficients must be used. For $\bar{u}^{(\pm)}(x, \varepsilon)$ the order
86 $\varepsilon^{1/2}$ is enough.

87 The denominators in series expansions for ξ , $\tilde{\xi}$ and τ are different due to the
88 difference in boundary conditions. For inner layer we have only Neumann boundary
89 conditions. We multiply the power series in equations (3.8),(3.9) by $\varepsilon^{\frac{3}{4}}$ to balance
90 the boundary conditions due to specific choice of ξ . By x_* we denote the transition
91 point, which can be also expressed in the form of power series:

$$x_* = x_0 + \varepsilon^{\frac{1}{4}} x_1 + \cdots + \varepsilon^{\frac{i}{4}} x_i + \cdots. \quad (3.10)$$

92 Then, we find the coefficients of asymptotic expansions (3.6)-(3.9) for right and
93 left problems. We also prove that there exists $x_* \in (0, 1)$ where

$$\frac{du^{(-)}}{dx}(x_*, \varepsilon) = \frac{du^{(+)}}{dx}(x_*, \varepsilon). \quad (3.11)$$

94 The value x_* is used by seaming method for matching solutions of left and right
95 problems.

96 Finally, we prove that the solution of problem (2.1) - (2.2) is also the solution
97 of (3.2) - (3.3) with contrast structure near $x = x_*$.

98 4. The asymptotic of the right and left problems 99 solution

Let us consider the left problem (3.2). Substituting series (3.4) to (3.2), we get

$$\begin{cases} \varepsilon^2 \frac{d^2 \bar{u}^{(-)}}{dx^2} + \frac{d^2 Q^{(-)}}{d\tau^2} + \varepsilon^{\frac{1}{2}} \frac{d^2 \Pi^{(-)}}{d\xi^2} = \bar{f}^{(-)} + Q^{(-)} f + \Pi^{(-)} f, \\ \frac{d\bar{u}^{(-)}}{dx}(0, \varepsilon) + \varepsilon^{\frac{4}{3}} \frac{d\Pi^{(-)}}{d\xi}(0, \varepsilon) = 0, \quad \bar{u}^{(-)}(x_*, \varepsilon) + Q^{(-)}(0, \varepsilon) = \varphi_2(x_*), \end{cases}$$

where

$$\begin{aligned} \bar{f}^{(-)} &= f(\bar{u}^{(-)}(x, \varepsilon), x, \varepsilon), \\ \Pi^{(-)} f &= f(\bar{u}^{(-)}(\xi \varepsilon^{3/4}, \varepsilon) + \Pi^{(-)}(\tau, \varepsilon), \xi \varepsilon^{3/4}, \varepsilon) - f(\bar{u}^{(-)}(\xi \varepsilon^{3/4}, \varepsilon), \xi \varepsilon^{3/4}, \varepsilon), \\ Q^{(-)} f &= f(\bar{u}^{(-)}(x_* + \tau \varepsilon, \varepsilon) + Q^{(-)}(\tau, \varepsilon), x_* + \tau \varepsilon, \varepsilon) \\ &\quad - f(\bar{u}^{(-)}(x_* + \tau \varepsilon, \varepsilon), x_* + \tau \varepsilon, \varepsilon). \end{aligned}$$

The regular part of asymptotics can be found from the equation

$$\varepsilon^2 \frac{d^2 \bar{u}^{(-)}}{dx^2} = \bar{f}^{(-)}, \quad 0 < x < x_*,$$

the inner layer from

$$\frac{d^2 Q^{(-)}}{d\tau^2} = Q^{(-)} f, \quad \tau < 0,$$

and boundary layer from

$$\varepsilon^{\frac{1}{2}} \frac{d^2 \Pi^{(-)}}{d\xi^2} = \Pi^{(-)} f, \quad \xi > 0.$$

100 The right problem (3.3) is solved similarly.

101 4.1. The regular part of asymptotics

102 Let us consider the regular part $u^{(-)}(x, \varepsilon)$ of asymptotics for left problem solution.

103 It satisfies

$$\varepsilon^2 \frac{d^2 \bar{u}^{(-)}}{dx^2} = f(\bar{u}^{(-)}, x, \varepsilon). \quad (4.1)$$

Substituting series (3.6) to equation (4.1), we get

$$\begin{aligned}
& \varepsilon^2 \frac{d^2 \bar{u}_0^{(-)}(x)}{dx^2} + \varepsilon^{\frac{5}{2}} \frac{d^2 \bar{u}_1^{(-)}(x)}{dx^2} + \dots + \varepsilon^{\frac{k+4}{2}} \frac{d^2 \bar{u}_k^{(-)}(x)}{dx^2} + \dots = \\
& = -(\bar{u}^{(-)} - \varphi_1(x))^2 (\bar{u}^{(-)} - \varphi_2(x)) (\bar{u}^{(-)} - \varphi_3(x))^2 - \varepsilon f_1(\bar{u}^{(-)}, x, \varepsilon) \\
& = -(\bar{u}_0^{(-)} - \varphi_1(x))^2 (\bar{u}_0^{(-)} - \varphi_2(x)) (\bar{u}_0^{(-)} - \varphi_3(x))^2 + \\
& \quad + \varepsilon^{\frac{1}{2}} [-(\bar{u}_0^{(-)} - \varphi_1(x))^2 \bar{u}_1^{(-)} (\bar{u}_0^{(-)} - \varphi_3(x))^2] \\
& \quad + \varepsilon [-(\bar{u}_1^{(-)})^2 (\bar{u}_0^{(-)} - \varphi_2(x)) (\bar{u}_0^{(-)} - \varphi_3(x))^2 - (\bar{u}_0^{(-)} - \varphi_1(x))^2 \bar{u}_2^{(-)} (\bar{u}_0^{(-)} - \varphi_3(x))^2 \\
& \quad - (\bar{u}_0^{(-)} - \varphi_1(x))^2 (\bar{u}_0^{(-)} - \varphi_2(x)) (\bar{u}_1^{(-)})^2 - f_1(\bar{u}_0^{(-)}, x, 0)] + \dots \\
& \quad + \varepsilon^{\frac{k}{2}} \sum_{\substack{2i+2j+2h+2l \leq k \\ i > 0, j, h, l \geq 0}} [-\bar{u}_i^{(-)} \bar{u}_l^{(-)} (\bar{u}_{k-2i-2j}^{(-)} - \tilde{\varphi}_2) (\bar{u}_j^{(-)} - \tilde{\varphi}_{3j}) (\bar{u}_h^{(-)} - \tilde{\varphi}_{3h}) - \bar{f}_k^{(-)}(x)] + \\
& \quad + \dots
\end{aligned}$$

where

$$\begin{aligned}
\tilde{\varphi}_{3j} &= \begin{cases} \varphi_3(x), & j = 0, \\ 0, & j \neq 0, \end{cases} & \tilde{\varphi}_{3h} &= \begin{cases} \varphi_3(x), & h = 0, \\ 0, & h \neq 0, \end{cases} \\
\tilde{\varphi}_2 &= \begin{cases} \varphi_2(x), & k - 2i - 2j = 0, \\ 0, & k - 2i - 2j \neq 0, \end{cases}
\end{aligned}$$

By equating the coefficients of the same powers of ε on both sides of the equation, for $\bar{u}_0^{(-)}(x)$ we obtain

$$-(\bar{u}_0^{(-)} - \varphi_1(x))^2 (\bar{u}_0^{(-)} - \varphi_2(x)) (\bar{u}_0^{(-)} - \varphi_3(x))^2 = 0.$$

For solution we can take $\varphi_1(x)$, i.e

$$\bar{u}_0^{(-)} = \varphi_1(x).$$

104 Then $\bar{u}_1^{(-)}$ is determined from the equation

$$\bar{h}^{(-)}(x) (\bar{u}_1^{(-)}(x))^2 - \bar{f}_1^{(-)}(x) = 0, \quad (4.2)$$

105 where

$$\begin{aligned}
\bar{h}^{(-)}(x) &= -(\varphi_1(x) - \varphi_3(x))^2 (\varphi_1(x) - \varphi_2(x)), \\
\bar{f}_1^{(-)}(x) &= f_1(\varphi_1(x), x, 0).
\end{aligned} \quad (4.3)$$

106 According to condition A2, the solution of equation (4.2) must exist. Then, we
107 can take a positive root of

$$\bar{u}_1^{(-)} = \sqrt{\frac{\bar{f}_1^{(-)}(x)}{\bar{h}^{(-)}(x)}} \quad (4.4)$$

108 as solution for (4.2).

109 Similarly, the higher order terms $\bar{u}_k^{(-)}(x)$ of series (3.6) can be obtained from
110 algebraic equations

$$[2\bar{h}^{(-)}(x) \bar{u}_1^{(-)}(x)] \bar{u}_k^{(-)}(x) - \bar{f}_k^{(-)}(x) = 0, \quad k > 1, \quad (4.5)$$

111 where $\bar{f}_k^{(-)}(x)$ is the known function that depends on $\bar{u}_j^{(-)}(x)$, ($j < k$). Hence, the
112 solution of (4.4)-(4.5) is unique and equal to $\bar{u}_k^{(-)}(x)$.

113 4.2. The inner layer of asymptotics

Now let us consider the inner layer term, namely, $Q^{(-)}(\tau, \varepsilon)$. Recall that it satisfies the following equation and boundary conditions

$$\begin{cases} \frac{d^2 Q^{(-)}(\tau, \varepsilon)}{d\tau^2} = Q^{(-)} f := f(\bar{u}^{(-)}(x_* + \varepsilon\tau, \varepsilon) + Q^{(-)}(\tau, \varepsilon), x_* + \varepsilon\tau, \varepsilon) \\ \quad - f(\bar{u}^{(-)}(x_* + \varepsilon\tau, \varepsilon), x_* + \varepsilon\tau, \varepsilon), \\ Q^{(-)}(0, \varepsilon) = \varphi_2(x_*) - \bar{u}^{(-)}(x_*, \varepsilon). \quad Q^{(-)}(-\infty, \varepsilon) = 0. \end{cases} \quad (4.6)$$

$$(4.7)$$

To find the coefficients of power series expansion for $Q^{(-)}(\tau, \varepsilon)$, we follow the Butuzov's nonstandard method from [11]. For convenience, for $Q_i^{(-)}$ we mention only dependance on τ . We have

$$h^{(-)}(u, x) = -(u - \varphi_2(x))(u - \varphi_3(x))^2, \quad \varphi_1(x) = \varphi^{(-)}(x).$$

Substituting series (3.7) into (4.6), we obtain the the following equation for $Q_i^{(-)}(\tau)$:

$$\begin{aligned} & \frac{d^2 Q_0^{(-)}(\tau)}{d\tau^2} + \varepsilon^{\frac{1}{4}} \frac{d^2 Q_1^{(-)}(\tau)}{d\tau^2} + \dots + \varepsilon^{\frac{k}{4}} \frac{d^2 Q_k^{(-)}(\tau)}{d\tau^2} + \dots \\ &= -(\bar{u}^{(-)} + Q^{(-)} - \varphi_1)^2 (\bar{u}^{(-)} + Q^{(-)} - \varphi_2) (\bar{u}^{(-)} + Q^{(-)} - \varphi_3)^2 - \varepsilon f_1(\bar{u}^{(-)} + Q^{(-)}, x, \varepsilon) - \\ & \quad - [-(\bar{u}^{(-)} - \varphi_1)^2 (\bar{u}^{(-)} - \varphi_2) (\bar{u}^{(-)} - \varphi_3)^2 - \varepsilon f_1(\bar{u}^{(-)}, x, \varepsilon)] \\ &= -(\bar{u}_0^{(-)} + Q_0^{(-)} - \varphi_2) (\bar{u}_0^{(-)} + Q_0^{(-)} - \varphi_3)^2 (Q_0^{(-)})^2 + \\ & \quad + \varepsilon^{\frac{1}{4}} \{ -2Q_1^{(-)} Q_0^{(-)} (\bar{u}_0^{(-)} + Q_0^{(-)} - \varphi_2) (\bar{u}_0^{(-)} + Q_0^{(-)} - \varphi_3)^2 \\ & \quad - (Q_0^{(-)})^2 [Q_1^{(-)} (\bar{u}_0^{(-)} + Q_0^{(-)} - \varphi_3)^2 + 2Q_1^{(-)} (\bar{u}_0^{(-)} + Q_0^{(-)} - \varphi_2) (\bar{u}_0^{(-)} + Q_0^{(-)} - \varphi_3)] \} + \dots \\ & \quad + \varepsilon^{\frac{k}{2}} \left\{ \sum_{i,j,l,h \geq 0}^{2i+2j+2h+2l \leq k} -[(\bar{u}_{i/2}^{(-)} + Q_i^{(-)} - \tilde{\varphi}_1) (\bar{u}_{l/2}^{(-)} + Q_l^{(-)} - \tilde{\varphi}_1) \times \right. \\ & \quad \times (\bar{u}_{k-2i-2j-2l-2h/2}^{(-)} + Q_{k-2i-2j-2l-2h}^{(-)} - \tilde{\varphi}_2) (\bar{u}_{j/2}^{(-)} + Q_j^{(-)} - \tilde{\varphi}_3) (\bar{u}_{h/2}^{(-)} + Q_h^{(-)} - \tilde{\varphi}_3)] - \\ & \quad \left. - \sum_{i>0, j, h, l \geq 0}^{2i+2j+2h+2l \leq k} -[\bar{u}_i^{(-)} \bar{u}_l^{(-)} (\bar{u}_{k-2i-2j}^{(-)} - \tilde{\varphi}_2) (\bar{u}_j^{(-)} - \tilde{\varphi}_{3j}) (\bar{u}_h^{(-)} - \tilde{\varphi}_{3h})] - Q f_k \right\} \\ & \quad + \dots \end{aligned}$$

where

$$\tilde{\varphi}_1 = \begin{cases} \varphi_1(x_* + \varepsilon\tau), & i = 0, \\ 0, & i \neq 0, \end{cases} \quad \bar{u}_\alpha^{(-)} = \begin{cases} \bar{u}_\alpha^{(-)}, & \text{if } \alpha \text{ is odd,} \\ 0, & \text{if } \alpha \text{ is even.} \end{cases}$$

According to Butuzov's nonstandard method, we get the following equation and boundary conditions that determine $(Q_0^{(-)}(\tau))$:

$$\begin{cases} \frac{d^2 Q_0^{(-)}}{d\tau^2} = h^{(-)}(\varphi^{(-)}(x_*) + Q_0^{(-)}, x_*) [(Q_0^{(-)})^2 + 2\sqrt{\varepsilon} \bar{u}_1^{(-)}(x_*) Q_0^{(-)}], & (4.8) \\ Q_0^{(-)}(0) = \varphi_2(x_*) - \varphi^{(-)}(x_*), \quad Q_0^{(-)}(-\infty) = 0. & (4.9) \end{cases}$$

This problem can be reduced to the first-order boundary value problem

$$\begin{cases} \frac{dQ_0^{(-)}}{d\tau} = \left[2 \int_0^{Q_0^{(-)}} h^{(-)}(\varphi^{(-)}(x_*) + s, x_*) (s^2 + 2\sqrt{\varepsilon}\bar{u}_1^{(-)}(x_*)s) ds \right]^{\frac{1}{2}}, & (4.10) \\ Q_0^{(-)}(0) = \varphi_2(x_*) - \varphi^{(-)}(x_*). & (4.11) \end{cases}$$

114 It is challenging to find the solution to the above boundary value problem di-
115 rectly. However, it can be estimated using a differential inequality. Since the
116 function $h^{(-)}(\varphi^{(-)}(x_*) + s, x_*)$ is bounded for $0 \leq s \leq Q_0^{(-)}$, let α_1 denote its
117 minimum value and α_2 its maximum value. Thus, the solution $Q_0^{(-)}(\tau)$ admits the
118 following estimate

$$Q_{\alpha_2}^{(-)}(\tau) \leq Q_0^{(-)}(\tau) \leq Q_{\alpha_1}^{(-)}(\tau). \quad (4.12)$$

119 where

$$Q_{\alpha}^{(-)}(\tau) = \frac{12\sqrt{\varepsilon}\bar{u}_1^{(-)}(x_*)(1 + O(\varepsilon^{1/4}))e^{\varepsilon^{1/4}\alpha k_0\tau}}{\left\{ 1 - \left[1 - \left(\frac{12\bar{u}_1^{(-)}(x_*)}{\varphi_2(x_*) - \varphi^{(-)}(x_*)} \right)^{1/2} \varepsilon^{1/4} + O(\varepsilon^{1/2}) \right] e^{\varepsilon^{1/4}\alpha k_0\tau} \right\}^2}. \quad (4.13)$$

120 Here we denote $k_0 = [2\bar{u}_1^{(-)}(x_*)]^{\frac{1}{2}} > 0$.

121 According to the different decay behaviors of $Q_0^{(-)}(\tau)$, the inner layer can be
122 divided into three regions. Here, we discuss only the left problem, i.e., when $\tau < 0$.
123 Specifically:

- 124 1. If $-\frac{1}{\varepsilon^\gamma} \leq \tau \leq 0$, where $0 \leq \gamma \leq \frac{1}{4}$, then $Q_0^{(-)}(\tau)$ decays according to a power
125 law as $\tau \rightarrow \infty$.
- 126 2. If $-\frac{1}{\varepsilon^{1/4}} \leq \tau \leq -\frac{1}{\varepsilon^\gamma}$, the decay of $Q_0^{(-)}(\tau)$ changes from a power-law decay to
127 an exponential decay.
- 128 3. If $\tau \leq -\frac{1}{\varepsilon^{1/4}}$, then $Q_0^{(-)}(\tau)$ exhibits exponential decay with respect to the new
129 variable $\theta = \frac{x}{\varepsilon^{3/4}}$.

130 There are also three analogous regions when $\tau > 0$.

131 For higher order coefficients of $Q_k^{(-)}(\tau)$ we have boundary value problem

$$\frac{d^2 Q_k^{(-)}}{d\tau^2} = \beta^{(-)}(\tau, \varepsilon) Q_k^{(-)} + q_k^{(-)}(\tau, \varepsilon), \quad (4.14)$$

132

$$Q_k^{(-)}(0) = \begin{cases} -\bar{u}_{\frac{k}{2}}^{(-)}(x_*), & \text{if } k \text{ is odd,} \\ 0, & \text{if } k \text{ is even,} \end{cases} \quad Q_k^{(-)}(\pm\infty) = 0, \quad (4.15)$$

133 where

$$\begin{aligned} \beta^{(-)}(\tau, \varepsilon) &= h_u^{(-)}(\tau) [(Q_0^{(-)}(\tau))^2 + 2\sqrt{\varepsilon}\bar{u}_1^{(-)}(x_*)Q_0^{(-)}(\tau)] + \\ &\quad + 2h^{(-)}(\tau) [Q_0^{(-)}(\tau) + \sqrt{\varepsilon}\bar{u}_1^{(-)}(x_*)], \\ h_u^{(-)}(\tau) &= \frac{\partial h}{\partial u}(\varphi^{(-)}(x_*) + Q_0^{(-)}(\tau), x_*, 0), \\ h^{(-)}(\tau) &= h^{(-)}(\varphi^{(-)}(x_*) + Q_0^{(-)}(\tau), x_*, 0), \end{aligned} \quad (4.16)$$

139 By equating the coefficients of the same powers of ε on both sides of the equation,
 140 for $\Pi_k^{(-)}$, $k = 0, 1, 2, \dots$ we get

$$\frac{d^2 \Pi_k^{(-)}}{d\xi^2} = 2\bar{h}^{(-)}(0)\bar{u}_1^{(-)}(0)\Pi_k^{(-)} + \pi_k^{(-)}(\xi), \quad \xi > 0, \quad k = 0, 1, \dots \quad (4.22)$$

141

$$\frac{d\Pi_k^{(-)}}{d\xi}(0) = \begin{cases} -\frac{d\bar{u}_{k/2}^{(-)}}{dx}(0), & \text{if } k \text{ is odd,} \\ 0, & \text{if } k \text{ is even,} \end{cases} \quad \Pi_k^{(-)}(+\infty) = 0, \quad (4.23)$$

142 where $\pi_k^{(-)}$ depends on known functions $\Pi_j^{(-)}(\xi)$, $j < k$. In particular, $\pi_0^{(-)} \equiv 0$.
 143 Since the equations (4.22) are linear, the solution $\Pi_0^{(-)}(\xi)$ can be written as

$$\Pi_0^{(-)}(\xi) = \varphi^{(-)'}(0)[2\bar{h}^{(-)}(0)\bar{u}_1^{(-)}(0)]^{-\frac{1}{2}} \exp\left(- (2\bar{h}^{(-)}(0)\bar{u}_1^{(-)}(0))^{\frac{1}{2}} \xi\right). \quad (4.24)$$

144

We also have the estimate:

$$\|\Pi_k^{(-)}(\xi)\| \leq ce^{-\kappa\xi}, \quad \xi > 0, \quad k = 0, 1, \dots \quad (4.25)$$

145

146

147

Currently, each term of the asymptotic expansion for the left problem has been determined and satisfies the exponential decay estimate. The right problem can be analyzed in the same way, so we skip the details here.

148

4.4. Proof of existence of solution to the problem (2.1)-(2.2)

149

150

In this section, we will prove the existence of a smooth solution to the original problem (2.1)-(2.2) using the sewing method.

Theorem 4.1. *Suppose we have the boundary value problem (2.1)-(2.2) and the conditions (A1)-(A3) are met. Then for sufficiently small $\varepsilon > 0$, there exist a smooth solution $u(x, \varepsilon)$ in the asymptotic form*

$$u(x, \varepsilon) = \begin{cases} U_n^{(-)}(x, \varepsilon) + O(\varepsilon^{\frac{n+1}{2}}), & 0 \leq x < (x_*)_{2n+5}, \\ U_n^{(+)}(x, \varepsilon) + O(\varepsilon^{\frac{n+1}{2}}), & (x_*)_{2n+5} \leq x \leq 1. \end{cases}$$

151

152

where $n \in \mathbb{N}$, $(x_*)_{2n+5} = \sum_{k=0}^{2n+5} \varepsilon^{\frac{k}{4}} \bar{x}_k$, the functions $U_n^{(\pm)}(x, \varepsilon)$ can be found from (4.27) with $\tau = [(x - x_*)_{2n+5}]/\varepsilon$.

153

To prove this theorem, we first introduce two lemmas.

154

155

156

Lemma 4.1. *If (A1) and (A2) are satisfied, then for sufficiently small $\varepsilon > 0$, the solution $u^{(-)}(x, \varepsilon)$ of the left problem (3.2) and the solution $u^{(+)}(x, \varepsilon)$ of the right problem (3.3) are respectively:*

$$\begin{aligned} u^{(-)}(x, \varepsilon) &= U_n^{(-)}(x, \varepsilon) + O(\varepsilon^{\frac{n+1}{2}}), \quad 0 \leq x \leq x_*, \\ u^{(+)}(x, \varepsilon) &= U_n^{(+)}(x, \varepsilon) + O(\varepsilon^{\frac{n+1}{2}}), \quad x_* \leq x \leq 1. \end{aligned} \quad (4.26)$$

157

where

$$\begin{aligned} U_n^{(-)}(x, \varepsilon) &= \sum_{k=0}^n \varepsilon^{\frac{k}{2}} \bar{u}_k^{(-)}(x) + \sum_{k=0}^{2n+1} \varepsilon^{\frac{k}{4}} Q_k^{(-)}(\tau) + \varepsilon^{\frac{3}{4}} \sum_{k=0}^{2n} \varepsilon^{\frac{k}{4}} \Pi_k^{(-)}(\xi), \\ U_n^{(+)}(x, \varepsilon) &= \sum_{k=0}^n \varepsilon^{\frac{k}{2}} \bar{u}_k^{(+)}(x) + \sum_{k=0}^{2n+1} \varepsilon^{\frac{k}{4}} Q_k^{(+)}(\tau) + \varepsilon^{\frac{3}{4}} \sum_{k=0}^{2n} \varepsilon^{\frac{k}{4}} \Pi_k^{(+)}(\tilde{\xi}). \end{aligned} \quad (4.27)$$

158 The proof is analogous to the one in [8] and skipped here.

159 **Lemma 4.2.** For the derivative $\frac{du^{(-)}}{dx}(x, \varepsilon)$, $\frac{du^{(+)}}{dx}(x, \varepsilon)$, the asymptotic repre-
160 sentations

$$\begin{aligned} \frac{du^{(-)}}{dx}(x, \varepsilon) &= \frac{dU_n^{(-)}}{dx}(x, \varepsilon) + O(\varepsilon^{\frac{n-1}{2}}), & 0 \leq x < x_*, \\ \frac{du^{(+)}}{dx}(x, \varepsilon) &= \frac{dU_n^{(+)}}{dx}(x, \varepsilon) + O(\varepsilon^{\frac{n-1}{2}}), & x_* \leq x \leq 1. \end{aligned} \quad (4.28)$$

161 are true.

162 The proof is also analogous to the one in [8].

163 Now, we prove the original theorem.

164 We rewrite the smooth seaming condition (3.11) as

$$I(x_*, \varepsilon) = \varepsilon \frac{du^{(-)}}{dx}(x_*, \varepsilon) - \varepsilon \frac{du^{(+)}}{dx}(x_*, \varepsilon) = 0. \quad (4.29)$$

Substituting expansion (3.10) for x_* in at $\tau = 0$, we get

$$\frac{dQ_0^{(-)}}{d\tau}(0) = I^{(-)}(x_0) + \sum_{i=1}^{+\infty} \varepsilon^{\frac{i}{4}} g_i^{(-)}(x),$$

where

$$I^{(-)}(x_0) = \left[2 \int_{\varphi^{(-)}(x_0)}^{\varphi_2(x_0)} f(u, x_0, 0) du \right]^{\frac{1}{2}},$$

and $g_i(x)$ is sufficiently smooth functions that depend on $x_k, k \leq i$. Similarly for the right problem solution, we have

$$\frac{dQ_0^{(+)}}{d\tau}(0) = I^{(+)}(x_0) + \sum_{i=1}^{+\infty} \varepsilon^{\frac{i}{4}} g_i^{(+)}(x),$$

where

$$I^{(+)}(x_0) = \left[2 \int_{\varphi^{(+)}(x_0)}^{\varphi_2(x_0)} f(u, x_0, 0) du \right]^{\frac{1}{2}}.$$

165 Substituting power series expansion of (3.4) and (3.5) with respect to $\varepsilon^{\frac{1}{4}}$ to
166 (4.29), we obtain

$$\begin{aligned} I(x_*, \varepsilon) &= \varepsilon \left(\frac{d\varphi^{(-)}}{dx}(x_*) + \varepsilon^{\frac{1}{2}} \frac{d\bar{u}_1^{(-)}}{dx}(x_*) + \dots \right) + \left(\frac{dQ_0^{(-)}}{d\tau}(0) + \varepsilon^{\frac{1}{4}} \frac{dQ_1^{(-)}}{d\tau}(0) + \dots \right) \\ &\quad - \varepsilon \left(\frac{d\varphi^{(+)}}{dx}(x_*) + \varepsilon^{\frac{1}{2}} \frac{d\bar{u}_1^{(+)}}{dx}(x_*) + \dots \right) - \left(\frac{dQ_0^{(+)}}{d\tau}(0) + \varepsilon^{\frac{1}{4}} \frac{dQ_1^{(+)}}{d\tau}(0) + \dots \right) = \\ &= \left(\frac{dQ_0^{(-)}}{d\tau}(0) - \frac{dQ_0^{(+)}}{d\tau}(0) \right) + \varepsilon^{\frac{1}{4}} \left(\frac{dQ_1^{(-)}}{d\tau}(0) - \frac{dQ_1^{(+)}}{d\tau}(0) \right) + \dots \\ &= [I^{(-)}(x_0) - I^{(+)}(x_0) + \varepsilon^{\frac{1}{4}} (I^{(-)'}(x_0)x_1 - I^{(+)'}(x_0)x_1) + \dots] + \dots \\ &= H(x_0) + \varepsilon^{\frac{1}{4}} [H'(x_0)x_1 + m_1] + \dots + \varepsilon^{\frac{k}{4}} [H'(x_0)x_k + m_k] + \dots = 0. \end{aligned} \quad (4.30)$$

167 Here $H(x) = I^{(-)}(x) - I^{(+)}(x)$. Then, according to (A3), $x = x_0$ is the root of the
 168 equation $H(x) = 0$.

169 Since $H'(x_0) \neq 0$, the higher order coefficients $x_k, k \geq 1$ can be uniquely
 170 determined from the following linear algebraic equations

$$H'(x_0)x_k + m_k = 0, \quad k \geq 1, \quad (4.31)$$

171 where m_k depend on the known numbers $x_{j, j < k}$. Note that

$$m_1 = [H(x_0)^{-1}(J^{(+)} - J^{(-)})]. \quad (4.32)$$

172 Since $q_1^{(-)} = 0$, we have $m_1 = 0$ and then $x_1 = 0$.

173 To prove the existence of a solution with contrast structures for the problem
 174 (2.1)-(2.2), we reconsider the left (3.2) and right (3.3) problems and modify x_* to
 175 the following form:

$$x_* = x_\delta^{(\pm)} := \bar{x}_0 + \varepsilon^{\frac{2}{4}}\bar{x}_1 + \cdots + \varepsilon^{\frac{2m+1}{4}}(\bar{x}_{m+1} \pm \delta), \quad (4.33)$$

176 where δ is an arbitrary real number, and δ is bounded as $\varepsilon \rightarrow 0$. Lemma 1 shows
 177 that the solutions of the left and right problems exist and have uniform asymptotic
 178 expansions. It only requires replacing the variable τ with $\tau = \frac{x - x_\delta^{(\pm)}}{\varepsilon}$, and the
 179 estimate in (4.19) still holds.

180 Rewriting (4.29) and taking $n = \left\lfloor \frac{2m+1}{4} \right\rfloor$ in (4.28), we have

$$\begin{aligned} I(x_\delta^{(\pm)}, \varepsilon) &= \varepsilon \frac{du^{(-)}}{dx}(x_\delta^{(\pm)}, \varepsilon, \delta) - \varepsilon \frac{du^{(+)}}{dx}(x_\delta^{(\pm)}, \varepsilon, \delta) \\ &= H(\bar{x}_0) + \sum_{k=2}^{2m} \varepsilon^{\frac{k}{4}} [H'(\bar{x}_0)\bar{x}_k + m_k] + \varepsilon^{\frac{2m+1}{4}} H'(\bar{x}_0)(\pm\delta) + O(\varepsilon^{\frac{2m+2}{4}}). \end{aligned} \quad (4.34)$$

181 where x_0 and x_k can be determined by (A2) and (4.31). The last two terms on the
 182 right-hand side of (4.34) depend on δ and are uniformly small as $\delta \rightarrow 0$. Thus,

$$I(x_\delta^{(\pm)}, \varepsilon) = \varepsilon^{\frac{2m+1}{4}} (H'(\bar{x}_0)(\pm\delta) + O(\varepsilon^{\frac{1}{4}})). \quad (4.35)$$

Since $H'(\bar{x}_0) \neq 0$, the sign of (4.35) depends on δ . Therefore, there exists a δ , such
 that

$$(H'(\bar{x}_0)\delta + O(\varepsilon^{\frac{1}{4}}))(-H'(\bar{x}_0)\delta + O(\varepsilon^{\frac{1}{4}})) < 0.$$

According to the intermediate value theorem, there exists $\delta = \bar{\delta}(\varepsilon) = O(\varepsilon^{1/4})$ such
 that (4.35) is equal to 0 when ε is sufficiently small. So, we have

$$\varepsilon \frac{du^{(-)}}{dx}(x_{\bar{\delta}}, \varepsilon, \delta) - \varepsilon \frac{du^{(+)}}{dx}(x_{\bar{\delta}}, \varepsilon, \delta) = 0.$$

It means that the function

$$u(x, \varepsilon) = \begin{cases} u^{(-)}(x, \varepsilon), & 0 \leq x < x_*, \\ u^{(+)}(x, \varepsilon), & x_* \leq x \leq 1, \end{cases}$$

183 is the solution of (2.1)-(2.2) with contrast structure near x_* . This completes the
 184 proof of the theorem.

185 5. Example

186 In this section we consider the following boundary value problem:

$$\begin{cases} \varepsilon^2 \frac{d^2 u}{dx^2} = -(u-x)^2(u-\frac{6}{5})(u-2)^2 - \varepsilon(\frac{6}{5}-u), & 0 < x < 1, \\ \frac{du}{dx}(0, \varepsilon) = 0, \quad \frac{du}{dx}(1, \varepsilon) = 0. \end{cases} \quad (5.1)$$

Here

$$\begin{aligned} \varphi_1(x) &= \varphi^{(-)} = x, \quad \varphi_2(x) = \frac{6}{5}, \quad \varphi_3(x) = \varphi^{(+)} = 2, \\ \bar{h}^{(-)}(x) &= -(x-2)^2(x-\frac{6}{5}), \quad \bar{h}^{(+)}(x) = -(x-2)^2(2-\frac{6}{5}), \\ \bar{f}_1^{(-)}(x) &= \frac{6}{5} - x, \quad \bar{f}_1^{(+)}(x) = -\frac{4}{5}. \end{aligned}$$

It is easy to verify that the conditions (A1) and (A2) are satisfied. Then $\bar{x}_0 = \frac{2}{5}$ is the root of the equation

$$I(\bar{x}_0) := \int_{x_0}^2 -(u-x_0)^2(u-\frac{6}{5})(u-2)^2 du = 0,$$

187 where $I'(\bar{x}_0) \neq 0$. Therefore, condition (A3) also holds.

188 From (4.4), we have

$$\bar{u}_1^{(-)} = \frac{1}{2-x}, \quad \bar{u}_1^{(+)} = \frac{-1}{2-x}, \quad (5.2)$$

We have the following equations for determining $Q_0^{(\pm)}(\tau)$ and their boundary conditions:

$$\begin{cases} \frac{dQ_0^{(-)}}{d\tau} = \left[2 \int_0^{Q_0^{(-)}} -(x_* + s - \frac{6}{5})(x_* + s - 2)^2 (s^2 + \frac{2\sqrt{\varepsilon}}{2-x}s) ds \right]^{\frac{1}{2}}, \\ Q_0^{(-)}(0) = \frac{6}{5} - x_*. \end{cases}$$

and

$$\begin{cases} \frac{dQ_0^{(+)}}{d\tau} = \left[2 \int_0^{Q_0^{(+)}} -(2 + s - x_*)^2 (2 + s - \frac{6}{5})(s^2 - \frac{2\sqrt{\varepsilon}}{2-x}s) ds \right]^{\frac{1}{2}}, \\ Q_0^{(+)}(0) = -\frac{4}{5}. \end{cases}$$

189 It is difficult to obtain the analytical solutions for these Cauchy problems, so we
190 provide only numerical simulation.

For $\Pi_0^{(\pm)}(\xi)$ we have

$$\begin{cases} \frac{d\Pi_0^{(-)}}{d\xi} = \frac{24}{5}\Pi_0^{(-)}, \\ \frac{d\Pi_0^{(-)}}{d\xi}(0) = -1, \quad \Pi_0^{(-)}(+\infty) = 0, \end{cases}$$

and

$$\begin{cases} \frac{d\Pi_0^{(+)}}{d\tilde{\xi}} = \frac{4}{5}\Pi_0^{(+)}, \\ \frac{d\Pi_0^{(+)}}{d\tilde{\xi}}(0) = 0, \quad \Pi_0^{(-)}(+\infty) = 0. \end{cases}$$

Here, we can solve these Cauchy problems analytically, and the solution is

$$\Pi_0^{(-)}(\xi) = \sqrt{5/24}\exp(-\sqrt{24/5}\xi), \quad \Pi_0^{(+)}(\tilde{\xi}) = 0.$$

191 At fig.1 you can see the results of numerical simulation for zero approximation
192 $U_0(x, \varepsilon)$ of the solution.

The whole solution can be get from theorem 4.1 and has the form

$$u(x, \varepsilon) = \begin{cases} x + Q_0^{(-)}(\tau) + \varepsilon^{\frac{3}{4}}\Pi_0^{(-)}(\xi) + O(\varepsilon^{\frac{1}{2}}), & 0 \leq x < (x_*)_{2n+5}, \\ 2 + Q_0^{(+)}(\tau) + O(\varepsilon^{\frac{1}{2}}), & (x_*)_{2n+5} \leq x \leq 1. \end{cases}$$




Figure 1. zero order approximation $U_0(x, \varepsilon)$ of problem (2.1)-(2.2)

193

194 6. Conclusion

195 In this paper, we discuss the singularly perturbed boundary value problem for a
196 degenerate equation with two double roots. The formal asymptotic solution is
197 constructed using the modified boundary layer function method. The existence
198 of multiple inner layers is established, and the existence of smooth solutions to the
199 problem is proven. Moreover, the obtained results are illustrated through numerical
200 simulation for particular right hand part of the equation. The theoretical results
201 can be extended to handle the contrast structure between repeated roots.

References

- 202
- 203 [1] N. N. Nefedov and E. I. Nikulin, Existence and asymptotic stability of periodic
204 two-dimensional contrast structures in the problem with weak linear advection,
205 *Mathematical Notes*, 2019, 106(5), 771–783.
- 206 [2] N. N. Nefedov, E. I. Nikulin and L. Recke, On the existence and asymptotic stability
207 of periodic contrast structures in quasilinear reaction–advection–diffusion
208 equations, *Russian Journal of Mathematical Physics*, 2019, 26(1), 55–69.
- 209 [3] N. N. Nefedov, L. Recke and K. R. Schneider, Existence and asymptotic stability
210 of periodic solutions with an internal layer of reaction–advection–diffusion
211 equations, *Journal of Mathematical Analysis and Applications*, 2013, 405(1),
- 212 [4] M. Ni, Y. Pang and M. A. Davydova, Contrast structures in problems for
213 a stationary equation of reaction–diffusion–advection type with discontinuous
214 nonlinearity, *Mathematical Notes*, 2018, 104(5), 118–127.
- 215 [5] A. B. Vasil’eva and V. F. Butuzov, *Asymptotic Methods in Singular Perturbation
216 Theory*, Vysshaya Shkola, Moscow, 1990.
- 217 [6] M. A. Davydova and S. A. Zakharova, Singularly perturbed stationary diffusion
218 model with a cubic nonlinearity, *Differential Equations*, 2020, 56(7), 819–830.
- 219 [7] V. F. Butuzov, On periodic solutions to singularly perturbed parabolic problems
220 in the cases of multiple roots of the degenerate equation, *Computational
221 Mathematics and Mathematical Physics*, 2011, 51(1), 40–50.
- 222 [8] V. F. Butuzov, On the special properties of the boundary layer in singularly
223 perturbed problems with multiple root of the degenerate equation, *Mathematical
224 Notes*, 2013, 94(1), 60–70.
- 225 [9] V. F. Butuzov, On the stability and the attraction domain of the stationary
226 solution of a singularly perturbed parabolic Equation with a multiple root of
227 the degenerate equation, *Partial Differential Equations*, 2015, 51(12), 1569–
228 1582.
- 229 [10] V. F. Butuzov, N. N. Nefedov, O. E. Omel’chenko and L. Recke, Partly dissipative
230 system with multizonal initial and boundary layers, *Journal of Physics:
231 Conference Series*, 2019, 1205, Article ID 012009, 7 pages.
- 232 [11] V. F. Butuzov, N. N. Nefedov, L. Recke and K. R. Schneider, On a singularly
233 perturbed initial value problem in the case of a double root of the degenerate
234 equation, *Nonlinear Analysis*, 2013, 83, 1–11.
- 235 [12] Q. Yang and M. K. Ni. Asymptotics of the solution to a stationary piecewise-
236 smooth reaction-diffusion equation with a multiple root of the degenerate equation,
237 *Science China Mathematics*, 2022, 65(2), 291-308.
- 238 [13] Q. Yang and M. K. Ni. Asymptotics of the solution to a stationary piecewise-
239 smooth reaction-diffusion-advection equation, *Chinese Annals of Mathematics
240 Ser.B*, 2023, 44(1), 91-98.
- 241 [14] Q. Yang and M. K. Ni. Multizonal internal layers in the singularly perturbed
242 equation with a discontinuous right-hand side, *Computational Mathematics
243 and Mathematical Physics*, 2021, 61(6), 953-963.
- 244 [15] Q. Yang and M. K. Ni. Asymptotics of the solution to a piecewise-smooth
245 quasilinear second-order differential equation, *Journal of Applied Analysis and
246 Computation*, 2022, 12(1), 256-269.

- 247 [16] Q. Yang, M. K. Ni. Asymptotics of a multizonal internal layer solution to
248 a piecewise-smooth singularly perturbed equation with a triple root of the
249 degenerate equation[J]. *Journal of Applied Analysis and Computation*, 2022,
250 12(6): 2441-2457.
- 251 [17] V. A. Beloshapko, V. F. Butuzov. Asymptotics of the solution to a singularly
252 perturbed elliptic problem with a three-zone boundary layer[J]. *Comput. Math.*
253 *Math. Phys.* 2016, 56(8):1414-1425.
- 254 [18] V. F. Butuzov, V. A. Beloshapko. Singularly perturbed elliptic dirichlet prob-
255 lem with a multiple root of the degenerate equation[J]. *Modelirovanie i analiz*
256 *informacionnyh sistem*. 2016, 23(5):515-528.
- 257 [19] V. F. Butuzov. Asymptotic behaviour and stability of solutions of a singularly
258 perturbed elliptic problem with a triple root of the degenerate equation[J].
259 *Izvestiya: Mathematics*. 2017, 81(3):481-504.
- 260 [20] Chen S L, Zhang R R, Asymptotic analysis for the singularly perturbed initial
261 value problem with degenerate equation having triple root[J]. *Qual. Theory*
262 *Dyn. Syst.* 2018, 17:91-101